

**FBI
NOTES**

GENERAL MATH

Federal Board Islamabad

Presented by:

**Urdu Books Whatsapp Group
STUDY GROUP**

**10TH
CLASS**

0333-8033313

راؤ ایاز

0343-7008883

پاکستان زندہ باد

0306-7163117

محمد سلمان سلیم

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

7

MAKING IT EASY

Solve:

1- If $P(x) = x^4 + 3x^2 - 5x + 9$, then find $P(x)$, for $x = 0, x = 1$.

Sol. $P(x) = x^4 + 3x^2 - 5x + 9$
 $P(0) = (0)^4 + 3(0)^2 - 5(0) + 9$
 $= 0 + 0 - 0 + 9$
 $= 9$

and $P(1) = (1)^4 + 3(1)^2 - 5(1) + 9$
 $= 1 + 3 - 5 + 9$
 $= 13 - 5$
 $= 8$

2- If $P(x) = 2x^3 + 2x^2 + x - 1$, then find $P(-2)$

Sol. $P(x) = 2x^3 + 2x^2 + x - 1$
Therefore, $P(-2) = 2(-2)^3 + 2(-2)^2 + (-2) - 1$
 $= 2(-8) + 2(4) - 2 - 1$
 $= -16 + 8 - 2 - 1$
 $= -8 - 1$
 $= -9$

3- If $P(y) = 3y^2 + \frac{y}{4} + 9$, then find $P(0)$.

Sol. $P(y) = 3y^2 + \frac{y}{4} + 9$
 $P(0) = 3(0)^2 + \frac{0}{4} + 9$
 $= 3(0) + 0 + 9$
 $= 9$

4- If $P(x) = 9x^3 - 2x^2 + 3x + 1$, then find $P(1)$ and $P(2)$.

Sol. $P(x) = 9x^3 - 2x^2 + 3x + 1$
 $P(1) = 9(1)^3 - 2(1)^2 + 3(1) + 1$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

8

$$\begin{aligned} &= 9(1) - 2(1) + 3 + 1 \\ &= 9 - 2 + 3 + 1 \\ &= 13 - 2 \\ &= 11 \end{aligned}$$

and $P(2) = 9(2)^3 - 2(2)^2 + 3(2) + 1$

$$\begin{aligned} &= 9(8) - 2(4) + 6 + 1 \\ &= 72 - 8 + 6 + 1 \\ &= 79 - 8 \\ &= 71 \end{aligned}$$

5- If $P(x) = \frac{x^2 - 5x + 6}{x + 1}$, then find $P(1)$ and $P(2)$.

Sol. $P(x) = \frac{x^2 - 5x + 6}{x + 1}$

$$\begin{aligned} P(1) &= \frac{(1)^2 - 5(1) + 6}{1 + 1} \\ &= \frac{1 - 5 + 6}{1 + 1} \\ &= \frac{2}{2} \\ &= 1 \end{aligned}$$

and $P(2) = \frac{(2)^2 - 5(2) + 6}{2 + 1}$

$$\begin{aligned} &= \frac{4 - 10 + 6}{3} \\ &= \frac{0}{3} \\ &= 0 \end{aligned}$$

6- If $P(r) = 2\pi r$, then find $P(r)$, for $r = 3$ and $\pi = \frac{22}{7}$.

Sol. $P(r) = 2\pi r$

By putting the values of r and π

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

9

$$\begin{aligned} P(r) &= 2 \times \frac{22}{7} \times 3 \\ &= \frac{132}{7} \\ &= 18.9 \text{ (approximately)} \end{aligned}$$

7- If $P(r) = 4\pi r^2$, then find $P(r)$, for $r = 8$ and $\pi = \frac{22}{7}$.

Sol. $P(r) = 4\pi r^2$

By putting the values of r and π

$$\begin{aligned} P(r) &= 4 \times \frac{22}{7} \times (8)^2 \\ &= 4 \times \frac{22}{7} \times 64 \\ &= \frac{5632}{7} \\ &= 804.57 \end{aligned}$$

8- If $P(y) = y^4 + \frac{3y^3}{2} - y^2 + 1$, then find $P(y)$, for $y = 2$ and $y = -2$.

Sol. $P(y) = y^4 + \frac{3y^3}{2} - y^2 + 1$

for $y = 2$

$$\begin{aligned} P(2) &= (2)^4 + \frac{3(2)^3}{2} - (2)^2 + 1 \\ &= 16 + \frac{3 \times 8}{2} - 4 + 1 \\ &= 16 + 12 - 4 + 1 \\ &= 25 \end{aligned}$$

$$P(-2) = (-2)^4 + \frac{3(-2)^3}{2} - (-2)^2 + 1 \quad \text{for } y = -2$$

ختم نبوت ﷺ زندہ باد

عظمت صحابہ زندہ باد

السلام علیکم ورحمۃ اللہ وبرکاتہ:

معزز ممبران: آپ کا وٹس ایپ گروپ ایڈمن "اردو بکس" آپ سے مخاطب ہے۔

آپ تمام ممبران سے گزارش ہے کہ:

- ❖ گروپ میں صرف PDF کتب پوسٹ کی جاتی ہیں لہذا کتب کے متعلق اپنے کمنٹس / ریویوز ضرور دیں۔ گروپ میں بغیر ایڈمن کی اجازت کے کسی بھی قسم کی (اسلامی و غیر اسلامی، اخلاقی، تحریری) پوسٹ کرنا سختی سے منع ہے۔
- ❖ گروپ میں معزز، پڑھے لکھے، سلجھے ہوئے ممبرز موجود ہیں اخلاقیات کی پابندی کریں اور گروپ رولز کو فالو کریں بصورت دیگر معزز ممبرز کی بہتری کی خاطر ریموو کر دیا جائے گا۔
- ❖ کوئی بھی ممبر کسی بھی ممبر کو انباکس میں میسج، مس کال، کال نہیں کرے گا۔ رپورٹ پر فوری ریموو کر کے کارروائی عمل میں لائے جائے گی۔
- ❖ ہمارے کسی بھی گروپ میں سیاسی و فرقہ واریت کی بحث کی قطعاً کوئی گنجائش نہیں ہے۔
- ❖ اگر کسی کو بھی گروپ کے متعلق کسی قسم کی شکایت یا تجویز کی صورت میں ایڈمن سے رابطہ کیجئے۔
- ❖ سب سے اہم بات:

گروپ میں کسی بھی قادیانی، مرزائی، احمدی، گستاخ رسول، گستاخ امہات المؤمنین، گستاخ صحابہ و خلفائے راشدین حضرت ابو بکر

صدیق، حضرت عمر فاروق، حضرت عثمان غنی، حضرت علی المرتضیٰ، حضرت حسنین کریمین رضوان اللہ تعالیٰ اجمعین، گستاخ اہلبیت یا

ایسے غیر مسلم جو اسلام اور پاکستان کے خلاف پراپیگنڈا میں مصروف ہیں یا ان کے روحانی و ذہنی سپورٹرز کے لئے کوئی گنجائش نہیں

ہے لہذا ایسے اشخاص بالکل بھی گروپ جو ان کرنے کی زحمت نہ کریں۔ معلوم ہونے پر فوراً ریموو کر دیا جائے گا۔

❖ تمام کتب انٹرنیٹ سے تلاش / ڈاؤنلوڈ کر کے فری آف کاسٹ وٹس ایپ گروپ میں شیئر کی جاتی ہیں۔ جو کتاب نہیں ملتی اس کے لئے معذرت کر

لی جاتی ہے۔ جس میں محنت بھی صرف ہوتی ہے لیکن ہمیں آپ سے صرف دعاؤں کی درخواست ہے۔

❖ عمران سیریز کے شوقین کیلئے علیحدہ سے عمران سیریز گروپ موجود ہے۔

❖ لیڈرز کے لئے الگ گروپ کی سہولت موجود ہے جس کے لئے ویریفیکیشن ضروری ہے۔

❖ اردو کتب / عمران سیریز یا سٹیڈی گروپ میں ایڈ ہونے کے لئے ایڈمن سے وٹس ایپ پر بذریعہ میسج رابطہ کریں اور جواب کا انتظار فرمائیں۔ برائے

مہربانی اخلاقیات کا خیال رکھتے ہوئے موبائل پر کال یا ایم ایس کرنے کی کوشش ہرگز نہ کریں۔ ورنہ گروپس سے توریوو کیا ہی جائے گا بلاک بھی کیا

جائے گا۔

نوٹ: ہمارے کسی گروپ کی کوئی فیس نہیں ہے۔ سب فی سبیل اللہ ہے

0333-8033313

0343-7008883

0306-7163117

راؤ ایاز

پاکستان زندہ باد

محمد سلمان سلیم

پاکستان پائمنڈہ باد

پاکستان زندہ باد

اللہ تبارک تعالیٰ ہم سب کا حامی و ناصر ہو

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

10

$$= 16 + \frac{3(-8)}{2} - 4 + 1$$

$$= 16 - 12 - 4 + 1$$

$$= 1$$

Reduce the given rational expressions to lowest terms.

Q.9 $\frac{8x^2y^2}{12x^4y}$

Sol. $= \frac{2y^{2-1}}{3x^{4-2}}$

$$= \frac{2y}{3x^2}$$

Q.10 $\frac{25a^3b^2}{14a^2b^4}$

Sol. $= \frac{25a^{3-2}}{14b^{4-2}}$

$$= \frac{25a}{14b^2}$$

Q.11 $\frac{16a^6b^7}{12a^3b^3 + 20a^5b^4}$

Sol. $= \frac{16a^6b^7}{4a^3b^4(3b + 5a^2)}$

$$= \frac{4a^{6-3}b^{7-4}}{(5a^2 + 3b)}$$

$$= \frac{4a^3b^3}{5a^2 + 3b}$$

Q.12 $\frac{18m^3x^3}{27m^4x^5 - 36m^6x^6}$

Sol. $= \frac{18m^3x^3}{9m^4x^6(3x^2 - 4m^2)}$

$$= \frac{2m^{3-4}}{x^{6-3}(3x^2 - 4m^2)}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

11

$$= \frac{2m}{x^3(3x^2 - 4m^2)}$$

$$= \frac{2m}{3x^3 - 4m^2x^3}$$

Q.13 $\frac{5c - 5d}{c^2 - d^2}$

Sol. $= \frac{5(c - d)}{(c - d)(c + d)}$

$$= \frac{5}{(c + d)}$$

Q.14 $\frac{x^2 - y^2}{3y - 3x}$

Sol. $= \frac{(x - y)(x + y)}{3(y - x)}$

$$= \frac{(x - y)(x + y)}{-3(x - y)}$$

$$= \frac{x + y}{-3}$$

Simplify:

Q.15 $\frac{x}{x - y} + \frac{x^2}{x^2 + y^2}$

Sol. $= \frac{x(x^2 + y^2) + x^2(x - y)}{(x - y)(x^2 + y^2)}$

$$= \frac{x^3 + xy^2 + x^3 - x^2y}{x^3 + xy^2 - x^2y - y^3}$$

$$= \frac{2x^3 - x^2y + xy^2}{x^3 - x^2y + xy^2 - y^3}$$

Q.16 $\frac{x^2 + 2x}{x^2 + x - 2} + \frac{3x}{x + 1}$

Sol. $= \frac{(x^2 + 2x)(x + 1) + 3x(x^2 + x - 2)}{(x^2 + x - 2)(x + 1)}$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

12

$$= \frac{x^3 + x^3 + 2x^2 + 2x + 3x^3 + 3x^2 - 6x}{x^3 + x^2 - 2x + x^2 + x - 2}$$

$$= \frac{4x^3 + 6x^2 - 4x}{x^3 + 2x^2 - x - 2}$$

Q.17 $\frac{x+2}{x^2+3x+2} - \frac{x-5}{x^2-x-6}$

Sol. $= \frac{x+2}{x^2+x+2x+2} - \frac{x-5}{x^2-x-6}$

$$= \frac{x+2}{x(x+1)+2(x+1)} - \frac{x-5}{x^2-x-6}$$

$$= \frac{x+2}{(x+2)(x+1)} - \frac{x-5}{x^2-x-6}$$

$$= \frac{1}{x+1} - \frac{x-5}{x^2-x-6}$$

$$= \frac{1(x^2-x-6) - (x+1)(x-5)}{(x+1)(x^2-x-6)}$$

$$= \frac{x^2-x-6 - (x^2-5x+x-5)}{(x+1)(x^2-x-6)}$$

$$= \frac{x^2-x-6-x^2+5x-x+5}{x^3-x^2-6x+x^2-x-6}$$

$$= \frac{3x-1}{x^3-7x-6}$$

Q.18 $\frac{8x^2+18y^2}{4x^2-9y^2} - \frac{2x+3y}{2x-3y}$

Sol. $= \frac{8x^2+18y^2}{(2x)^2-(3y)^2} - \frac{2x+3y}{2x-3y}$

$$= \frac{8x^2+18y^2}{(2x+3y)(2x-3y)} - \frac{2x+3y}{2x-3y}$$

$$= \frac{8x^2+18y^2 - (2x+3y)(2x+3y)}{(2x+3y)(2x-3y)}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

13

$$= \frac{8x^2 + 18y^2 - (4x^2 + 12xy + 9y^2)}{(2x + 3y)(2x - 3y)}$$

$$= \frac{8x^2 + 18y^2 - 4x^2 - 12xy - 9y^2}{(2x + 3y)(2x - 3y)}$$

$$= \frac{4x^2 - 12xy + 9y^2}{(2x + 3y)(2x - 3y)}$$

$$= \frac{(2x)^2 - 2(2x)(3y) + (3y)^2}{(2x + 3y)(2x - 3y)}$$

$$= \frac{(2x - 3y)^2}{(2x + 3y)(2x - 3y)}$$

$$= \frac{(2x - 3y)(2x - 3y)}{(2x + 3y)(2x - 3y)}$$

$$= \frac{2x - 3y}{2x + 3y}$$

Q.19 $\frac{x}{x^2 + xy} - \frac{y}{x^2 - y^2}$

Sol. $= \frac{x}{x(x + y)} - \frac{y}{(x + y)(x - y)}$

$$= \frac{1}{x + y} - \frac{y}{(x + y)(x - y)}$$

$$= \frac{(x - y) - y}{(x + y)(x - y)}$$

$$= \frac{x - y - y}{(x + y)(x - y)}$$

$$= \frac{x - 2y}{x^2 - y^2}$$

Q.20 $\frac{x + y}{xy + y^2} - \frac{x}{x^2 - xy}$

Sol. $= \frac{x + y}{y(x + y)} - \frac{x}{x(x - y)}$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

14

$$\begin{aligned} &= \frac{1}{y} - \frac{1}{x-y} \\ &= \frac{1(x-y) - 1(y)}{(y)(x-y)} \\ &= \frac{x-y-y}{y(x-y)} \\ &= \frac{x-2y}{xy-y^2} \end{aligned}$$

Q.21 $\frac{(x+1)^2}{x^2-1} - \frac{x^2+1}{x^2+1}$

Sol. $\begin{aligned} &= \frac{(x+1)(x+1)}{(x+1)(x-1)} - \frac{x^2+1}{x^2+1} \\ &= \frac{x+1}{x-1} - 1 \\ &= \frac{x+1-1(x-1)}{x-1} \\ &= \frac{x+1-x+1}{x-1} \\ &= \frac{2}{x-1} \end{aligned}$

Q.22 $\frac{5x}{x-9} + \frac{x^2-2x+1}{x^2-12x+27} - \frac{6x}{x-3}$

Sol. $\begin{aligned} &= \frac{5x}{x-9} + \frac{x^2-2x+1}{x^2-9x-3x+27} - \frac{6x}{x-3} \\ &= \frac{5x}{x-9} + \frac{x^2-2x+1}{x(x-9)-3(x-9)} - \frac{6x}{x-3} \\ &= \frac{5x}{x-9} + \frac{x^2-2x+1}{(x-9)(x-3)} - \frac{6x}{x-3} \\ &= \frac{5x(x-3) + x^2-2x+1 - 6x(x-9)}{(x-9)(x-3)} \end{aligned}$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

15

$$= \frac{5x^2 - 15x + x^2 - 2x + 1 - 6x^2 + 54x}{(x-9)(x-3)}$$

$$= \frac{37x+1}{x^2-12x+27}$$

Q.23 $\frac{x^2-4x+4}{x^2-4} + \frac{x}{x-2}$

Sol. $= \frac{x^2-2x-2x+4}{(x)^2-(2)^2} \times \frac{x-2}{x}$
 $= \frac{x(x-2)-2(x-2)}{(x+2)(x-2)} \times \frac{x-2}{x}$
 $= \frac{(x-2)(x-2)}{x+2} \times \frac{1}{x}$
 $= \frac{x^2-4x+4}{x^2+2x}$

Q.24 $\frac{x^2-36}{x^2-1} + \frac{x-6}{1-x}$

Sol. $= \frac{(x)^2-(6)^2}{(x)^2-(1)^2} \times \frac{1-x}{x-6}$
 $= \frac{(x-6)(x+6)}{(x-1)(x+1)} \times \frac{-(x-1)}{x-6}$
 $= \frac{x+6}{x+1} \times (-1)$
 $= -\frac{x+6}{x+1}$

Q.25 $\frac{x^2-5x}{x-1} + \frac{x^2-25}{x^2+x+20}$

Sol. $= \frac{x(x-5)}{x-1} \times \frac{x^2+x+20}{x^2-25}$
 $= \frac{x(x-5)}{x-1} \times \frac{x^2+x+20}{(x)^2-(5)^2}$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

16

$$= \frac{x(x-5)}{x-1} \times \frac{x^2+x+20}{(x-5)(x+5)}$$

$$= \frac{x(x^2+x+20)}{(x-1)(x+5)}$$

$$= \frac{x^3+x^2+20x}{x^2+4x-5}$$

Q.26 $\frac{2x^2-5x-12}{4x^2+4x-3} + \frac{2x^2-7x-4}{6x^2+5x-4}$

Sol. $= \frac{2x^2-8x+3x-12}{4x^2+6x-2x-3} \times \frac{6x^2+5x-4}{2x^2-7x-4}$
 $= \frac{2x(x-4)+3(x-4)}{2x(2x+3)-1(2x+3)} \times \frac{6x^2+8x-3x-4}{2x^2-8x+x-4}$
 $= \frac{(x-4)(2x+3)}{(2x+3)(2x-1)} \times \frac{2x(3x+4)-1(3x+4)}{2x(x-4)+1(x-4)}$
 $= \frac{x-4}{2x-1} \times \frac{(3x+4)(2x-1)}{(x-4)(2x+1)}$
 $= \frac{3x+4}{2x+1}$

Q.27 $\frac{x(2x-1)^2}{2x^2-1} + \frac{4x^2-1}{4x^2+4x+1}$

Sol. $= \frac{x(2x-1)(2x-1)}{2x^2-1} \times \frac{4x^2+4x+1}{4x^2-1}$
 $= \frac{x(2x-1)(2x-1)}{(2x^2-1)} \times \frac{4x^2+2x+2x+1}{(2x)^2-(1)^2}$
 $= \frac{x(2x-1)(2x-1)}{(2x^2-1)} \times \frac{2x(2x+1)+1(2x+1)}{(2x+1)(2x-1)}$
 $= \frac{x(2x-1)(2x-1)}{(2x^2-1)} \times \frac{(2x+1)(2x+1)}{(2x+1)(2x-1)}$
 $= \frac{x(2x-1)(2x+1)}{2x^2-1}$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

17

$$= \frac{x(4x^2 - 1)}{2x^2 - 1}$$

$$= \frac{4x^3 - x}{2x^2 - 1}$$

Q.28 $\frac{x^2 + x}{x^2 - 1} \times \frac{x + 1}{x^3 + 1}$

Sol. $= \frac{x(x+1)}{(x+1)(x-1)} \times \frac{(x+1)}{(x+1)(x^2 - x + 1)}$

$$= \frac{x}{(x-1)(x^2 - x + 1)}$$

$$= \frac{x}{x^3 - x^2 + x - x^2 + x - 1}$$

$$= \frac{x}{x^3 - 2x^2 + 2x - 1}$$

Q.29 $\frac{x^2 - 9}{x^2 - 6x + 9} \times \frac{x}{3x + 9}$

Sol. $= \frac{(x)^2 - (3)^2}{x^2 - 3x - 3x + 9} \times \frac{x}{3(x+3)}$

$$= \frac{(x+3)(x-3)}{x(x-3) - 3(x-3)} \times \frac{x}{3(x+3)}$$

$$= \frac{(x+3)(x-3)}{(x-3)(x-3)} \times \frac{x}{3(x+3)}$$

$$= \frac{x}{3(x-3)}$$

$$= \frac{x}{3x - 9}$$

Q.30 $\frac{x+5}{x^2+6x} \times \frac{x^3+6x^2}{x+5}$

Sol. $= \frac{(x+5)}{x(x+6)} \times \frac{x^2(x+6)}{(x+5)}$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

18

$$= \frac{x^2}{x}$$

$$= x^{2-1}$$

$$= x$$

Q.31 $\frac{x^2 - 2x + 1}{x^2 - 1} \times \frac{x+1}{x-1}$

Sol. $= \frac{x^2 - x - x + 1}{(x)^2 - (1)^2} \times \frac{x+1}{x-1}$
 $= \frac{x(x-1) - 1(x-1)}{(x+1)(x-1)} \times \frac{(x+1)}{(x-1)}$
 $= \frac{(x-1)(x-1)}{(x+1)(x-1)} \times \frac{(x+1)}{(x-1)}$
 $= 1$

Q.32 $\frac{x^2 + 4x + 3}{x+3} \times \frac{x^2 - 2x + 1}{x^2 - 1}$

Sol. $= \frac{x^2 + x + 3x + 3}{x+3} \times \frac{x^2 - x - x + 1}{(x)^2 - (1)^2}$
 $= \frac{x(x+1) + 3(x+1)}{(x+3)} \times \frac{x(x-1) - 1(x-1)}{(x-1)(x+1)}$
 $= \frac{(x+1)(x+3)}{(x+3)} \times \frac{(x-1)(x-1)}{(x-1)(x+1)}$
 $= x - 1$

Formulae

(i) $(a + b)^2 = a^2 + 2ab + b^2$

(ii) $(a - b)^2 = a^2 - 2ab + b^2$

(iii) $(a + b)^2 + (a - b)^2 = 2(a^2 + b^2)$

(iv) $(a + b)^2 - (a - b)^2 = 4ab$

(v) $(a + b + c)^2 = a^2 + b^2 + c^2 + 2ab + 2bc + 2ca$
 $= a^2 + b^2 + c^2 + 2(ab + bc + ca)$

(vi) $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

19

$$\begin{aligned} &= a^3 + b^3 + 3ab(a + b) \\ \text{(vii)} \quad (a - b)^3 &= a^3 - 3a^2b + 3ab^2 - b^3 \\ &= a^3 - b^3 - 3ab(a - b) \\ \text{(viii)} \quad a^3 + b^3 &= (a + b)(a^2 - ab + b^2) \\ \text{(ix)} \quad a^3 - b^3 &= (a - b)(a^2 + ab + b^2) \end{aligned}$$

EXERCISE-10

Solve the following questions using formulas.

Q.1 $(x + 2y)^2 + (x - 2y)^2$

Sol. $= 2[(x)^2 + (2y)^2]$ [Formula: $(a + b)^2 + (a - b)^2 = 2(a^2 + b^2)$]
 $= 2[x^2 + 4y^2]$
 $= 2x^2 + 8y^2$

Q.2 $(5x + 3y)^2 + (5x - 3y)^2$

Sol. $= 2[(5x)^2 + (3y)^2]$ [Formula: $(a + b)^2 + (a - b)^2 = 2(a^2 + b^2)$]
 $= 2[25x^2 + 9y^2]$
 $= 50x^2 + 18y^2$

Q.3 $(3l + 2m)^2 - (3l - 2m)^2$

Sol. $= 4(3l)(2m)$ [Formula: $(a + b)^2 - (a - b)^2 = 4ab$]
 $= 24lm$

Q.4 $(l + m)(l - m)(l^2 + m^2)(l^4 + m^4)$

Sol. $= [(l + m)(l - m)](l^2 + m^2)(l^4 + m^4)$
 [Formula: $(a + b)(a - b) = a^2 - b^2$]
 $= (l^2 - m^2)(l^2 + m^2)(l^4 + m^4)$
 $= [(l^2)^2 - (m^2)^2](l^4 + m^4)$
 $= (l^4 - m^4)(l^4 + m^4)$
 $= (l^4)^2 - (m^4)^2$
 $= l^8 - m^8$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

20

Q.5 $\left(ab - \frac{1}{ab}\right)^3$

Sol. [Formula: $(a - b)^3 = a^3 - 3a^2b + 3ab^2 - b^3$]

$$= (ab)^3 - 3(ab)^2 \left(\frac{1}{ab}\right) + 3ab \left(\frac{1}{ab}\right)^2 - \left(\frac{1}{ab}\right)^3$$

$$= a^3b^3 - 3ab + \frac{3}{ab} - \frac{1}{a^3b^3}$$

Q.6 $(2x + 3y + 2)^3$

Sol. [(a + b + c)² = a² + b² + c² + 2ab + 2bc + 2ca]

$$= (2x)^2 + (3y)^2 + (2)^2 + 2(2x)(3y) + 2(3y)(2) + 2(2)(2x)$$

$$= 4x^2 + 9y^2 + 4 + 12xy + 12y + 8x$$

Q.7 $(2p + q)^3$

Sol. [Formula: $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$]

$$= (2p)^3 + 3(2p)^2(q) + 3(2p)(q)^2 + (q)^3$$

$$= 8p^3 + 3(4p^2)(q) + 6pq^2 + q^3$$

$$= 8p^3 + 12p^2q + 6pq^2 + q^3$$

Q.8 $(3p + q + r)^2$

Sol. [Formula: $(a + b + c)^2 = a^2 + b^2 + c^2 + 2ab + 2bc + 2ca$]

$$= (3p)^2 + (q)^2 + (r)^2 + 2(3p)(q) + 2(q)(r) + 2(r)(3p)$$

$$= 9p^2 + q^2 + r^2 + 6pq + 2qr + 6rp$$

Q.9 $(2x + 3y)^3$

Sol. [Formula: $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$]

$$= (2x)^3 + 3(2x)^2(3y) + 3(2x)(3y)^2 + (3y)^3$$

$$= 8x^3 + 3(4x^2)(3y) + 3(2x)(9y^2) + 9y^3$$

$$= 8x^3 + 36x^2y + 54xy^2 + 9y^3$$

Q.10 $(x + y)^3 - 1$

Sol. $= (x + y)^3 - (1)^3$
 [Formula: $a^3 - b^3 = (a - b)(a^2 + ab + b^2)$]

$$= (x + y - 1)((x + y)^2 + (x + y)(1) + (1)^2)$$

$$= (x + y - 1)(x^2 + y^2 + 2xy + x + y + 1)$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

21

Q.11 $(x - y)^3 + 64$

Sol. $= (x - y)^3 + (4)^3$

$$\begin{aligned} & \text{[Formula: } a^3 + b^3 = (a + b)(a^2 - ab + b^2)] \\ & = (x - y + 4)[(x - y)^2 - (x - y)(4) + (4)^2] \\ & = (x - y + 4)(x^2 + y^2 + 2xy - 4x + 4y + 16) \end{aligned}$$

Q.12 $8x^3 + 27y^3$

Sol. $= (2x)^3 + (3y)^3$

$$\begin{aligned} & \text{[Formula: } a^3 + b^3 = (a + b)(a^2 - ab + b^2)] \\ & = (2x + 3y)[(2x)^2 - (2x)(3y) + (3y)^2] \\ & = (2x + 3y)(4x^2 - 6xy + 9y^2) \end{aligned}$$

Q.13 $x^6 - 729y^6$

Sol. $= (x^3)^2 - (27y^3)^2$ [Formula: $a^2 - b^2 = (a + b)(a - b)$]
 $= (x^3 + 27y^3)(x^3 - 27y^3)$
 $= [(x)^3 + (3y)^3][(x)^3 - (3y)^3]$

(Formulas)

$$\begin{aligned} a^3 + b^3 &= (a + b)(a^2 - ab + b^2) \\ a^3 - b^3 &= (a - b)(a^2 + ab + b^2) \\ &= (x + 3y)(x^2 - 3xy + 9y^2)(x - 3y)(x^2 + 3xy + 9y^2) \\ &= (x + 3y)(x - 3y)(x^2 - 3xy + 9y^2)(x^2 + 3xy + 9y^2) \end{aligned}$$

Q.14 $64a^6 - b^6$

Sol. $= (8a^3)^2 - (b^3)^2$
 $= (8a^3 - b^3)(8a^3 + b^3)$ [Formula: $a^2 - b^2 = (a - b)(a + b)$]
 $= [(2a)^3 - (b)^3][(2a)^3 + (b)^3]$
 $a^3 + b^3 = (a + b)(a^2 - ab + b^2)$

(Formulas)

$$\begin{aligned} a^3 - b^3 &= (a - b)(a^2 + ab + b^2) \\ &= (2a - b)(4a^2 + 2ab + b^2)(2a + b)(4a^2 - 2ab + b^2) \\ &= (2a - b)(2a + b)(4a^2 + 2ab + b^2)(4a^2 - 2ab + b^2) \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

22

Q.15 Find the value of $a^3 - b^3$ when $a - b = 4$ and $ab = -5$.

Sol. $a - b = 4$
 $ab = -5$
 $a^3 - b^3 = ?$
 $a - b = 4$
 $(a - b)^3 = (4)^3$
 $a^3 - b^3 - 3(-5)(4) = 64$
 $a^3 - b^3 + 60 = 64$
 $a^3 - b^3 = 64 - 60$
 $a^3 - b^3 = 4$

Taking cube

Q.16 Show that $\left(z + \frac{1}{z}\right)^2 - \left(z - \frac{1}{z}\right)^2 = 4$

L.H.S. $= \left(z + \frac{1}{z}\right)^2 - \left(z - \frac{1}{z}\right)^2$
 $= \left(z^2 + 2(z)\left(\frac{1}{z}\right) + \frac{1}{z^2}\right) - \left(z^2 - 2(z)\left(\frac{1}{z}\right) + \frac{1}{z^2}\right)$
 $= \left(z^2 + 2 + \frac{1}{z^2}\right) - \left(z^2 - 2 + \frac{1}{z^2}\right)$
 $= z^2 + 2 + \frac{1}{z^2} - z^2 + 2 - \frac{1}{z^2}$
 $= 4 = \text{R.H.S.}$

Q.17 Find the value of $a^2 + b^2$ and ab when $a + b = 5$ and $a - b = 3$.

Sol. $a + b = 5$
 $a - b = 3$
 $a^2 + b^2 = ?$
 $ab = ?$

Formula.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

23

$$(a + b)^2 + (a - b)^2 = 2(a^2 + b^2)$$

$$(5)^2 + (3)^2 = 2(a^2 + b^2)$$

$$25 + 9 = 2(a^2 + b^2)$$

$$34 = 2(a^2 + b^2)$$

$$\text{Thus, } a^2 + b^2 = \frac{34}{2} = 17$$

$$\text{Formula: } (a + b)^2 - (a - b)^2 = 4ab$$

$$(5)^2 - (3)^2 = 4ab$$

$$25 - 9 = 4ab$$

$$16 = 4ab$$

$$ab = \frac{16}{4}$$

$$ab = 4$$

Q.18 Find the value of $a^2 + b^2 + c^2$ if $ab + bc + ca = 11$ and

$$a + b + c = 6$$

Sol. $a + b + c = 6$

$$ab + bc + ca = 11$$

$$a^2 + b^2 + c^2 = ?$$

$$a + b + c = 6$$

$$(a + b + c)^2 = (6)^2 \quad \text{Taking square root}$$

$$a^2 + b^2 + c^2 + 2(ab + bc + ca) = 36$$

$$\text{By putting values } ab + bc + ca = 11$$

$$a^2 + b^2 + c^2 + 2(11) = 36$$

$$a^2 + b^2 + c^2 + 22 = 36$$

$$a^2 + b^2 + c^2 = 36 - 22$$

$$a^2 + b^2 + c^2 = 14$$

Q.19 Find the value of $x^3 + y^3$ if $xy = 10$ and $x + y = 7$.

Sol. $x + y = 7$

$$xy = 10$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

24

$$x^3 + y^3 = ?$$

$$x + y = 7$$

$$(x + y)^3 = (7)^3 \quad \text{Taking cube}$$

$$x^3 + y^3 + 3xy(x + y) = 343$$

$$x^3 + y^3 + 3(10)(7) = 343$$

$$x^3 + y^3 + 210 = 343$$

$$x^3 + y^3 = 343 - 210$$

$$x^3 + y^3 = 133$$

Q.20 Find the value of $(x - y)^2$ if $x^2 + y^2 = 86$ and $xy = -16$.

Sol. $xy = -16 \quad (i)$

$$x^2 + y^2 = 86 \quad (ii)$$

$$(x - y)^2 = ?$$

$$(x - y)^2 = x^2 + y^2 - 2xy$$

$$= (x^2 + y^2) - 2(xy)$$

$$= 86 - 2(-16) \quad \text{From (i) and (ii)}$$

$$= 86 + 32$$

$$= 118$$

Q.21 Find the value of $ab + bc + ca$ when the values of

$$a^2 + b^2 + c^2 = 81, a + b + c = 11$$

Sol. $a + b + c = 11 \dots\dots\dots(i)$

$$a^2 + b^2 + c^2 = 81 \dots\dots\dots(ii)$$

$$ab + bc + ca = ?$$

$$a + b + c = 11$$

$$(a + b + c)^2 = (11)^2 \quad \text{Taking square root}$$

$$a^2 + b^2 + c^2 + 2(ab + bc + ca) = 121$$

$$81 + 2(ab + bc + ca) = 121 \quad \text{from (ii)}$$

$$2(ab + bc + ca) = 121 - 81$$

$$2(ab + bc + ca) = 40$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

25

$$ab + bc + ca = \frac{40}{2}$$

$$ab + bc + ca = 20$$

Q.22 Find the value of $(a + b + c)^2$ when the values of $a^2 + b^2 + c^2 = 32$ and $ab + bc + ca = 7$.

Sol. $ab + bc + ca = 7$ (i)

$$a^2 + b^2 + c^2 = 32$$
 (ii)

$$(a + b + c)^2 = ?$$

$$\begin{aligned} (a + b + c)^2 &= a^2 + b^2 + c^2 + 2(ab + bc + ca) \\ &= 32 + 2(7) \quad \text{From (i) and (ii)} \\ &= 32 + 14 \end{aligned}$$

$$(a + b + c)^2 = 46$$

Laws of Radicals

$$(i) (\sqrt[n]{a})^n = a \quad (ii) \sqrt[n]{ab} = \sqrt[n]{a} \sqrt[n]{b}$$

$$(iii) \sqrt[n]{\frac{a}{b}} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}} \quad (iv) (\sqrt[n]{a})^m = \sqrt[n]{a^m}$$

Q.1 Remove the radical sign from the denominator:

(i) $\frac{1}{\sqrt{5}}$

Sol.

$$= \frac{1}{\sqrt{5}} \times \frac{\sqrt{5}}{\sqrt{5}}$$

$$= \frac{\sqrt{5}}{(\sqrt{5})^1}$$

$$= \frac{\sqrt{5}}{5}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

26

(ii) $\frac{2}{\sqrt{2}} \cdot \frac{7}{\sqrt{3}}$

Sol.

$$\begin{aligned} &= \frac{2 \times 7}{\sqrt{2} \times \sqrt{3}} \\ &= \frac{14}{\sqrt{6}} \\ &= \frac{14}{\sqrt{6}} \times \frac{\sqrt{6}}{\sqrt{6}} \\ &= \frac{14 \times \sqrt{6}}{6} = \frac{7\sqrt{6}}{3} \end{aligned}$$

(iii) $\frac{\sqrt{6}}{\sqrt{7}}$

Sol.

$$\begin{aligned} &= \frac{\sqrt{6}}{\sqrt{7}} \times \frac{\sqrt{7}}{\sqrt{7}} \\ &= \frac{\sqrt{6 \times 7}}{\sqrt{7 \times 7}} \\ &= \frac{\sqrt{42}}{7} \end{aligned}$$

Q.2 Simplify these expressions:

(i) $\sqrt{2} + \sqrt{8}$

Sol.

$$\begin{aligned} &= \sqrt{2} + \sqrt{2 \times 2 \times 2} \\ &= \sqrt{2} + 2\sqrt{2} \\ &= \sqrt{2}(1 + 2) \\ &= 3\sqrt{2} \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

27

(ii) $4\sqrt{50} + \sqrt{200} + \sqrt{50}$

Sol.

$$\begin{aligned} &= 4\sqrt{5 \times 5 \times 2} + \sqrt{5 \times 5 \times 2 \times 2 \times 2} + \sqrt{5 \times 5 \times 2} \\ &= 4 \times 5\sqrt{2} + 5 \times 2\sqrt{2} + 5\sqrt{2} \\ &= 20\sqrt{2} + 10\sqrt{2} + 5\sqrt{2} \\ &= \sqrt{2}(20 + 10 + 5) \\ &= \sqrt{2}(35) \\ &= 35\sqrt{2} \end{aligned}$$

(iii) $(\sqrt{12} - \sqrt{2})(\sqrt{20} - 3\sqrt{2})$

Sol.

$$\begin{aligned} &= \sqrt{12} \times \sqrt{20} - \sqrt{12} \times 3\sqrt{2} - \sqrt{2} \times \sqrt{20} + \sqrt{2} \times 3\sqrt{2} \\ &= \sqrt{2 \times 2 \times 3} \times \sqrt{2 \times 2 \times 5} - \sqrt{2 \times 2 \times 3} \times 3\sqrt{2} \\ &\quad - \sqrt{2} \times \sqrt{2 \times 2 \times 5} + 3\sqrt{2 \times 2} \\ &= 2\sqrt{3} \times 2\sqrt{5} - 2\sqrt{3} \times 3\sqrt{2} - \sqrt{2} \times 2\sqrt{5} + 3 \times 2 \\ &= 4\sqrt{3 \times 5} - 6\sqrt{3 \times 2} - 2\sqrt{2 \times 5} + 6 \\ &= 4\sqrt{15} - 6\sqrt{6} - 2\sqrt{10} + 6 \end{aligned}$$

(iv) $(6 + \sqrt{2})(5 - \sqrt{5})$

Sol.

$$\begin{aligned} &= 6(5 - \sqrt{5}) + \sqrt{2}(5 - \sqrt{5}) \\ &= 30 - 6\sqrt{5} + 5\sqrt{2} - \sqrt{2} \times 5 \\ &= 30 - 6\sqrt{5} + 5\sqrt{2} - \sqrt{10} \end{aligned}$$

(v) $(\sqrt{3} - 2)(5 - \sqrt{5})$

Sol.

$$\begin{aligned} &= \sqrt{3}(5 - \sqrt{5}) - 2(5 - \sqrt{5}) \\ &= 5\sqrt{3} - \sqrt{3} \times 5 - 10 + 2\sqrt{5} \\ &= 5\sqrt{3} - \sqrt{15} - 10 + 2\sqrt{5} \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

=====

Pilot Super One "General Math" 10th

28

(vi) $(7 + \sqrt{3})(5 + \sqrt{2})$

Sol.

$$\begin{aligned} &= 7(5 + \sqrt{2}) + \sqrt{3}(5 + \sqrt{2}) \\ &= 35 + 7\sqrt{2} + 5\sqrt{3} + \sqrt{3 \times 2} \\ &= 35 + 7\sqrt{2} + 5\sqrt{3} + \sqrt{6} \end{aligned}$$

Q.3 Rationalize the denominators of the following:

(i) $\frac{1}{\sqrt{3} + 2}$

Sol.

$$\begin{aligned} &= \frac{1}{\sqrt{3} + 2} \times \frac{\sqrt{3} - 2}{\sqrt{3} - 2} \\ &= \frac{\sqrt{3} - 2}{(\sqrt{3})^2 - (2)^2} \\ &= \frac{\sqrt{3} - 2}{3 - 4} \\ &= \frac{\sqrt{3} - 2}{-1} \\ &= 2 - \sqrt{3} \end{aligned}$$

(ii) $\frac{1}{4 - \sqrt{5}}$

Sol.

$$\begin{aligned} &= \frac{1}{4 - \sqrt{5}} \times \frac{4 + \sqrt{5}}{4 + \sqrt{5}} \\ &= \frac{4 + \sqrt{5}}{(4)^2 - (\sqrt{5})^2} \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

29

$$= \frac{4 + \sqrt{5}}{16 - 5}$$

$$= \frac{4 + \sqrt{5}}{11}$$

(iii) $\frac{4\sqrt{3}}{\sqrt{7} + \sqrt{5}}$

Sol.

$$= \frac{4\sqrt{3}}{\sqrt{7} + \sqrt{5}} \times \frac{\sqrt{7} - \sqrt{5}}{\sqrt{7} - \sqrt{5}}$$

$$= \frac{(4\sqrt{3})(\sqrt{7} - \sqrt{5})}{(\sqrt{7})^2 - (\sqrt{5})^2}$$

$$= \frac{(4\sqrt{3})(\sqrt{7} - \sqrt{5})}{7 - 5}$$

$$= \frac{4\sqrt{3}(\sqrt{7} - \sqrt{5})}{2}$$

$$2\sqrt{3}(\sqrt{7} - \sqrt{5})$$

(iv) $\frac{\sqrt{x} - \sqrt{y}}{\sqrt{x} + \sqrt{y}}$

Sol.

$$= \frac{\sqrt{x} - \sqrt{y}}{\sqrt{x} + \sqrt{y}} \times \frac{\sqrt{x} - \sqrt{y}}{\sqrt{x} - \sqrt{y}}$$

$$= \frac{(\sqrt{x} - \sqrt{y})^2}{(\sqrt{x})^2 - (\sqrt{y})^2}$$

$$= \frac{x + y - 2\sqrt{xy}}{x - y}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

30

(v) $\frac{5\sqrt{7}}{2+3\sqrt{7}}$

Sol.

$$\begin{aligned} &= \frac{5\sqrt{7}}{2+3\sqrt{7}} \times \frac{2-3\sqrt{7}}{2-3\sqrt{7}} \\ &= \frac{5\sqrt{7}(2-3\sqrt{7})}{(2)^2 - (3\sqrt{7})^2} \\ &= \frac{(5\sqrt{7})(2) - (5\sqrt{7})(3\sqrt{7})}{4 - 9 \times 7} \\ &= \frac{10\sqrt{7} - 105}{4 - 63} \\ &= \frac{10\sqrt{7} - 105}{-59} \\ &= \frac{105 - 10\sqrt{7}}{59} \end{aligned}$$

(vi) $\frac{\sqrt{3} + \sqrt{2}}{\sqrt{3} - \sqrt{2}}$

Sol.

$$\begin{aligned} &= \frac{\sqrt{3} + \sqrt{2}}{\sqrt{3} - \sqrt{2}} \times \frac{\sqrt{3} + \sqrt{2}}{\sqrt{3} + \sqrt{2}} \\ &= \frac{(\sqrt{3} + \sqrt{2})^2}{(\sqrt{3})^2 - (\sqrt{2})^2} \\ &= \frac{3 + 2 + 2\sqrt{3} \times 2}{3 - 2} \\ &= \frac{5 + 2\sqrt{6}}{1} \\ &= 5 + 2\sqrt{6} \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

31

(vii) $\frac{29}{11+3\sqrt{5}}$

Sol.

$$\begin{aligned} &= \frac{29}{11+3\sqrt{5}} \times \frac{11-3\sqrt{5}}{11-3\sqrt{5}} \\ &= \frac{29(11-3\sqrt{5})}{(11)^2 - (3\sqrt{5})^2} \\ &= \frac{29(11-3\sqrt{5})}{121 - 9 \times 5} \\ &= \frac{29(11-3\sqrt{5})}{121 - 45} \\ &= \frac{29(11-3\sqrt{5})}{76} \end{aligned}$$

(viii) $\frac{17}{3\sqrt{7}+2\sqrt{3}}$

Sol.

$$\begin{aligned} &= \frac{17}{3\sqrt{7}+2\sqrt{3}} \times \frac{3\sqrt{7}-2\sqrt{3}}{3\sqrt{7}-2\sqrt{3}} \\ &= \frac{17(3\sqrt{7}-2\sqrt{3})}{(3\sqrt{7})^2 - (2\sqrt{3})^2} \\ &= \frac{17(3\sqrt{7}-2\sqrt{3})}{9 \times 7 - 4 \times 3} \\ &= \frac{17(3\sqrt{7}-2\sqrt{3})}{63 - 12} \\ &= \frac{17(3\sqrt{7}-2\sqrt{3})}{51} \\ &= \frac{3\sqrt{7}-2\sqrt{3}}{3} \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

32

Q.4 If $x = \sqrt{5} + 2$, then find the values of (i) $x + \frac{1}{x}$ and

(ii) $x^2 + \frac{1}{x^2}$.

Sol. (i) $x = \sqrt{5} + 2$(1)

$$\begin{aligned}\frac{1}{x} &= \frac{1}{\sqrt{5} + 2} \\ &= \frac{1}{\sqrt{5} + 2} \times \frac{\sqrt{5} - 2}{\sqrt{5} - 2} \\ &= \frac{\sqrt{5} - 2}{(\sqrt{5})^2 - (2)^2} \\ &= \frac{\sqrt{5} - 2}{5 - 4}\end{aligned}$$

$$\frac{1}{x} = \sqrt{5} - 2$$

$$x + \frac{1}{x} = \sqrt{5} + 2 + \sqrt{5} - 2 \quad \text{from (1) + (2)}$$

Thus

$$x + \frac{1}{x} = 2\sqrt{5}$$

(ii) $\left(x + \frac{1}{x}\right)^2 = (2\sqrt{5})^2$

Taking square root

$$x^2 + \frac{1}{x^2} + 2 = 4 \times 5$$

$$x^2 + \frac{1}{x^2} + 2 = 20$$

$$x^2 + \frac{1}{x^2} = 20 - 2$$

Hence,

$$x^2 + \frac{1}{x^2} = 18$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

33

- Q.5 If $x = 2 + \sqrt{3}$, then find the values of (i) $x - \frac{1}{x}$ and
 (ii) $x^2 + \frac{1}{x^2}$.

Sol. (i) $x = 2 + \sqrt{3}$ (1)

$$\frac{1}{x} = \frac{1}{2 + \sqrt{3}}$$

$$= \frac{1}{2 + \sqrt{3}} \times \frac{2 - \sqrt{3}}{2 - \sqrt{3}} \quad (\text{Rationalize the denominators})$$

$$= \frac{2 - \sqrt{3}}{(2)^2 - (\sqrt{3})^2}$$

$$= \frac{2 - \sqrt{3}}{4 - 3} = \frac{2 - \sqrt{3}}{1} = 2 - \sqrt{3} \quad \text{.....(2)}$$

$$x - \frac{1}{x} = (2 + \sqrt{3}) - (2 - \sqrt{3}) \quad \text{from (1), (2)}$$

$$x - \frac{1}{x} = 2 + \sqrt{3} - 2 + \sqrt{3}$$

Hence, $x - \frac{1}{x} = 2\sqrt{3}$

(ii) $\left(x - \frac{1}{x}\right)^2 = (2\sqrt{3})^2$ Taking square root

$$x^2 + \frac{1}{x^2} - 2 = 4 \times 3$$

$$x^2 + \frac{1}{x^2} - 2 = 12$$

Hence, $x^2 + \frac{1}{x^2} = 12 + 2 = 14$

- Q.6 If $x = \sqrt{3} - \sqrt{2}$, then find the values of (i) $x - \frac{1}{x}$ and

(ii) $x^2 + \frac{1}{x^2}$.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

34

Sol. (i) $x = \sqrt{3} - \sqrt{2}$ (1)

$$\begin{aligned}\frac{1}{x} &= \frac{1}{\sqrt{3} - \sqrt{2}} \\ &= \frac{1}{(\sqrt{3} - \sqrt{2})} \times \frac{(\sqrt{3} + \sqrt{2})}{(\sqrt{3} + \sqrt{2})} \\ &= \frac{(\sqrt{3} + \sqrt{2})}{(\sqrt{3})^2 - (\sqrt{2})^2} \\ &= \frac{(\sqrt{3} + \sqrt{2})}{3 - 2}\end{aligned}$$

$$\frac{1}{x} = \sqrt{3} + \sqrt{2} \text{(2)}$$

$$\begin{aligned}x - \frac{1}{x} &= (\sqrt{3} - \sqrt{2}) - (\sqrt{3} + \sqrt{2}) \quad \text{from (1), (2)} \\ &= \sqrt{3} - \sqrt{2} - \sqrt{3} - \sqrt{2}\end{aligned}$$

Hence, $\boxed{x - \frac{1}{x} = -2\sqrt{2}}$

(ii) $\left(x - \frac{1}{x}\right)^2 = (-2\sqrt{2})^2$ Taking square root

$$x^2 + \frac{1}{x^2} - 2 = 4 \times 2$$

$$\left(x^2 + \frac{1}{x^2} - 2 = 8\right)$$

$$x^2 + \frac{1}{x^2} = 8 + 2$$

Hence, $\boxed{x^2 + \frac{1}{x^2} = 10}$

Q.7 If $\frac{1}{x} = 3 - \sqrt{2}$, then evaluate (i) $x + \frac{1}{x}$ and (ii) $x - \frac{1}{x}$.

Sol. (i) $\frac{1}{x} = 3 - \sqrt{2}$ (1)

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

35

$$\begin{aligned}x &= \frac{1}{3-\sqrt{2}} \\&= \frac{1}{3-\sqrt{2}} \times \frac{3+\sqrt{2}}{3+\sqrt{2}} \\&= \frac{3+\sqrt{2}}{(3)^2 - (\sqrt{2})^2} \\&= \frac{3+\sqrt{2}}{9-2} \\x &= \frac{3+\sqrt{2}}{7} \dots\dots\dots(2)\end{aligned}$$

$$\begin{aligned}x + \frac{1}{x} &= \frac{3+\sqrt{2}}{7} + 3-\sqrt{2} \quad \text{Now from (1) + (2)} \\&= \frac{3+\sqrt{2} + 7(3-\sqrt{2})}{7} \\&= \frac{3+\sqrt{2} + 21-7\sqrt{2}}{7} \\&= \frac{24+\sqrt{2}-7\sqrt{2}}{7} \\&= \frac{24+\sqrt{2}(1-7)}{7} \\&= \frac{24+\sqrt{2}(-6)}{7}\end{aligned}$$

Hence, $x + \frac{1}{x} = \frac{24-6\sqrt{2}}{7}$

$$\begin{aligned}\text{(ii)} \quad x - \frac{1}{x} &= \frac{3+\sqrt{2}}{7} - (3-\sqrt{2}) \quad \text{from (1),(2)} \\&= \frac{3+\sqrt{2}}{7} - 3 + \sqrt{2}\end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

36

$$= \frac{3 + \sqrt{2} - 21 + 7\sqrt{2}}{7}$$

$$= \frac{-18 + \sqrt{2} + 7\sqrt{2}}{7}$$

$$= \frac{-18 + \sqrt{2}(1+7)}{7}$$

Hence,
$$x - \frac{1}{x} = \frac{-18 + 8\sqrt{2}}{7}$$

Q.8 If $\frac{1}{p} = \sqrt{10} + 3$, then evaluate (i) $\left(p + \frac{1}{p}\right)^2$ and

(ii) $\left(p - \frac{1}{p}\right)^2$

Sol.

(i) $\frac{1}{p} = \sqrt{10} + 3 \dots\dots\dots (1)$

$$p = \frac{1}{\sqrt{10} + 3}$$

$$p = \frac{1}{\sqrt{10} + 3} \times \frac{\sqrt{10} - 3}{\sqrt{10} - 3} \text{ (Rationalize the denominators)}$$

$$= \frac{\sqrt{10} - 3}{(\sqrt{10})^2 - (3)^2}$$

$$= \frac{\sqrt{10} - 3}{10 - 9}$$

$$= \frac{\sqrt{10} - 3}{1}$$

$$= \sqrt{10} - 3 \dots\dots\dots (2)$$

$$p + \frac{1}{p} = (\sqrt{10} - 3) + (\sqrt{10} + 3) \text{ from (1)+(2)}$$

$$= \sqrt{10} - 3 + \sqrt{10} + 3$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

37

$$p + \frac{1}{p} = 2\sqrt{10}$$

$$\left(p + \frac{1}{p}\right)^2 = (2\sqrt{10})^2 \quad \text{Taking square root}$$

$$= 4 \times 10$$

$$\text{Hence, } \boxed{\left(p + \frac{1}{p}\right)^2 = 40}$$

$$(ii) \quad p - \frac{1}{p} = (\sqrt{10} - 3) - (\sqrt{10} + 3) \quad \text{Now from (1) - (2)}$$

$$= \sqrt{10} - 3 - \sqrt{10} - 3$$

$$p - \frac{1}{p} = -6$$

$$\left(p - \frac{1}{p}\right)^2 = (-6)^2 \quad \text{Taking square root}$$

$$\text{Hence, } \boxed{\left(p - \frac{1}{p}\right)^2 = 36}$$

Q.9 Rationalize

$$(i) \quad \frac{b + \sqrt{b^2 - a^2}}{b - \sqrt{b^2 - a^2}}$$

$$\text{Sol. } = \frac{b + \sqrt{b^2 - a^2}}{b - \sqrt{b^2 - a^2}} \times \frac{b + \sqrt{b^2 - a^2}}{b + \sqrt{b^2 - a^2}}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 1)

Pilot Super One "General Math" 10th

38

$$\begin{aligned} &= \frac{(h + \sqrt{h^2 - a^2})^2}{(h)^2 - (\sqrt{h^2 - a^2})^2} \\ &= \frac{h^2 + h^2 - a^2 + 2h\sqrt{h^2 - a^2}}{h^2 - (h^2 - a^2)} \\ &= \frac{2h^2 - a^2 + 2h\sqrt{h^2 - a^2}}{h^2 - h^2 + a^2} \\ &= \frac{2h^2 - a^2 + 2h\sqrt{h^2 - a^2}}{a^2} \end{aligned}$$

(ii) $\frac{\sqrt{a+3} - \sqrt{a-3}}{\sqrt{a+3} + \sqrt{a-3}}$

Sol.

$$\begin{aligned} &= \frac{\sqrt{a+3} - \sqrt{a-3}}{\sqrt{a+3} + \sqrt{a-3}} \times \frac{\sqrt{a+3} - \sqrt{a-3}}{\sqrt{a+3} - \sqrt{a-3}} \\ &= \frac{(\sqrt{a+3} - \sqrt{a-3})^2}{(\sqrt{a+3})^2 - (\sqrt{a-3})^2} \\ &= \frac{(a+3) + (a-3) - 2\sqrt{(a+3)(a-3)}}{(a+3) - (a-3)} \\ &= \frac{a+3+a-3-2\sqrt{(a)^2 - (3)^2}}{a+3-a+3} \\ &= \frac{2a-2\sqrt{a^2-9}}{6} \\ &= \frac{2(a-\sqrt{a^2-9})}{6} \\ &= \frac{a-\sqrt{a^2-9}}{3} \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

Pilot Super One "General Math" 10th

43

Exercise 2.10

Factorize:

Q.1 $3a(x + y) - 7b(x + y)$

Sol: $= (x + y)(3a - 7b)$

Q.2 $ax + ay - x^2 - xy$

Sol: $= (ax + ay) - (x^2 + xy)$

$= a(x + y) - x(x + y)$

$= (x + y)(a - x)$

Q.3 $a^3 + a - 3a^2 - 3$

Sol: $= (a^3 + a) - (3a^2 + 3)$

$= a(a^2 + 1) - 3(a^2 + 1)$

$= (a^2 + 1)(a - 3)$

Q.4 $x^3 + y - xy - x$

Sol: $= x^3 - x - xy + y$ (writing in order)

$= x(x^2 - 1) - y(x - 1)$

$= x(x - 1)(x + 1) - y(x - 1)$

$= (x - 1)[x(x + 1) - y]$

$= (x - 1)(x^2 + x - y)$

Q.5 $3ax + 6ay - 8bx - 4by$

Sol: $= 3ax - 4bx + 6ay - 8by$ (writing in order)

$= x(3a - 4b) + 2y(3a - 4b)$

$= (3a - 4b)(x + 2y)$

Q.6 $2a^2 - bc - 2ab + ac$

Sol: $= 2a^2 + ac - 2ab - bc$ (writing in order)

$= a(2a + c) - b(2a + c)$

$= (2a + c)(a - b)$

$= (a - b)(2a + c)$

Q.7 $a(a - b + c) - bc$

Sol: $= a^2 - ab + ca - bc$

$= a(a - b) + c(a - b)$

$= (a - b)(a + c)$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

Pilot Super One "General Math" 10th

44

Q.8 $8 - 4a - 2a^3 + a^4$

Sol: $= 4(2 - a) - a^3(2 - a)$
 $= (2 - a)(4 - a^3)$
 $= (4 - a^3)(2 - a)$

Q.9 $16x^2 - 24xa + 9a^2$

Sol: $= 16x^2 - 12xa - 12xa + 9a^2$
 $= 4x(4x - 3a) - 3a(4x - 3a)$
 $= (4x - 3a)(4x - 3a)$
 $= (4x - 3a)^2$

Q.10 $1 - 14x + 49x^2$

Sol: $= (1)^2 - 2(1)(7x) + (7x)^2$
 $= (1 - 7x)^2$

Q.11 $20x^2 + 5 - 20x$

Sol: $= 20x^2 - 20x + 5$
 $= 5[4x^2 - 4x + 1]$
 $= 5[(2x)^2 - 2(2x)(1) + (1)^2]$
 $= 5(2x - 1)^2$

Q.12 $2a^3b + 2ab^3 - 4a^2b^2$

Sol: $= 2ab[a^2 + b^2 - 2ab]$
 $= 2ab[a^2 - 2ab + b^2]$
 $= 2ab[(a)^2 - 2(a)(b) + (b)^2]$
 $= 2ab(a - b)^2$

Q.13 $(x^2)^2 + x + \frac{1}{4}$

Sol: $= (x)^2 + 2(x)\left(\frac{1}{2}\right) + \left(\frac{1}{2}\right)^2$
 $= \left(x + \frac{1}{2}\right)^2$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

Pilot Super One "General Math" 10th

45

Q.14 $x^3 + \frac{1}{x^3} - 2$

Sol:
$$= x^3 - 2 + \frac{1}{x^3}$$

$$= (x)^3 - 2(x)\left(\frac{1}{x}\right) + \left(\frac{1}{x}\right)^3$$

$$= \left(x - \frac{1}{x}\right)^3$$

Q.15 $5x^3 - 30x^2 + 45x$

Sol:
$$= 5x[x^2 - 6x + 9]$$

$$= 5x[(x)^2 - 2(x)(3) + (3)^2]$$

$$= 5x(x - 3)^2$$

Q.16 $a^2 + b^2 + 2ab + 2bc + 2ac$

Sol:
$$= (a^2 + 2ab + b^2) + 2ac + 2bc$$

$$= [(a)^2 + 2(a)(b) + (b)^2] + 2c(a + b)$$

$$= (a + b)^2 + 2c(a + b)$$

$$= (a + b)[(a + b) + 2c]$$

$$= (a + b)(a + b + 2c)$$

Exercise 2.4

Resolve into factors.

Q.1 $x^2 + 2xy + y^2 - a^2$

Sol:
$$= (x^2 + 2xy + y^2) - a^2$$

$$= [(x)^2 + 2(x)(y) + (y)^2] - (a)^2$$

$$= (x + y)^2 - (a)^2$$

$$= (x + y + a)(x + y - a)$$

Q.2 $4a^2 - 4ab + b^2 - 9c^2$

Sol:
$$= (4a^2 - 4ab + b^2) - 9c^2$$

$$= [(2a)^2 - 2(2a)(b) + (b)^2] - (3c)^2$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

Pilot Super One "General Math" 10th

46

$$= (2a - b)^2 - (3c)^2$$

$$= (2a - b + 3c)(2a - b - 3c)$$

Q.3 $x^2 + 6ax + 9a^2 - 16b^2$

Sol: $= (x^2 + 6ax + 9a^2) - 16b^2$

$$= [(x)^2 + 2(x)(3a) + (3a)^2] - (4b)^2$$

$$= (x + 3a)^2 - (4b)^2$$

$$= (x + 3a + 4b)(x + 3a - 4b)$$

Q.4 $y^2 - c^2 + 2cx - x^2$

Sol: $= y^2 - (c^2 - 2cx + x^2)$

$$= (y)^2 - [(c)^2 - 2(c)(x) + (x)^2]$$

$$= (y)^2 - (c - x)^2 \quad \text{Formula: } [a^2 - b^2 = (a + b)(a - b)]$$

$$= [y + (c - x)][y - (c - x)]$$

$$= (y + c - x)(y - c + x)$$

$$= (y - x + c)(y + x - c)$$

Q.5 $x^2 + y^2 + 2xy - 4x^2y^2$

Sol: $= (x^2 + 2xy + y^2) - 4x^2y^2$

$$= [(x)^2 + 2(x)(y) + (y)^2] - (2xy)^2$$

$$= (x + y)^2 - (2xy)^2$$

$$= (x + y + 2xy)(x + y - 2xy)$$

Q.6 $a^2 - 4ab + 4b^2 - 9a^2c^2$

Sol: $= (a^2 - 4ab + 4b^2) - (9a^2c^2)$

$$= [(a)^2 - 2(a)(2b) + (2b)^2] - [3ac]^2$$

$$= (a - 2b)^2 - (3ac)^2$$

$$= (a - 2b + 3ac)(a - 2b - 3ac)$$

Q.7 $x^2 - 2xy + y^2 - a^2 + 2ab - b^2$

Sol: $= (x^2 - 2xy + y^2) - (a^2 - 2ab + b^2)$

$$= [(x)^2 - 2(x)(y) + (y)^2] [(a)^2 - 2(a)(b) + (b)^2]$$

$$= (x - y)^2 - (a - b)^2$$

$$= [(x - y) + (a - b)][(x - y) - (a - b)]$$

$$= (x - y + a - b)(x - y - a + b)$$

Q.8 $y^4 + 4$

Sol: $= (y^2)^2 + (2)^2 + 2(y^2)(2) - 2(y^2)(2) \text{ (completing square root)}$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

Pilot Super One "General Math" 10th

47

$$\begin{aligned} &= [(y^2)^2 + (2)^2 + 4y^2] - 4y^2 \\ &= (y^2 + 2)^2 - (2y)^2 \\ &= (y^2 + 2 + 2y)(y^2 + 2 - 2y) \\ &= (y^2 + 2y + 2)(y^2 - 2y + 2) \end{aligned}$$

Q.9 $z^4 + 64y^4$

Sol: $\begin{aligned} &= (z^2)^2 + (8y^2)^2 + 2(z^2)(8y^2) - 2(z^2)(8y^2) \text{ (completing square root)} \\ &= (z^2 + 8y^2)^2 - 16z^2y^2 \\ &= (z^2 + 8y^2)^2 - (4zy)^2 \\ &= (z^2 + 8y^2 - 4zy)(z^2 + 8y^2 + 4zy) \end{aligned}$

Q.10 $x^4 + 324$

Sol: $\begin{aligned} &= (x^2)^2 + (18)^2 + 2(x^2)(18) - 2(x^2)(18) \text{ (completing square root)} \\ &= (x^2 + 18)^2 - 36x^2 \\ &= (x^2 + 18)^2 - (6x)^2 \\ &= (x^2 + 18 + 6x)(x^2 + 18 - 6x) \\ &= (x^2 + 6x + 18)(x^2 - 6x + 18) \end{aligned}$

Q.11 $z^4 - z^2 + 16$

Sol: $\begin{aligned} &= (z^2)^2 + (4)^2 + 2(z^2)(4) - 9z^2 \text{ (completing square root)} \\ &= (z^2 + 4)^2 - (3z)^2 \\ &= (z^2 + 4 - 3z)(z^2 + 4 + 3z) \\ &= (z^2 - 3z + 4)(z^2 + 3z + 4) \end{aligned}$

Q.12 $4x^4 - 5x^2y^2 + y^4$

Sol: $\begin{aligned} &= (2x^2)^2 - 5x^2y^2 + (y^2)^2 \text{ (completing square root)} \\ &= (2x^2)^2 + 2(2x^2)(y^2) + (y^2)^2 - 9x^2y^2 \text{ (completing square root)} \\ &= (2x^2 + y^2)^2 - (3xy)^2 \\ &= (2x^2 + y^2 - 3xy)(2x^2 + y^2 + 3xy) \end{aligned}$

Exercise 2.3

Factorize:

Q.1 $x^2 + 9x + 20$

Sol: $\begin{aligned} &= x^2 + 4x + 5x + 20 \\ &= (x^2 + 4x) + (5x + 20) \end{aligned}$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

=====

Pilot Super One "General Math" 10th

48

$$= x(x + 4) + 5(x + 4)$$

$$= (x + 4)(x + 5)$$

Q.2 $x^2 + 5x - 14$

Sol: $= x^2 + 7x - 2x - 14$
 $= (x^2 + 7x) - (2x + 14)$
 $= x(x + 7) - 2(x + 7)$
 $= (x + 7)(x - 2)$

Q.3 $x^2 + 5x - 6$

Sol: $= x^2 + 6x - x - 6$
 $= (x^2 + 6x) - (x + 6)$
 $= x(x + 6) - 1(x + 6)$
 $= (x + 6)(x - 1)$

Q.4 $x^2 - 7x + 12$

Sol: $= x^2 - 3x - 4x + 12$
 $= (x^2 - 3x) - (4x - 12)$
 $= x(x - 3) - 4(x - 3)$
 $= (x - 3)(x - 4)$

Q.5 $x^2 - x - 156$

Sol: $= x^2 - 13x + 12x - 156$
 $= (x^2 - 13x) + (12x - 156)$
 $= x(x - 13) + 12(x - 13)$
 $= (x - 13)(x + 12)$

Q.6 $x^2 - x - 2$

Sol: $= x^2 - 2x + x - 2$
 $= (x^2 - 2x) + (x - 2)$
 $= x(x - 2) + 1(x - 2)$
 $= (x - 2)(x + 1)$

Q.7 $x^2 - 9x - 90$

Sol: $= x^2 - 15x + 6x - 90$
 $= (x^2 - 15x) + (6x - 90)$
 $= x(x - 15) + 6(x - 15)$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

Pilot Super One "General Math" 10th

49

$$= (x - 15)(x + 6)$$

Q.8 $a^2 - 12a - 85$

Sol: $= a^2 - 17a + 5a - 85$
 $= (a^2 - 17a) + (5a - 85)$
 $= a(a - 17) + 5(a - 17)$
 $= (a - 17)(a + 5)$

Q.9 $98 - 7x - x^2$

Sol: $= 98 - 14x + 7x - x^2$
 $= (98 - 14x) + (7x - x^2)$
 $= 14(7 - x) + x(7 - x)$
 $= (7 - x)(14 + x)$

Q.10 $y^2 - 11y - 152$

Sol: $= y^2 - 19y + 8y - 152$
 $= (y^2 - 19y) + (8y - 152)$
 $= y(y - 19) + 8(y - 19)$
 $= (y - 19)(y + 8)$

Q.11 $2x^2 + 3x + 1$

Sol: $= 2x^2 + 2x + x + 1$
 $= (2x^2 + 2x) + (x + 1)$
 $= 2x(x + 1) + 1(x + 1)$
 $= (x + 1)(2x + 1)$

Q.12 $3x^2 + 5x + 2$

Sol: $= 3x^2 + 3x + 2x + 2$
 $= (3x^2 + 3x) + (2x + 2)$
 $= 3x(x + 1) + 2(x + 1)$
 $= (x + 1)(3x + 2)$

Q.13 $2x^2 - x - 1$

Sol: $= 2x^2 - 2x + x - 1$
 $= (2x^2 - 2x) + (x - 1)$
 $= 2x(x - 1) + 1(x - 1)$
 $= (x - 1)(2x + 1)$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

Pilot Super One "General Math" 10th

50

Q.14 $6x^2 + 7x - 3$

Sol: $= 6x^2 + 9x - 2x - 3$
 $= (6x^2 + 9x) - (2x + 3)$
 $= 3x(2x + 3) - 1(2x + 3)$
 $= (2x + 3)(3x - 1)$

Q.15 $2 - 3x - 2x^2$

Sol: $= 2 - 4x + x - 2x^2$
 $= (2 - 4x) + (x - 2x^2)$
 $= 2(1 - 2x) + x(1 - 2x)$
 $= (1 - 2x)(2 + x)$

Q.16 $8 + 6x - 5x^2$

Sol: $= 8 + 10x - 4x - 5x^2$
 $= (8 + 10x) - (4x + 5x^2)$
 $= 2(4 + 5x) - x(4 + 5x)$
 $= (4 + 5x)(2 - x)$

Q.17 $3u^2 - 10u + 8$

Sol: $= 3u^2 - 6u - 4u + 8$
 $= (3u^2 - 6u) - (4u - 8)$
 $= 3u(u - 2) - 4(u - 2)$
 $= (u - 2)(3u - 4)$

Q.18 $10x^2 - 7x - 12$

Sol: $= 10x^2 - 15x + 8x - 12$
 $= (10x^2 - 15x) + (8x - 12)$
 $= 5x(2x - 3) + 4(2x - 3)$
 $= (2x - 3)(5x + 4)$

Q.19 $5x^2 - 32x + 12$

Sol: $= 5x^2 - 30x - 2x + 12$
 $= (5x^2 - 30x) - (2x - 12)$
 $= 5x(x - 6) - 2(x - 6)$
 $= (x - 6)(5x - 2)$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

Pilot Super One "General Math" 10th

51

Q.20 $4\sqrt{3}x^2 + 5x - 2\sqrt{3}$

Sol: $4\sqrt{3}x^2 + 8x - 3x - 2\sqrt{3}$
 $(4\sqrt{3}x^2 + 8x) - (3x + 2\sqrt{3})$
 $4x(\sqrt{3}x + 2) - \sqrt{3}(\sqrt{3}x + 2)$
 $= (4\sqrt{3}x + 2)(4x - \sqrt{3})$

Formulae

- (i) $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$
- (ii) $(a - b)^3 = a^3 - 3a^2b + 3ab^2 - b^3$
- (iii) $a^3 + b^3 = (a + b)(a^2 - ab + b^2)$
- (iv) $a^3 - b^3 = (a - b)(a^2 + ab + b^2)$

EXERCISE 6.4

Factorize:

Q.1 $8x^3 - y^3$

Sol: $(2x)^3 - (y)^3$
 $(2x - y)[(2x)^2 + (2x)(y) + (y)^2]$
 $(2x - y)(4x^2 + 2xy + y^2)$

Q.2 $27x^3 + 1$

Sol: $=(3x)^3 + (1)^3$
 $=(3x + 1)[(3x)^2 - (3x)(1) + (1)^2]$
 $=(3x + 1)(9x^2 - 3x + 1)$

Q.3 $1 - 343x^3$

Sol: $=(1)^3 - (7x)^3$
 $(1 - 7x)[(1)^2 + 1(7x) + (7x)^2]$
 $(1 - 7x)(1 + 7x + 49x^2)$

Q.4 $a^3b^3 + 512$

Sol: $=(ab)^3 + (8)^3$
 $(ab + 8)[(ab)^2 - (ab)(8) + (8)^2]$
 $(ab + 8)(a^2b^2 - 8ab + 64)$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

Pilot Super One "General Math" 10th

52

Q.5 $27 - 1000y^3$

Sol: $= (3)^3 - (10y)^3$
 $= (3 - 10y)[(3)^2 + (3)(10y) + (10y)^2]$
 $= (3 - 10y)(9 + 30y + 100y^2)$

Q.6 $27x^3 - 64y^3$

Sol: $= (3x)^3 - (4y)^3$
 $= (3x - 4y)[(3x)^2 + (3x)(4y) + (4y)^2]$
 $= (3x - 4y)(9x^2 + 12xy + 16y^2)$

Q.7 $x^3y^3 + z^3$

Sol: $= (xy)^3 + (z)^3$
 $= (xy + z)[(xy)^2 + (xy)(z) + (z)^2]$
 $= (xy + z)(x^2y^2 + xyz + z^2)$

Q.8 $216p^3 - 343$

Sol: $= (6p)^3 - (7)^3$
 $= (6p - 7)[(6p)^2 + (6p)(7) + (7)^2]$
 $= (6p - 7)(36p^2 + 42p + 49)$

Q.9 $8x^3 - \frac{1}{27}$

Sol: $= (2x)^3 - \left(\frac{1}{3}\right)^3$
 $= \left(2x - \frac{1}{3}\right) \left[(2x)^2 + (2x)\left(\frac{1}{3}\right) + \left(\frac{1}{3}\right)^2 \right]$
 $= \left(2x - \frac{1}{3}\right) \left[4x^2 + \frac{2}{3}x + \frac{1}{9} \right]$

Q.10 $a^3 + b^3 + a + b$

Sol: $= (a^3 + b^3) + (a + b)$
 $= (a + b)(a^2 - ab + b^2) + (a + b)$
 $= (a + b)(a^2 - ab + b^2 + 1)$
 $= (a + b)(a^2 - ab + b^2 + 1)$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

Pilot Super One "General Math" 10th

53

Q.11 $a - b - a^3 + b^3$

Sol: $= (a - b) - (a^3 - b^3)$
 $= (a - b) - [(a - b)(a^2 + ab + b^2)]$
 $= (a - b) [1 - (a^2 + ab + b^2)]$
 $= (a - b)(1 - a^2 - ab - b^2)$

Q.12 $x - 8xy^3$

Sol: $= x(1 - 8y^3)$
 $= x[(1)^3 - (2y)^3]$
 $= x[(1 - 2y)\{(1)^2 + (1)(2y) + (2y)^2\}]$
 $= x[(1 - 2y)(1 + 2y + 4y^2)]$
 $= x(1 - 2y)(1 + 2y + 4y^2)$

Q.13 $x^{12} - y^{12}$

Sol: $= (x^6)^2 - (y^6)^2$
 $= (x^6 - y^6)(x^6 + y^6)$
 $= [(x^3)^2 - (y^3)^2][(x^2)^3 + (y^2)^3]$
 $= (x^3 - y^3)(x^3 + y^3)(x^2 + y^2)(x^4 - x^2y^2 + y^4)$
 $= (x - y)(x^2 + xy + y^2)(x + y)(x^2 - xy + y^2)(x^2 + y^2)$
 $= (x - y)(x + y)(x^2 + y^2)(x^2 + xy + y^2)(x^2 - xy + y^2)$
 $= (x - y)(x + y)(x^2 + y^2)(x^2 + xy + y^2)(x^2 - xy + y^2)$

Q.14 $1 - \frac{64p^3}{q^3}$

Sol: $= (1)^3 - \left(\frac{4p}{q}\right)^3$
 $= \left(1 - \frac{4p}{q}\right) \left[(1)^2 + (1)\left(\frac{4p}{q}\right) + \left(\frac{4p}{q}\right)^2\right]$
 $= \left(1 - \frac{4p}{q}\right) \left[1 + \frac{4p}{q} + \frac{16p^2}{q^2}\right]$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

Pilot Super One "General Math" 10th

54

Q.15 $1 + 64u^3$

Sol: $= (1)^3 + (4u)^3$
 $= (1 + 4u)[(1)^2 - (1)(4u) + (4u)^2]$
 $= (1 + 4u)(1 - 4u + 16u^2)$

Q.16 $8x^3 - 6x - 9y + 27y^3$

Sol: $= 8x^3 + 27y^3 - 6x - 9y$
 $= (8x^3 + 27y^3) - (6x + 9y)$
 $= [(2x)^3 + (3y)^3] - 3(2x + 3y)$
 $= (2x + 3y)[(2x)^2 - (2x)(3y) + (3y)^2] - 3(2x + 3y)$
 $= (2x + 3y)[4x^2 - 6xy + 9y^2] - 3(2x + 3y)$
 $= (2x + 3y)[4x^2 - 6xy + 9y^2 - 3]$

Q.17 $z^3 + 125$

Sol: $= (z)^3 + (5)^3$
 $= (z + 5)[(z)^2 - (z)(5) + (5)^2]$
 $= (z + 5)(z^2 - 5z + 25)$

Q.18 $x^9 + y^9$

Sol: $= (x^3)^3 + (y^3)^3$
 $= (x^3 + y^3)[(x^3)^2 - (x^3)(y^3) + (y^3)^2]$
 $= (x^3 + y^3)(x^6 - x^3y^3 + y^6)$
 $= (x + y)(x^2 - xy + y^2)(x^6 - x^3y^3 + y^6)$

Q.19 $m^6 - n^6$

Sol: $= (m^3)^2 - (n^3)^2$
 $= (m^3 + n^3)(m^3 - n^3)$
 $= (m + n)(m^2 - mn + n^2)(m - n)(m^2 + mn + n^2)$
 $= (m + n)(m - n)(m^2 - mn + n^2)(m^2 + mn + n^2)$

Q.20 $64x^7 - xa^6$

Sol: $= x(64x^6 - a^6)$
 $= x\{(8x^3)^2 - (a^3)^2\}$
 $= x(8x^3 + a^3)(8x^3 - a^3)$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

Pilot Super One "General Math" 10th

55

$$x\{(2x)^3 + (a)^3\} \{(2x)^3 - (a)^3\}$$

$$x(2x + a)(4x^2 - 2ax + a^2)(2x - a)(4x^2 + 2ax + a^2)$$

$$x(2x + a)(2x - a)(4x^2 - 2ax + a^2)(4x^2 + 2ax + a^2)$$

Q.21 $x^3 - 27a^3$

Sol: $= (x)^3 - (3a)^3$

$$= (x - 3a)\{(x)^2 + (x)(3a) + (3a)^2\}$$

$$= (x - 3a)(x^2 + 3ax + 9a^2)$$

Q.22 $x^3 + 27a^3$

Sol: $= (x)^3 + (3a)^3$

$$= (x + 3a)\{(x)^2 + (x)(3a) + (3a)^2\}$$

$$= (x + 3a)(x^2 + 3ax + 9a^2)$$

EXERCISE # 2.1

1. Evaluate each of the polynomials for the value indicated.

Q.1 $P(x) = 2x^3 - 5x^2 + 7x - 7$; $P(2)$

Sol: $P(x) = 2x^3 - 5x^2 + 7x - 7$

$$P(2) = 2(2)^3 - 5(2)^2 + 7(2) - 7$$

$$= 2 \times 8 - 5 \times 4 + 7 \times 2 - 7$$

$$= 16 - 20 + 14 - 7$$

$$= 3$$

Q.2 $P(x) = x^4 - 10x^2 + 25x - 2$; $P(-4)$

Sol: $P(x) = x^4 - 10x^2 + 25x - 2$

$$P(-4) = (-4)^4 - 10(-4)^2 + 25(-4) - 2$$

$$= 256 - 160 - 100 - 2$$

$$= -6$$

Q.3 $P(x) = x^4 + 5x^3 - 13x^2 - 30$; $P(-1)$

Sol: $P(x) = x^4 + 5x^3 - 13x^2 - 30$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

Pilot Super One "General Math" 10th

56

$$\begin{aligned}P(-1) &= (-1)^4 + 5(-1)^3 - 13(-1)^2 - 30 \\&= 1 - 5 - 13 - 30 \\&= -47\end{aligned}$$

Q.4 $P(x) = x^5 - 10x^3 + 7x + 6$; $P(3)$

Sol: $P(x) = x^5 - 10x^3 + 7x + 6$

$$\begin{aligned}P(3) &= (3)^5 - 10(3)^3 + 7(3) + 6 \\&= 243 - 270 + 21 + 6 \\&= 0\end{aligned}$$

Q.5 $P(x) = x^4 + 4x^3 - 9x^2 + 19x + 6$; $P(-2)$

Sol: $P(x) = x^4 + 4x^3 - 9x^2 + 19x + 6$

$$\begin{aligned}P(-2) &= (-2)^4 + 4(-2)^3 - 9(-2)^2 + 19(-2) + 6 \\&= 16 + 4(-8) - 9(4) - 38 + 6 \\&= 16 - 32 - 36 - 38 + 6 \\&= -106 + 22 \\&= -84\end{aligned}$$

II. *Determine whether the second polynomial is a factor of the first polynomial without dividing (Hint: evaluate directly and use the factor theorem).*

Q.6 $x^{18} - 1$; $x + 1$

Sol: Let $P(x) = x^{18} - 1$

Here $x - a = x + 1$

Thus $a = -1$

$$\begin{aligned}\text{Now } P(-1) &= (-1)^{18} - 1 \\&= 1 - 1 = 0\end{aligned}$$

Since $P(-1) = 0$

Then by factor theorem $x + 1$ is a factor of $x^{18} - 1$.

Q.7 $x^{18} - 1$; $x - 1$

Sol: Let $P(x) = x^{18} - 1$

Here $x - a = x - 1$

Thus $a = 1$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

Pilot Super One "General Math" 10th

57

$$\begin{aligned}\text{Now } P(-1) &= (1)^{18} - 1 \\ &= 1 - 1 = 0\end{aligned}$$

Since $P(-1) = 0$

Then by factor theorem, $x - 1$ is a factor of $x^{18} - 1$.

Q.8 $x^9 - 2^9$; $x + 2$

Sol: Let $P(x) = x^9 - 2^9$

Here $x - a = x + 2$

Thus $a = -2$

$$\begin{aligned}\text{Now } P(-2) &= (-2)^9 - (2)^9 \\ &= -(2)^9 - (2)^9 \\ &= -ve \neq 0\end{aligned}$$

Since $P(-2) \neq 0$

Therefore, $x + 2$ is not a factor of $x^9 - 2^9$.

Q.9 $x^9 + 2^9$; $x - 2$

Sol: Let $P(x) = x^9 + 2^9$

Here $x - a = x - 2$

Thus $a = 2$

$$\begin{aligned}\text{Now } P(2) &= (2)^9 + (2)^9 \\ &\neq 0\end{aligned}$$

Since $P(2) \neq 0$

Therefore, $x - 2$ is not a factor of $x^9 + 2^9$.

Q.10 $3x^4 - 2x^3 + 5x - 6$; $x - 1$

Sol: Let $P(x) = 3x^4 - 2x^3 + 5x - 6$

Here $x - a = x - 1$

Thus $a = 1$

$$\begin{aligned}\text{Now } P(1) &= 3(1)^4 - 2(1)^3 + 5(1) - 6 \\ &= 3 - 2 + 5 - 6 \\ &= 8 - 8 = 0\end{aligned}$$

Since $P(1) = 0$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

=====

Pilot Super One "General Math" 10th

58

Then by factor theorem, $x - 1$ is a factor of $3x^4 - 2x^3 + 5x - 6$.

Q.11 $5x^6 - 7x^3 - 6x + x ; x - 1$

Sol: Let $P(x) = 5x^6 - 7x^3 - 6x + x$

Here $x - a = x - 1$

Thus $a = 1$

$$\begin{aligned}\text{Now } P(1) &= 5(1)^6 - 7(1)^3 - 6(1) + (1) \\ &= 5 - 7 - 6 + 1 \\ &= 6 - 13 \\ &= -7 \neq 0\end{aligned}$$

Since $P(1) \neq 0$

Then by factor theorem, $x - 1$ is not a factor of $5x^6 - 7x^3 - 6x + x$.

Q.12 $3x^3 - 7x^2 - 8x + 2 ; x + 1$

Sol: Let $P(x) = 3x^3 - 7x^2 - 8x + 2$

Here $x - a = x + 1$

therefore $a = -1$

$$\begin{aligned}\text{Now } P(-1) &= 3(-1)^3 - 7(-1)^2 - 8(-1) + 2 \\ &= -3 - 7 + 8 + 2 \\ &= 0\end{aligned}$$

Since $P(-1) = 0$

Then by factor theorem, $x + 1$ is a factor of $3x^3 - 7x^2 - 8x + 2$.

Q.13 $5x^5 - 2x^5 + 3x^3 + 6x + 2 ; x + 1$

Sol: Let $P(x) = 5x^5 - 2x^5 + 3x^3 + 6x + 2$

Here $x - a = x + 1$

therefore $a = -1$

$$\begin{aligned}\text{Now } P(-1) &= 5(-1)^5 - 2(-1)^5 + 3(-1)^3 + 6(-1) + 2 \\ &= 5 + 2 - 3 - 6 + 2 \\ &= 9 - 9 = 0\end{aligned}$$

Since $P(-1) = 0$

Then by factor theorem, $x + 1$ is a factor of $5x^5 - 2x^5 + 3x^3 + 6x + 2$.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

Pilot Super One "General Math" 10th

59

Q.14 $6x^3 + 2x^2 - x + 9 ; x - 1$

Sol: Let $P(x) = 6x^3 + 2x^2 - x + 9$

Here $x = a \quad x = 1$

therefore $a = 1$

Now $P(1) = 6(1)^3 + 2(1)^2 - (1) + 9$

$= 6 + 2 - 1 + 9$

$17 - 1 = 16 \neq 0$

Since $P(1) \neq 0$

Then by factor theorem, $x - 1$ is not a factor of $6x^3 + 2x^2 - x + 9$.

Q.15 $4x^3 - 3x^2 - 8x + 4 ; x - 2$

Sol: Let $P(x) = 4x^3 - 3x^2 - 8x + 4$

Here $x = a \quad x = 2$

therefore $a = 2$

Now $P(2) = 4(2)^3 - 3(2)^2 - 8(2) + 4$

$= 4(8) - 3(4) - 8(2) + 4$

$= 32 - 12 - 16 + 4$

$= 36 - 28 = 8 \neq 0$

Since $P(2) \neq 0$

Then by factor theorem, $x - 2$ is not a factor of $4x^3 - 3x^2 - 8x + 4$.

Q.16 $5x^3 + 3x^2 - x + 1 ; x + 1$

Sol: Let $P(x) = 5x^3 + 3x^2 - x + 1$

Here $x = a \quad x = -1$

therefore $a = -1$

Now $P(-1) = 5(-1)^3 + 3(-1)^2 - (-1) + 1$

$= 5(-1) + 3(1) + 1 + 1$

$= -5 + 3 + 1 + 1$

$= -5 + 5$

$= 0$

Since $P(-1) = 0$

Then by factor theorem, $x+1$ is a factor of $5x^3 + 3x^2 - x + 1$.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

=====

Pilot Super One "General Math" 10th

60

Q.17 $2y^3 - 8y^2 + y - 4 ; y - 4$

Sol: Let $P(y) = 2y^3 - 8y^2 + y - 4$

Here $y - a = y - 4$

therefore $a = 4$

$$\begin{aligned}\text{Now } P(4) &= 2(4)^3 - 8(4)^2 + (4) - 4 \\ &= 2(64) - 8(16) + 4 - 4 \\ &= 128 - 128 + 4 - 4 \\ &= 132 - 132 \\ &= 0\end{aligned}$$

Since $P(4) = 0$

Then by factor theorem, $y - 4$ is not a factor of $2y^3 - 8y^2 + y - 4$.

Q.18 $z^3 - 5z^2 - 4z - 4 ; z + 2$

Sol: Let $P(z) = z^3 - 5z^2 - 4z - 4$

Here $z - a = z + 2$

therefore $a = -2$

$$\begin{aligned}\text{Now } P(-2) &= (-2)^3 - 5(-2)^2 - 4(-2) - 4 \\ &= -8 - 5(4) + 8 - 4 \\ &= -8 - 20 + 8 - 4 \\ &= -32 + 8 \\ &= -24 \neq 0\end{aligned}$$

Since $P(-2) \neq 0$

Then by factor theorem, $z+2$ is not a factor of $z^3 - 5z^2 - 4z - 4$.

III. Solve

Q.19. If $P(x) = x^3 - kx^2 + 3x + 5$ is divided by $x - 1$, find k , if remainder is 8.

Sol: Let $P(x) = x^3 - kx^2 + 3x + 5$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 2)

Pilot Super One "General Math" 10th

61

Here $x = a = x - 1$

therefore $a = 1$

$$\text{Now } P(1) = (1)^3 - k(1)^2 + 3(1) + 5$$

$$= 1 - k + 3 + 5$$

$$= 9 - k = \text{Remainder}$$

$$\text{But } 9 - k = 8$$

$$-k = 8 - 9$$

$$-k = -1$$

$$k = 1$$

Q.20. If $P(x) = 3x^3 + kx - 26$ is divided by $x - 2$, find k , if remainder is 0.

Sol: Let's $P(x) = 3x^3 + kx - 26$

Here $x = a = x - 2$

therefore $a = 2$

$$\text{Now } P(2) = 3(2)^3 + k(2) - 26$$

$$= 3(8) + 2k - 26$$

$$= 24 + 2k - 26$$

$$= 2k - 2 = \text{Remainder}$$

But Remainder = 0

$$\text{Thus, } 2k - 2 = 0$$

$$2k = 2$$

$$k = \frac{2}{2}$$

$$k = 1$$

UNIT 3 Algebraic Manipulation

- ▶ H.C.F and L.C.M
- ▶ Basic Operations on Algebraic Fractions
- ▶ Square Roots of Algebraic Fractions

After completion of this unit, the students will be able to:

- ▶ find highest common factor (HCF) and least common multiple (LCM) of algebraic expressions.
- ▶ use factor or division method to determine HCF and LCM.
- ▶ know the relationship between HCF and LCM.
- ▶ use HCF and LCM to reduce fractional expressions involving +, -, ×, ÷.
- ▶ find square root of an algebraic expression by factorization and division.

Exercise 3.1

Find H.C.F by factorization.

Q.1 $abxy, a^2bc$

Sol.

$$\text{Factorization of } abxy = \boxed{a} \times \boxed{b} \times x \times y$$

$$\text{Factorization of } a^2bc = \boxed{a} \times \boxed{b} \times a \times c$$

$$\text{common factors} = a, b$$

$$\begin{aligned}\text{Thus, H.C.F} &= a \times b \\ &= ab\end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

66

Q.2 $6pqr, 15qrs$

Sol.

$$\begin{aligned}\text{Factorization of } 6pqr &= 2 \times \boxed{3} \times p \times \boxed{q} \times \boxed{r} \\ \text{Factorization of } 15qrs &= 5 \times \boxed{3} \times \boxed{q} \times \boxed{r} \times s\end{aligned}$$

$$\text{Common factors} = 3, q, r$$

$$\begin{aligned}\text{Thus, H.C.F} &= 3 \times q \times r \\ &= 3qr\end{aligned}$$

Q.3 $8xy^2z^3, 12x^2y^2z^2$

Sol.

$$\begin{aligned}\text{Factorization of } 8xy^2z^3 &= 2 \times 2 \times 2 \times x \times y \times y \times z \times z \times z \\ \text{Factorization of } 12x^2y^2z^2 &= 2 \times 2 \times 3 \times x \times x \times y \times y \times z \times z\end{aligned}$$

$$\text{Common factors} = 2, 2, x, y, y, z, z$$

$$\begin{aligned}\text{Thus, H.C.F} &= 2 \times 2 \times x \times y \times y \times z \times z \\ &= 4xy^2z^2\end{aligned}$$

Q.4 $14a^2bc, 21ab^2$

Sol.

$$\begin{aligned}\text{Factorization of } 14a^2bc &= 2 \times \boxed{7} \times a \times a \times b \times c \\ \text{Factorization of } 21ab^2 &= 3 \times \boxed{7} \times a \times b \times b\end{aligned}$$

$$\text{Common factors} = 7, a, b$$

$$\begin{aligned}\text{Thus, H.C.F} &= 7 \times a \times b \\ &= 7ab\end{aligned}$$

Q.5 $3x^5y^2, 12x^2y^4, 15x^3y^2$

Sol.

$$\begin{aligned}\text{Factorization of } 3x^5y^2 &= 3 \times x \times x \times x \times x \times x \times y \times y \\ \text{Factorization of } 12x^2y^4 &= 2 \times 2 \times 3 \times x \times x \times y \times y \times y \times y \\ \text{Factorization of } 15x^3y^2 &= 3 \times 5 \times x \times x \times x \times y \times y\end{aligned}$$

$$\text{Common factors} = 3, x, x, y, y$$

$$\begin{aligned}\text{Thus, H.C.F} &= 3 \times x \times x \times y \times y \\ &= 3x^2y^2\end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

67

Q.6 $4abc^3, 8a^3bc, 6ab^3c$

Sol.

Factorization of $4abc^3 = 2 \times 2 \times a \times b \times c \times c \times c$

Factorization of $8a^3bc = 2 \times 2 \times 2 \times a \times a \times a \times b \times c$

Factorization of $6ab^3c = 2 \times 3 \times a \times b \times b \times b \times c$

Common factors = $2, a, b, c$

Thus, H.C.F = $2 \times a \times b \times c$
 $= 2abc$

Q.7 $x^3 + 64, x^2 - 16$

Sol.

Factorization of $x^3 + 64 = (x)^3 + (4)^3$
 $= (x + 4)[(x)^2 - (x)(4) + (4)^2]$
 $= (x + 4)(x^2 - 4x + 16) \dots\dots\dots (i)$

Factorization of $x^2 - 16 = (x)^2 - (4)^2$
 $= (x + 4)(x - 4) \dots\dots\dots (ii)$

Common factors are: $(x + 4)$

Thus, H.C.F = $x + 4$

Q.8 $x^2 - y^2, x^4 - y^4, x^6 - y^6$

Factorization of $x^2 - y^2 = (x + y)(x - y) \dots\dots\dots (i)$

Factorization of $x^4 - y^4 = (x^2)^2 - (y^2)^2$
 $= (x^2 + y^2)(x^2 - y^2)$
 $= (x^2 + y^2)(x + y)(x - y) \dots\dots\dots (ii)$

Factorization of $x^6 - y^6 = (x^3)^2 - (y^3)^2$
 $= (x^3 - y^3)(x^3 + y^3)$
 $= (x - y)(x^2 + xy + y^2)(x + y)(x^2 - xy + y^2) \dots\dots\dots (iii)$

Common factors are: $(x + y), (x - y)$

Thus, H.C.F = $(x + y)(x - y)$
 $= x^2 - y^2$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

68

Q.9 $t^2 - 9, (t + 3)^2, t^2 + t - 6$

Sol.

Factorization of $t^2 - 9 = (t)^2 - (3)^2$
 $= (t + 3)(t - 3)$(i)

Factorization of $(t + 3)^2 = (t + 3)(t + 3)$(ii)

Factorization of $t^2 + t - 6 = t^2 + 3t - 2t - 6$
 $= t(t + 3) - 2(t + 3)$
 $= (t + 3)(t - 2)$(iii)

Common factor is : $(t + 3)$

Thus, H.C.F = $(t + 3)$

Q.10 $x^2 - x - 2, x^2 + x - 6, x^2 - 3x + 2$

Sol.

Factorization of $x^2 - x - 2 = x^2 - 2x + x - 2$
 $= (x^2 - 2x) + (x - 2)$
 $= x(x - 2) + 1(x - 2)$
 $= (x - 2)(x + 1)$(i)

Factorization of $x^2 + x - 6 = x^2 + 3x - 2x - 6$
 $= (x^2 + 3x) - (2x + 6)$
 $= x(x + 3) - 2(x + 3)$
 $= (x + 3)(x - 2)$(ii)

Factorization of $x^2 - 3x + 2 = x^2 - x - 2x + 2$
 $= (x^2 - x) - (2x - 2)$
 $= x(x - 1) - 2(x - 1)$
 $= (x - 1)(x - 2)$(iii)

Common factor is : $(x - 2)$

Thus, H.C.F = $(x - 2)$

Q.11 $1 - x^2, x^3 + 1, 1 - x - 2x^2$

Sol.

Factorization of $1 - x^2 = (1)^2 - (x)^2$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

69

$$= (1 + x)(1 - x) \dots (i)$$

Factorization of $x^3 + 1 = (x)^3 + (1)^3$

$$= (x + 1)(x^2 - x + 1) \dots (ii)$$

Factorization of $1 - x - 2x^2 = 1 - 2x + x - 2x^2$

$$= (1 - 2x) + (x - 2x^2)$$

$$= 1(1 - 2x) + x(1 - 2x)$$

$$= (1 - 2x)(1 + x)$$

Common factor is $1 + x$

Thus, H.C.F = $1 + x$

Q.12 $x^3 - 8, x^2 - 7x + 10$

Sol:

Factorization of $x^3 - 8 = (x)^3 - (2)^3$

$$= (x - 2)[(x)^2 + (x)(2) + (2)^2]$$

$$= (x - 2)(x^2 + 2x + 4) \dots (i)$$

Factorization of $x^2 - 7x + 10 = x^2 - 2x - 5x + 10$

$$= (x^2 - 2x) - (5x - 10)$$

$$= x(x - 2) - 5(x - 2)$$

$$= (x - 2)(x - 5) \dots (ii)$$

Common factors are: $x - 2$

Thus, H.C.F = $x - 2$

Q.13 $x^2 + 3x + 2, x^2 + 4x + 3, x^2 + 5x + 4$

Sol:

Factorization of $x^2 + 3x + 2 = x^2 + x + 2x + 2$

$$= (x^2 + x) + (2x + 2)$$

$$= x(x + 1) + 2(x + 1)$$

$$= (x + 1)(x + 2) \dots (i)$$

Factorization of $x^2 + 4x + 3 = x^2 + x + 3x + 3$

$$= (x^2 + x) + (3x + 3)$$

$$= x(x + 1) + 3(x + 1)$$

$$= (x + 1)(x + 3) \dots (ii)$$

Factorization of $x^2 + 5x + 4 = x^2 + x + 4x + 4$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

70

$$\begin{aligned}
 &= (x^2 + x) + (4x + 4) \\
 &= x(x + 1) + 4(x + 1) \\
 &= (x + 1)(x + 4) \dots\dots\dots(iii)
 \end{aligned}$$

Common factors are: $(x + 1)$

Thus, H.C.F = $x + 1$

Q.14: $x^4 + x^3 - 6x^2$, $x^4 - 9x^2$, $x^3 + x^2 - 6x$

Sol:

Factorization of $x^4 + x^3 - 6x^2 = x^2(x^2 + x - 6)$

$$\begin{aligned}
 &= x^2(x^2 + 3x - 2x - 6) \\
 &= x^2[(x^2 + 3x) - (2x + 6)] \\
 &= x^2[x(x + 3) - 2(x + 3)] \\
 &= x^2(x + 3)(x - 2) \\
 &= x \times x(x + 3)(x - 2) \dots\dots\dots(i)
 \end{aligned}$$

Factorization of $x^4 - 9x^2 = x^2(x^2 - 9)$

$$\begin{aligned}
 &= x^2[(x)^2 - (3)^2] \\
 &= x^2(x - 3)(x + 3) \\
 &= x \times x(x - 3)(x + 3) \dots\dots\dots(ii)
 \end{aligned}$$

Factorization of $x^3 + x^2 - 6x = x(x^2 + x - 6)$

$$\begin{aligned}
 &= x(x^2 + 3x - 2x - 6) \\
 &= x[(x^2 + 3x) - (2x + 6)] \\
 &= x[x(x + 3) - 2(x + 3)] \\
 &= x(x + 3)(x - 2) \dots\dots\dots(iii)
 \end{aligned}$$

Common factors are: $x, x + 3$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

• Pilot Super One "General Math" 10th

71

Thus, H.C.F = $x(x + 3)$

= $x^2 + 3x$

Q.15 $35a^2c^3h$, $45a^3cb^2$, $30ac^2b^3$

Sol:

Factorization of $35a^2c^3h = 5 \times 7 \times \begin{bmatrix} a \\ a \\ a \end{bmatrix} \times \begin{bmatrix} c \\ c \\ c \end{bmatrix} \times \begin{bmatrix} h \\ h \\ h \end{bmatrix}$
 Factorization of $45a^3cb^2 = 5 \times 3 \times 3 \times \begin{bmatrix} a \\ a \\ a \end{bmatrix} \times \begin{bmatrix} c \\ c \\ c \end{bmatrix} \times \begin{bmatrix} b \\ b \\ b \end{bmatrix}$
 Factorization of $30ac^2b^3 = 5 \times 2 \times 3 \times \begin{bmatrix} a \\ a \\ a \end{bmatrix} \times \begin{bmatrix} c \\ c \\ c \end{bmatrix} \times \begin{bmatrix} b \\ b \\ b \end{bmatrix}$

Common factors = 5, a, b, c

Thus, H.C.F = $5 \times a \times b \times c$
 = $5abc$



Find the H.C.F by Division Method.

Q.1 $x^4 + x^2 + 1$, $x^4 + x^3 + x + 1$

Sol:

$$\begin{array}{r} x^4 + x^2 + 1 \overline{) x^4 + x^3 + x + 1} \\ \underline{+ x^3 + x + 1} \\ x^3 - x^2 + x + 1 \end{array}$$

Take common x

$$\begin{array}{r} x^3 - x^2 + x + 1 \overline{) x^4 + x^3 + x + 1} \\ \underline{+ x^4 - x^3 + x^2 + 1} \\ x^3 - x^2 + x + 1 \end{array}$$

$$\begin{array}{r} x^3 - x^2 + x + 1 \overline{) x^3 - x^2 + x + 1} \\ \underline{+ x^3 - x^2 + x + 1} \\ 0 \end{array}$$

H.C.F = $x^2 - x + 1$

Sold:

$$\begin{array}{r} 3x-1 \\ 2x^2+3x-2 \overline{) 6x^3+7x^2-9x+2} \\ \underline{+6x^3+9x^2-6x} \\ -2x^2-3x+2 \\ \underline{+2x^2+3x-2} \\ 0 \end{array}$$

Q.3 $4x^3 + 2x^2 - 6x$, $4x^3 - 8x + 4$

Sol: $\frac{2 \sqrt{4x^3 + 2x^2 - 6x}}{2x^3 + x^2 - 3x} \quad 4 \sqrt{4x^3 - 8x + 4}$
 $\frac{2 \sqrt{4x^3 + 2x^2 - 6x}}{2x^3 + x^2 - 3x} \quad \frac{4 \sqrt{4x^3 - 8x + 4}}{x^3 - 2x + 1}$

Now we find H.C.F of $x^3 - 2x + 1$ and $2x^3 + x^2 - 3x$ and

H.C.F of 2, 4 is 2.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

73

$$\begin{array}{r} 2 \\ x^3 - 2x + 1 \overline{) 2x^3 + x^2 - 3x} \\ \underline{\pm 2x^3 \quad \mp 4x \pm 2} \\ x^2 + x - 2 \end{array} \quad \begin{array}{r} x^3 - 2x + 1 \\ \underline{\pm x^3 \mp 2x \pm x^2} \\ -x^2 + 1 \end{array}$$

$$\begin{array}{r} -1 \\ -x^2 + 1 \overline{) x^2 + x - 2} \\ \underline{\pm x^2 \mp 1} \\ x - 1 \end{array} \quad \begin{array}{r} -x^2 + 1 \\ \underline{\mp x^2 \pm x} \\ -x + 1 \\ \underline{\mp x \pm 1} \\ 0 \end{array}$$

Required H.C.F = $2(x - 1)$

Q.4 $x^3 + 7x^2 + 12x$, $x^3 + 2x^2 - 15x$

Sol:

$$x \mid x^3 + 7x^2 + 12x$$

$$x \mid x^3 + 2x^2 - 15x$$

$$\begin{array}{r} x^3 + 7x^2 + 12x \\ \hline x^2 + 7x + 12 \end{array}$$

$$\begin{array}{r} x^3 + 2x^2 - 15x \\ \hline x^2 - 2x - 15 \end{array}$$

H.C.F of x and x is x .

Now we find H.C.F of $x^2 - 2x - 15$ and $x^2 + 7x + 12$.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One 'General Math' 10th

74

$$\begin{array}{r}
 x^2 + 7x + 12 \overline{) \begin{array}{l} x^2 - 2x - 15 \\ + x^2 + 7x + 12 \\ \hline - 9 \quad - 9x - 27 \\ + 9x + 27 \\ \hline 0 \end{array}} \\
 \hline
 x + 3 \overline{) \begin{array}{l} x^2 + 7x + 12 \\ + x^2 + 3x \\ \hline + 4x + 12 \\ + 4x + 12 \\ \hline 0 \end{array}} \\
 \hline
 x + 4
 \end{array}$$

Required H.C.F = $x(x + 3)$

Q.5 $x^3 - x^2 - x + 1$, $x^4 - 2x^3 + 2x - 1$

Sol:

$$\begin{array}{r}
 x^3 - x^2 - x + 1 \overline{) \begin{array}{l} x^4 - 2x^3 + 2x - 1 \\ + x^4 - x^3 + x^2 + x \\ \hline - x^3 + x^2 + x - 1 \\ + x^3 + x^2 + x + 1 \\ \hline 0 \end{array}} \\
 \hline
 x - 1
 \end{array}$$

Required H.C.F = $x^3 - x^2 - x + 1$

$$= x^2(x - 1) - 1(x - 1)$$

$$= (x^2 - 1)(x - 1)$$

$$= [(x)^2 - (1)^2](x - 1)$$

$$= (x - 1)(x + 1)(x - 1)$$

$$= (x + 1)(x - 1)^2$$

Q.6 $x^3 - x^2 - x - 2$, $x^3 + 3x^2 - 6x - 8$

Sol:

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

75

$$\begin{array}{r}
 x^3 - x^2 - x - 2 \overline{) x^3 + 3x^2 - 6x - 8} \\
 \underline{\pm x^3 \mp x^2 \mp x \mp 2} \\
 4x^2 - 5x - 6 \overline{) x^3 - x^2 - x - 2}
 \end{array}$$

$$\begin{array}{r}
 4x^2 - 5x - 6 \overline{) 4x^3 - 4x^2 - 4x - 8} \quad \begin{array}{l} x \\ 4 \end{array} \\
 \underline{\pm 4x^3 \mp 5x^2 \mp 6x} \\
 x^2 + 2x - 8 \overline{) 4x^2 - 5x - 6} \\
 \underline{\pm 4x^2 \pm 8x \pm 32} \\
 -13 \overline{) -13x + 26} \\
 \underline{x - 2}
 \end{array}$$

$$\begin{array}{r}
 x - 2 \overline{) x^2 + 2x - 8} \\
 \underline{\pm x^2 \mp 2x} \\
 4x - 8 \\
 \underline{\pm 4x \mp 8} \\
 0
 \end{array}$$

Required H.C.F = $x - 2$

Q.7 $x^2 + 3x - 4$, $x^3 - 2x^2 - 2x + 3$

Sol:

$$\begin{array}{r}
 x^2 + 3x - 4 \overline{) x^3 - 2x^2 - 2x + 3} \quad \begin{array}{l} x - 5 \\ x - 1 \end{array} \\
 \underline{\pm x^3 \pm 3x^2 \mp 4x} \\
 -5x^2 + 2x + 3 \\
 \underline{\mp 5x^2 \mp 15x \pm 20} \\
 17 \overline{) 17x - 17} \\
 x - 1 \overline{) x^2 + 3x - 4} \quad \begin{array}{l} x + 4 \\ 4x - 4 \end{array} \\
 \underline{\pm x^2 \mp x} \\
 4x - 4 \\
 \underline{\pm 4x \mp 4} \\
 0
 \end{array}$$

Required H.C.F = $x - 1$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

76

Q.8 $3x^3 - 14x^2 + 9x + 10$, $15x^3 - 34x^2 + 21x - 10$

Sol:

$$\begin{array}{r}
 \begin{array}{r}
 5 \\
 3x^3 - 14x^2 + 9x + 10 \overline{) 15x^3 - 34x^2 + 21x - 10} \\
 \underline{\pm 15x^3 \mp 70x^2 \pm 45x \pm 50} \\
 12 \overline{) 36x^2 - 24x - 60} \\
 \underline{3x^2 - 2x - 5} \overline{) 3x^3 - 14x^2 + 9x + 10} \\
 \underline{\pm 3x^3 \mp 2x^2 \mp 5x} \\
 -12x^2 + 14x + 10 \\
 \underline{\mp 12x^2 \pm 8x \pm 20} \\
 2 \overline{) 6x - 10} \\
 \underline{3x - 5}
 \end{array}
 \end{array}$$

$a \div b$

$$\begin{array}{r}
 3x - 5 \overline{) 3x^2 - 2x - 5} \\
 \underline{\pm 3x^2 \mp 5x} \\
 3x - 5 \\
 \underline{\pm 3x \mp 5} \\
 0
 \end{array}$$

Required H.C.F = $3x - 5$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

77

Q.9 $2x^4 + x^3 + 4x + 2$, $6x^3 + 5x^2 + x$, $2x^4 + 3x^3 + x^2 + 2x + 1$

Sol: First we will find the H.C.F of $2x^4 + x^3 + 4x + 2$ and $2x^4 + 3x^3 + x^2 + 2x + 1$

$$\begin{array}{r}
 2x^4 + x^3 + 4x + 2 \overline{) 2x^4 + 3x^3 + x^2 + 2x + 1} \\
 \underline{+ 2x^4 + x^3 } \\
 2x^3 + x^2 - 2x - 1 \\
 \underline{+ 2x^3 + x^3 } \\
 2x^2 + 5x + 2 \\
 \overline{) 2x^3 + x^2 - 2x - 1} \\
 \underline{+ 2x^3 + 5x^2 + 2x} \\
 -4x^2 - 4x - 1 \\
 \underline{+ 4x^2 + 10x + 4} \\
 3 \overline{) 6x + 3} \\
 \underline{3 } \\
 \text{H.C.F} \quad 2x + 1 \overline{) 2x^2 + 5x + 2} \\
 \underline{+ 2x^2 + x} \\
 4x + 2 \\
 \underline{+ 4x + 2} \\
 0
 \end{array}$$

Now we will find H.C.F of $2x + 1$ and $6x^3 + 5x^2 + x$

$$\begin{array}{r}
 3x^2 + x \\
 2x + 1 \overline{) 6x^3 + 5x^2 + x} \\
 \underline{+ 6x^3 + 3x^2} \\
 2x^2 + x \\
 \underline{+ 2x^2 + x} \\
 0
 \end{array}$$

Required H.C.F = $2x + 1$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

78

Q.10 $x^3 + x^2 - 5x + 3$, $x^3 - 7x + 6$, $x^3 + 2x^2 - 2x + 3$

Sol: First we will find the H.C.F of $x^3 - 7x + 6$ and $x^3 + x^2 - 5x + 3$.

$$\begin{array}{r}
 x^3 - 7x + 6 \quad | \quad \begin{array}{l} x^3 + x^2 - 5x + 3 \\ \pm x^3 \quad \mp 7x \pm 6 \end{array} \quad x - 2 \\
 \hline
 x^2 + 2x - 3 \quad | \quad \begin{array}{l} x^3 \quad - 7x + 6 \\ \pm x^3 \pm 2x^2 \mp 3x \end{array} \\
 \hline
 \quad \quad \quad \begin{array}{l} - 2x^2 - 4x + 6 \\ \mp 2x^2 \mp 4x \pm 6 \end{array} \\
 \hline
 \quad \quad \quad 0
 \end{array}$$

Now we will find H.C.F of $x^2 + 2x - 3$ and $x^3 + 2x^2 - 2x + 3$.

$$\begin{array}{r}
 x^2 + 2x - 3 \quad | \quad \begin{array}{l} x^3 + 2x^2 - 2x + 3 \\ \pm x^3 \pm 2x^2 \mp 3x \end{array} \quad x - 1 \\
 \hline
 \quad \quad \quad \begin{array}{l} x + 3 \quad | \quad x^2 + 2x - 3 \\ \pm x^2 \pm 3x \end{array} \\
 \hline
 \quad \quad \quad \begin{array}{l} - x - 3 \\ \mp x \mp 3 \end{array} \\
 \hline
 \quad \quad \quad 0
 \end{array}$$

H.C.F $x + 3$

Least Common Multiple (LCM)

The least common multiple of two or more algebraic expressions is the expression of lowest degree which is divisible by each of them without remainder.

The abbreviation of the words *least common multiple* is **L.C.M.**

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Ptmt Super One "General Math" 10th

79



Find L.C.M by Factorization.

Q.1 $21a^4x^3y$, $35a^2x^4y$, $28a^3xy^4$

Sol:

Factorization of $21a^4x^3y = 3 \times 7 \times a \times a \times a \times a \times x \times x \times x \times y$

Factorization of $35a^2x^4y = 5 \times 7 \times a \times a \times x \times x \times x \times x \times y$

Factorization of $28a^3xy^4 = 2 \times 2 \times 7 \times a \times a \times a \times x \times y \times y \times y \times y$

Product of common factors = $7 \times a \times a \times a \times x \times y \times x \times x$
 $= 7a^3x^3y \dots \dots \dots (i)$

Product of uncommon factors = $3 \times 5 \times 2 \times 2 \times a \times x \times y \times y \times y$
 $= 60axy^3 \dots \dots \dots (ii)$

L.C.M = (i) \times (ii)
 $= 7a^3x^3y \times 60axy^3$
 $= 420a^4x^4y^4$

Q.2 $3a^4b^2c^3$, $5a^2b^3c^5$

Sol:

Fac of $3a^4b^2c^3 = 3 \times a \times a \times a \times a \times b \times b \times c \times c \times c$

Fac of $5a^2b^3c^5 = 5 \times a \times a \times b \times b \times b \times c \times c \times c \times c \times c$

Product of common factors = $a \times a \times b \times b \times c \times c \times c$
 $= a^2b^2c^3 \dots \dots \dots (i)$

Product of uncommon factors = $3 \times 5 \times a \times a \times b \times c \times c$
 $= 15a^2bc^2 \dots \dots \dots (ii)$

L.C.M = (i) \times (ii)
 $= (a^2b^2c^3)(15a^2bc^2)$
 $= 15a^4b^3c^5$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

80

Q.3 $2ab, 3ab, 4ca$

Sol:

$$\begin{aligned} \text{Factorization of } 2ab &= 2 \times a \times b \\ \text{Factorization of } 3ab &= 3 \times a \times b \\ \text{Factorization of } 4ca &= 2 \times 2 \times a \times c \end{aligned}$$

$$\text{Product of common factors} = 2 \times a \times b = 2ab \dots\dots (i)$$

$$\text{Product of uncommon factors} = 3 \times 2 \times c = 6c \dots\dots (ii)$$

$$\begin{aligned} \text{L.C.M} &= (i) \times (ii) \\ &= (2ab)(6c) \\ &= 12abc \end{aligned}$$

Q.4 x^2yz, xy^2z, xyz^2

Sol:

$$\begin{aligned} \text{Factorization of } x^2yz &= x \times x \times y \times z \\ \text{Factorization of } xy^2z &= x \times y \times y \times z \\ \text{Factorization of } xyz^2 &= x \times y \times z \times z \end{aligned}$$

$$\text{Product of common factors} = x \times y \times y \times z = xy^2z \dots\dots (i)$$

$$\text{Product of uncommon factors} = x \times z = xz \dots\dots (ii)$$

$$\begin{aligned} \text{L.C.M} &= (i) \times (ii) \\ &= xy^2z \times xz \\ &= x^2y^2z^2 \end{aligned}$$

Q.5 $p^3q - pq^3, p^5q^2 - p^3q^5$

Sol:

$$\begin{aligned} \text{Factorization of } p^3q - pq^3 &= pq(p^2 - q^2) \\ &= pq(p - q)(p + q) \dots\dots (i) \end{aligned}$$

$$\text{Factorization of } p^5q^2 - p^2q^5 = p^2q^2(p^3 - q^3)$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

81

$$= pq(q(p - q)(p^2 + pq + q^2)) \dots (ii)$$

In (i) and (ii)

$$\text{Product of common factors} = pq(p - q) \dots (iii)$$

$$\text{Product of uncommon factors} = pq(p + q)(p^2 + pq + q^2) \dots (iv)$$

$$\text{L.C.M} = (iii) \times (iv)$$

$$= [pq(p - q)] [pq(p + q)(p^2 + pq + q^2)]$$

$$= p^2 q^2 (p - q)(p + q)(p^2 + pq + q^2)$$

Q. 7 $x^3 + 64, x^2 - 16$

Sol:

$$\text{Factorization of } x^3 + 64 = (x)^3 + (4)^3$$

$$= (x + 4) [(x)^2 - (x)(4) + (4)^2]$$

$$= (x + 4)(x^2 - 4x + 16) \dots (i)$$

$$\text{Factorization of } x^2 - 16 = (x)^2 - (4)^2$$

$$= (x + 4)(x - 4) \dots (ii)$$

In (i) and (ii)

$$\text{Product of common factors} = (x + 4) \dots (iii)$$

$$\text{Product of uncommon factors} = (x - 4)(x^2 - 4x + 16) \dots (iv)$$

$$\text{L.C.M} = (iii) \times (iv)$$

$$= (x + 4)(x - 4)(x^2 - 4x + 16)$$

Q. 10 $x^2 - x - 2, x^2 + x - 6, x^2 - 3x + 2$

Sol:

$$\text{Factorization of } x^2 - x - 2 = x^2 - 2x + x - 2$$

$$= (x^2 - 2x) + (x - 2)$$

$$= x(x - 2) + 1(x - 2)$$

$$= (x - 2)(x + 1) \dots (i)$$

$$\text{Factorization of } x^2 + x - 6 = x^2 + 3x - 2x - 6$$

$$= (x^2 + 3x) - (2x + 6)$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

82

$$= x(x + 3) - 2(x + 3)$$

$$= (x + 3)(x - 2) \dots \dots \dots (ii)$$

Factorization of $x^2 - 3x + 2 = x^2 - x - 2x + 2$

$$= (x^2 - x) - (2x + 2)$$

$$= x(x - 1) - 2(x + 1)$$

$$= (x - 1)(x - 2) \dots \dots \dots (iii)$$

In (i), (ii) and (iii)

Product of common factors = $(x - 2) \dots \dots \dots (iv)$

Product of uncommon factors = $(x + 1)(x + 3)(x - 1) \dots \dots \dots (v)$

$$L.C.M = (iv) \times (v)$$

$$= (x - 2)(x + 1)(x + 3)(x - 1)$$

Q.8 $y^2 - 9, (y + 3)^2, y^2 + y - 6$

Sol:

Factorization of $y^2 - 9 = (y)^2 - (3)^2$

$$= (y + 3)(y - 3) \dots \dots \dots (i)$$

Factorization of $(y + 3)^2 = (y + 3)(y + 3) \dots \dots \dots (ii)$

Factorization of $y^2 + y - 6 = y^2 + 3y - 2y - 6$

$$= (y^2 + 3y) - (2y + 6)$$

$$= y(y + 3) - 2(y + 3)$$

$$= (y + 3)(y - 2) \dots \dots \dots (iii)$$

In (i), (ii) and (iii)

Product of common factors = $(y + 3) \dots \dots \dots (iv)$

Product of uncommon factors = $(y - 3)(y + 3)(y - 2) \dots \dots \dots (v)$

$$L.C.M = (iv) \times (v)$$

$$= (y + 3)(y - 3)(y + 3)(y - 2)$$

Q.9 $1 - x^2, x^3 + 1, 1 - x - 2x^3$

Sol:

Factorization of $1 - x^2 = (1)^2 - (x)^2$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

83

$$= (1 + y)(1 - y) \dots (i)$$

Factorization of $y^3 + 1 = (y)^3 + (1)^3$
 $= (y + 1)(y^2 - y + 1) \dots (ii)$

Factorization of $1 - 2y^2 = 1 - 2y + y - 2y^2$
 $= (1 - 2y) + (y - 2y^2)$
 $= 1(1 - 2y) + y(1 - 2y)$
 $= (1 - 2y)(1 + y) \dots (iii)$

In (i), (ii) and (iii)

Product of common factors = $(1 + y) \dots (iv)$

Product of uncommon factors = $(1 - y)(1 - 2y)(y^2 - y + 1) \dots (v)$

Q. 10 $x^2 - y^2, x^4 - y^4, x^6 - y^6$
 L.C.M = $(iv) \times (v)$
 $= (1 + y)(1 - y)(1 - 2y)(y^2 - y + 1)$

Q.10 $x^2 - y^2, x^4 - y^4, x^6 - y^6$

Sol:

Factorization of $x^2 - y^2 = (x + y)(x - y) \dots (i)$
 $= (x)^2 - (y)^2$

Factorization of $x^4 - y^4 = (x^2)^2 - (y^2)^2$
 $= (x^2 + y^2)(x^2 - y^2)$
 $= (x^2 + y^2)[(x)^2 - (y)^2]$
 $= (x^2 + y^2)(x + y)(x - y) \dots (ii)$

Factorization of $x^6 - y^6 = (x^3)^2 - (y^3)^2$
 $= (x^3 + y^3)(x^3 - y^3)$
 $= (x + y)(x^2 - xy + y^2)(x - y)(x^2 + xy + y^2) \dots (iii)$

In (i), (ii) and (iii)

Product of common factors = $(x + y)(x - y) \dots (iv)$

Product of uncommon factors = $(x^2 + y^2)(x^2 - xy + y^2)(x^2 + xy + y^2) \dots (v)$

L.C.M = $(iv) \times (v)$

$= (x + y)(x - y)(x^2 + y^2)$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

84

$$\begin{aligned} & (x^2 - xy + y^2)(x^2 + xy + y^2) \\ & = (x + y)(x - y)(x^2 + y^2) \\ & (x^4 + x^2y^2 + y^4) \end{aligned}$$

Q.11 $x^3 + 1, x^4 + x^2 + 1, (x^2 + x + 1)^2$

Sol:

Factorization of $x^3 + 1 = (x)^3 + (1)^3$
 $= (x + 1)(x^2 - x + 1) \dots\dots\dots(i)$

Factorization of $x^4 + x^2 + 1 = x^4 + 2x^2 + 1 - x^2$ (completing square)
 $= (x^2 + 1)^2 - (x)^2$
 $= (x^2 + 1 + x)(x^2 + 1 - x)$
 $= (x^2 + x + 1)(x^2 - x + 1) \dots\dots(ii)$

Factorization of $(x^2 + x + 1)^2 = (x^2 + x + 1)(x^2 + x + 1) \dots\dots(iii)$

In (i), (ii) and (iii)

Product of common factors $= (x^2 + x + 1)(x^2 - x + 1) \dots(iv)$

Product of uncommon factors $= (x + 1)(x^2 + x + 1) \dots(v)$

L.C.M $= (iv) \times (v)$

$$\begin{aligned} & = (x^2 + x + 1)(x^2 - x + 1)(x + 1)(x^2 + x + 1) \\ & = (x + 1)(x^2 - x + 1)(x^2 + x + 1)^2 \end{aligned}$$

Q.12 $x^3 + y^3, x^4 - y^4, x^6 + y^6$

Sol:

Factorization of $x^3 + y^3 = (x + y)(x^2 - xy + y^2) \dots\dots(i)$

Factorization of $x^4 - y^4 = (x^2)^2 - (y^2)^2$
 $= (x^2 + y^2)(x^2 - y^2)$
 $= (x^2 + y^2)[(x^2) - (y^2)]$
 $= (x^2 + y^2)(x + y)(x - y) \dots\dots(ii)$

Factorization of $x^6 + y^6 = (x^2)^3 + (y^2)^3$
 $= (x^2 + y^2)[(x^2)^2 - (x^2)(y^2) + (y^2)^2]$
 $= (x^2 + y^2)(x^4 - x^2y^2 + y^4) \dots\dots(iii)$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

85

In (i), (ii) and (iii)

$$\text{Product of common factors} = (x + y)(x^2 + y^2) \dots (iv)$$

$$\text{Product of uncommon factors} = (x^2 - xy + y^2)(x - y)(x^4 - x^2y^2 + y^4) \dots (v)$$

$$\text{L.C.M} = (iv) \times (v)$$

$$= (x + y)(x^2 + y^2)(x^2 - x + y^2)(x - y)(x^4 - x^2y^2 + y^4)$$

$$= (x + y)(x - y)(x^2 + y^2)(x^2 - xy + y^2)(x^4 - x^2y^2 + y^4)$$

$$\text{Q.13 } 2x^2 + 5x + 3, x^2 + 2x + 1, 2x^2 + 9x + 9$$

Sol:

$$\text{Factorization of } 2x^2 + 5x + 3 = 2x^2 + 2x + 3x + 3$$

$$= (2x^2 + 2x) + (3x + 3)$$

$$= 2x(x + 1) + 3(x + 1)$$

$$= (x + 1)(2x + 3) \dots (i)$$

$$\text{Factorization of } x^2 + 2x + 1 = x^2 + x + x + 1$$

$$= (x^2 + x) + (x + 1)$$

$$= x(x + 1) + 1(x + 1)$$

$$= (x + 1)(x + 1) \dots (ii)$$

$$\text{Factorization of } 2x^2 + 9x + 9 = 2x^2 + 3x + 6x + 9$$

$$= (2x^2 + 3x) + (6x + 9)$$

$$= x(2x + 3) + 3(2x + 3)$$

$$= (2x + 3)(x + 3) \dots (iii)$$

In (i), (ii) and (iii)

$$\text{Product of common factors} = (x + 1)(2x + 3) \dots (iv)$$

$$\text{Product of uncommon factors} = (x + 1)(x + 3) \dots (v)$$

$$\text{L.C.M} = (iv) \times (v)$$

$$= (x + 1)(2x + 3)(x + 1)(x + 3)$$

$$= (x + 1)^2(2x + 3)(x + 3)$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

86

Q.14 $x^4 + x^3 - 6x^2$, $x^4 - 9x^2$, $x^3 + x^2 - 6x$

Sol:

$$\begin{aligned}\text{Factorization of } x^4 + x^3 - 6x^2 &= x^2(x^2 + x - 6) \\ &= x^2(x^2 + 3x - 2x - 6) \\ &= x^2[(x^2 + 3x) - (2x + 6)] \\ &= x^2[x(x + 3) - 2(x + 3)] \\ &= x^2(x + 3)(x - 2) \dots \dots \dots (i)\end{aligned}$$

$$\begin{aligned}\text{Factorization of } x^4 - 9x^2 &= x^2(x^2 - 9) \\ &= x^2[(x^2 - 3^2)] \\ &= x^2(x + 3)(x - 3) \dots \dots \dots (ii)\end{aligned}$$

$$\begin{aligned}\text{Factorization of } x^3 + x^2 - 6x &= x(x^2 + x - 6) \\ &= x(x^2 + 3x - 2x - 6) \\ &= x[x(x + 3) - 2(x + 3)] \\ &= x(x + 3)(x - 2) \dots \dots \dots (iii)\end{aligned}$$

In (i), (ii) and (iii)

$$\text{Product of common factors} = x^2(x + 3)(x - 2) \dots \dots \dots (iv)$$

$$\text{Product of uncommon factors} = (x - 3) \dots \dots \dots (v)$$

$$\begin{aligned}\text{L.C.M} &= (iv) \times (v) \\ &= x^2(x + 3)(x - 2)(x - 3)\end{aligned}$$

Q.15 $x^2 + 4xy + 4y^2$, $x^2 + 3xy + 2y^2$, $x^2 + 2xy + y^2$

Sol:

$$\begin{aligned}\text{Factorization of } x^2 + 4xy + 4y^2 &= x^2 + 2xy + 2xy + 4y^2 \\ &= (x^2 + 2xy) + (2xy + 4y^2) \\ &= x(x + 2y) + 2y(x + 2y) \\ &= (x + 2y)(x + 2y) \dots \dots \dots (i)\end{aligned}$$

$$\begin{aligned}\text{Factorization of } x^2 + 3xy + 2y^2 &= x^2 + xy + 2xy + 2y^2 \\ &= (x^2 + xy) + (2xy + 2y^2) \\ &= x(x + y) + 2y(x + y) \\ &= (x + y)(x + 2y) \dots \dots \dots (ii)\end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

87

$$\begin{aligned}\text{Factorization of } x^2 + 2xy + y^2 &= x^2 + xy + xy + y^2 \\ &= (x^2 + xy) + (xy + y^2) \\ &= x(x + y) + y(x + y) \\ &= (x + y)(x + y) \dots \dots \dots (iii)\end{aligned}$$

In (i), (ii) and (iii)

$$\text{Product of common factors} = (x + 2y)(x + y) \dots \dots \dots (iv)$$

$$\text{Product of uncommon factors} = (x + 2y)(x + y) \dots \dots \dots (v)$$

$$\begin{aligned}\text{L.C.M} &= (iv) \times (v) \\ &= (x + 2y)(x + y)(x + 2y)(x + y) \\ &= (x + y)(x + y)(x + 2y)(x + 2y) \\ &= (x + y)^2(x + 2y)^2\end{aligned}$$

Relationship between HCF and LCM

If A and B are two algebraic expressions and *H.C.F* and *L.C.M* of these is represented by H and L respectively, then the relation among them can be expressed as:

$$\boxed{A \times B = H \times L}$$

It is called a formula between *L.C.M* and *H.C.F*.

PROOF: Suppose that

$$\frac{A}{H} = x \quad \text{and} \quad \frac{B}{H} = y$$

$$A = Hx \quad \dots \dots \dots (i)$$

$$B = Hy \quad \dots \dots \dots (ii)$$

Since there is no common factor between x and y.

$$\text{Therefore } L = H \cdot x \cdot y$$

$$\begin{aligned}HL &= H(H \cdot x \cdot y) \text{ (multiplying both the sides by H)} \\ &= (Hx) \cdot (Hy) \\ HL &= A \cdot B.\end{aligned}$$

$$(i) \quad L = \frac{A \times B}{H}$$

$$(ii) \quad H = \frac{A \times B}{L}$$

$$(iii) \quad A = \frac{H \times L}{B}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

88

Exercise 3.2

Find the H.C.F and L.C.M of the following.

Q.1 $x^3 + x^2 + x + 1, x^3 - x^2 + x - 1$

Sol:

First we find H.C.F

$$\begin{array}{r}
 \overline{1} \\
 x^3 + x^2 + x + 1 \overline{) x^3 - x^2 + x - 1} \\
 \underline{+ x^3 + x^2 + x + 1} \\
 -2x^2 - 2 \\
 \underline{-2x^2 - 2} \\
 x + 1 \\
 \overline{x^2 + 1} \overline{) x^3 + x^2 + x + 1} \\
 \underline{+ x^3} \underline{+ x} \\
 x^2 + 1 \\
 \underline{+ x^2} \underline{+ 1} \\
 0
 \end{array}$$

H.C.F = $x^2 + 1$

Now we will find L.C.M

$$\begin{aligned}
 \text{L.C.M} &= \frac{(x^3 + x^2 + x + 1)(x^3 - x^2 + x - 1)}{(x^2 + 1)} \\
 &= (x^3 + x^2 + x + 1)(x - 1) \\
 &= x^4 + x^3 + x^2 + x - x^3 - x^2 - x - 1 \\
 \text{L.C.M} &= x^4 - 1
 \end{aligned}$$

Working
 $x - 1$

$$\begin{array}{r}
 x^2 + 1 \overline{) x^3 - x^2 + x - 1} \\
 \underline{+ x^3} \underline{+ x} \\
 -x^2 - 1 \\
 \underline{+ x^2} \underline{+ 1} \\
 0
 \end{array}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

89

Q.2 $x^3 - 3x^2 - 4x + 12$, $x^3 - x^2 - 4x + 4$

Sol: First we find H.C.F

$$\begin{array}{r}
 x^3 - 3x^2 - 4x + 12 \quad \begin{array}{l} \overline{) x^3 - x^2 - 4x + 4} \\ \underline{\pm x^3 \mp 3x^2 \mp 4x \pm 12} \\ 2x^2 - 8 \end{array} \\
 \begin{array}{l} \overline{) 2x^2 - 8} \quad x - 3 \\ \underline{x^2 - 4} \end{array} \quad \begin{array}{l} \overline{) x^3 - 3x^2 - 4x + 12} \\ \underline{\pm x^3 \mp 4x} \\ -3x^2 \quad + 12 \\ \underline{\mp 3x^2 \quad \pm 12} \\ 0 \end{array}
 \end{array}$$

H.C.F = $x^2 - 4$

Now we will find L.C.M

$$\text{L.C.M} = \frac{(x^3 - 3x^2 - 4x + 12)(x^3 - x^2 - 4x + 4)}{(x^2 - 4)}$$

= $(x - 3)(x^3 - x^2 - 4x + 4)$

= $x^4 - x^3 - 4x^2 + 4x - 3x^3 + 3x^2 + 12x - 12$

= $x^4 - 4x^3 - x^2 + 16x - 12$

Q.3 $2x^3 + 2x^2 + x + 1$, $2x^3 - 2x^2 + x - 1$

Sol: First we find H.C.F

$$\begin{array}{r}
 2x^3 + 2x^2 + x + 1 \quad \begin{array}{l} \overline{) 2x^3 - 2x^2 + x - 1} \\ \underline{\pm 2x^3 \pm 2x^2 \pm x \pm 1} \\ -2x^2 - 2 \end{array} \\
 \begin{array}{l} \overline{) -2x^2 - 2} \quad x + 1 \\ \underline{-2x^2 - 2} \\ 0 \end{array}
 \end{array}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

90

$$\begin{array}{r|l}
 2x^2 + 1 & 2x^3 + 2x^2 + x + 1 \\
 & \pm 2x^3 \quad \pm x \\
 \hline
 & 2x^2 \quad + 1 \\
 & \pm 2x^2 \quad \pm 1 \\
 \hline
 & 0
 \end{array}$$

$$\text{H.C.F} = 2x^2 + 1$$

Now we will find L.C.M

$$\begin{array}{l}
 x+1 \\
 \text{L.C.M} = \frac{(2x^3 + 2x^2 + x + 1)(2x^3 - 2x^2 + x - 1)}{(2x^2 + 1)}
 \end{array}$$

$$= (x+1)(2x^3 - 2x^2 + x - 1)$$

$$= 2x^4 - 2x^3 + x^2 - x + 2x^3 - 2x^2 + x - 1$$

$$= 2x^4 - x^2 - 1$$

Q.4 $6x^3 + 7x^2 - 9x + 2, 8x^4 + 6x^3 - 15x^2 + 9x - 2$

Sol:

First we find H.C.F

$$4x + 5$$

$$\begin{array}{r|l}
 6x^3 + 7x^2 - 9x + 2 & 8x^4 + 6x^3 - 15x^2 + 9x - 2 \\
 & \times 3 \\
 \hline
 & 24x^4 + 18x^3 - 45x^2 + 27x - 6 \\
 & \pm 24x^4 \pm 28x^3 \mp 36x^2 \pm 8x \\
 \hline
 & -10x^3 - 9x^2 + 19x - 6 \\
 & -1 \mid -10x^3 - 9x^2 + 19x - 6 \\
 \hline
 & 10x^3 + 9x^2 - 19x + 6 \\
 & \times 3 \\
 \hline
 & 30x^3 + 27x^2 - 57x + 18 \\
 & \pm 30x^3 \pm 35x^2 \mp 45x \pm 10 \\
 \hline
 \end{array}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

91

$$\begin{array}{r}
 -1 \overline{) -8x^2 - 12x + 8} \\
 4 \overline{) 8x^2 + 12x - 8} \qquad 3x - 1 \\
 \hline
 2x^2 + 3x - 2 \overline{) 6x^3 + 7x^2 - 9x + 2} \\
 \underline{\pm 6x^3 \pm 9x^2 \mp 6x} \\
 -2x^2 - 3x + 2 \\
 \underline{\mp 2x^2 \mp 3x \pm 2} \\
 0
 \end{array}$$

$$\text{H.C.F} = 2x^2 + 3x - 2$$

Now we will find L.C.M

$$3x - 1$$

$$\text{L.C.M} = \frac{(6x^3 + 7x^2 - 9x + 2)(8x^4 + 6x^3 - 15x^2 + 9x - 2)}{(2x^2 + 3x - 2)}$$

$$= (3x - 1)(8x^4 + 6x^3 - 15x^2 + 9x - 2)$$

Q.5 $3x^4 + 17x^3 + 27x^2 + 7x - 6$, $6x^4 + 7x^3 - 27x^2 + 17x - 3$

Sol: First we find H.C.F

$$\begin{array}{r}
 2 \\
 3x^4 + 17x^3 + 27x^2 + 7x - 6 \overline{) 6x^4 + 7x^3 - 27x^2 + 17x - 3} \\
 \underline{\pm 6x^4 \pm 34x^3 \pm 54x^2 \pm 14x \mp 12} \\
 -3 \overline{) -27x^3 - 81x^2 + 3x + 9} \\
 \underline{9x^3 + 27x^2 - x - 3}
 \end{array}$$

$$\begin{array}{r}
 x + 2 + 2 \\
 9x^3 + 27x^2 - x - 3 \overline{) 3x^4 + 17x^3 + 27x^2 + 7x - 6} \\
 \underline{\times 3} \\
 9x^4 + 51x^3 + 81x^2 + 21x - 18 \\
 \underline{\pm 9x^4 \pm 27x^3 \mp x^2 \mp 3x}
 \end{array}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

92

$$24x^3 + 82x^2 + 24x - 18$$

$$\pm 18x^3 \pm 54x^2 \mp 2x \mp 6$$

$$6x^3 + 28x^2 + 26x - 12$$

$\times 3$

$$18x^3 + 84x^2 + 78x - 36$$

$$\pm 18x^3 \pm 54x^2 \mp 2x \mp 6$$

$$2 \mid 30x^2 + 80x - 30$$

$$5 \mid 15x^2 + 40x - 15$$

$$3x^2 + 8x - 3 \mid 9x^3 + 27x^2 - x - 3$$

$$\pm 9x^3 \pm 24x^2 \mp 9x$$

$$3x^2 + 8x - 3$$

$$\pm 3x^2 \pm 8x \mp 3$$

0

$$\text{H.C.F} = 3x^2 + 8x - 3$$

Now we will find L.C.M

$$2x^2 - 3x + 1$$

$$\text{L.C.M} = \frac{(3x^4 + 17x^3 + 27x^2 + 7x - 6)(6x^4 + 7x^3 - 27x^2 + 17x - 3)}{(3x^2 + 8x - 3)}$$

$$= (2x^2 - 3x + 1)(3x^4 + 17x^3 + 27x^2 + 7x - 6)$$

Working

$$2x^2 - 3x + 1$$

$$3x^2 + 8x - 3$$

$$6x^4 + 7x^3 - 27x^2 + 17x - 3$$

$$\pm 6x^4 \pm 16x^3 \mp 6x^2$$

$$-9x^3 - 21x^2 + 17x - 3$$

$$\mp 9x^3 \mp 24x^2 \pm 9x$$

$$3x^2 + 8x - 3$$

$$\pm 3x^2 \pm 8x \mp 3$$

0

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

93

Q.6 $2x^4 + 3x^3 - 13x^2 - 7x + 15$, $2x^4 + x^3 - 20x^2 - 7x + 24$

Sol: First we find H.C.F

$$\begin{array}{r}
 1 \\
 2x^4 + 3x^3 - 13x^2 - 7x + 15 \overline{) 2x^4 + x^3 - 20x^2 - 7x + 24} \\
 \underline{\pm 2x^4 + 3x^3 \mp 13x^2 \mp 7x \pm 15} \\
 -1 \mid -2x^3 - 7x^2 + 9 \qquad x - 2 \\
 \underline{2x^3 + 7x^2 - 9} \quad 2x^4 + 3x^3 - 13x^2 - 7x + 15 \\
 \underline{\pm 2x^4 \pm 7x^3 \mp 9x} \\
 -4x^3 - 13x^2 + 2x + 15 \\
 \underline{\mp 4x^3 \mp 14x^2 \pm 18} \\
 x^2 + 2x - 3
 \end{array}$$

$$\begin{array}{r}
 2x^2 - x - 5 \\
 x^2 + 2x - 3 \overline{) 2x^4 + 3x^3 - 13x^2 - 7x + 15} \\
 \underline{\pm 2x^4 + 4x^3 \mp 6x^2} \\
 -x^3 - 7x^2 - 7x + 15 \\
 \underline{\mp x^3 \mp 2x^2 \pm 3x} \\
 -5x^2 - 10x + 15 \\
 \underline{\mp 5x^2 \mp 10x \pm 15} \\
 0
 \end{array}$$

H.C.F = $x^2 + 2x - 3$

Now we will find L.C.M

$$\begin{array}{l}
 2x^2 - x - 5 \\
 \text{L.C.M} = \frac{(2x^4 + 3x^3 - 13x^2 - 7x + 15)(2x^4 + x^3 - 20x^2 - 7x + 24)}{x^2 + 2x - 3} \\
 = (2x^2 - x - 5)(2x^4 + x^3 - 20x^2 - 7x + 24)
 \end{array}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

84

Q.7 $x^4 - x^3 - x + 1, x^4 + x^3 - x - 1$

Sol: First we find H.C.F

$$\begin{array}{r}
 \begin{array}{r}
 1 \\
 x^4 - x^3 - x + 1 \overline{) x^4 + x^3 - x - 1} \\
 \underline{\pm x^4 \mp x^3 \mp x \pm 1} \\
 2x^3 - 2
 \end{array}
 \qquad
 \begin{array}{r}
 x - 1 \\
 x^3 - 1 \overline{) x^4 - x^3 - x + 1} \\
 \underline{\pm x^4 \mp x} \\
 -x^3 + 1 \\
 \underline{-x^3 + 1} \\
 0
 \end{array}
 \end{array}$$

H.C.F = $x^3 - 1$

Now we will find L.C.M

$(x-1)$

$$\begin{aligned}
 \text{L.C.M} &= \frac{(x^4 - x^3 - x + 1)(x^4 + x^3 - x - 1)}{(x^3 - 1)} \\
 &= (x-1)(x^4 + x^3 - x - 1)
 \end{aligned}$$

Q.8 $x^4 + x^3 + x + 1, x^4 + x^3 - x - 1$

Sol: First we find H.C.F

$$\begin{array}{r}
 \begin{array}{r}
 1 \\
 x^4 + x^3 + x + 1 \overline{) x^4 + x^3 - x - 1} \\
 \underline{\pm x^4 \pm x^3 \pm x \pm 1} \\
 -2x - 2
 \end{array}
 \qquad
 \begin{array}{r}
 x^3 + 1 \\
 x + 1 \overline{) x^4 + x^3 + x + 1} \\
 \underline{\pm x^4 \pm x^3} \\
 x + 1 \\
 \underline{\pm x \pm 1} \\
 0
 \end{array}
 \end{array}$$

H.C.F = $x + 1$

Now we will find L.C.M.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

95

$$\begin{aligned} & (x^3 + 1) \\ \text{L.C.M} &= \frac{(x^4 + x^3 + x + 1)(x^4 + x^3 - x - 1)}{(x + 1)} \\ &= (x^3 + 1)(x^4 + x^3 - x + 1) \end{aligned}$$

Find the Required Polynomial

Q.9. $A = x^2 - 5x - 14$, $H = x - 7$, $L = x^3 - 10x^2 + 11x + 70$, $B = ?$

Sol: Formula: $B = \frac{H \times L}{A}$

$$\begin{aligned} & (x - 5) \\ &= \frac{(x - 7)(x^3 - 10x^2 + 11x + 70)}{(x^2 - 5x - 14)} \\ &= (x - 7)(x - 5) \\ &= x^2 - 12x + 35 \end{aligned}$$

Working
 $x - 5$

$$\begin{array}{r} x^2 - 5x - 14 \overline{) x^3 - 10x^2 + 11x + 70} \\ \underline{+ x^3 + 5x^2 + 14x} \\ -5x^2 + 25x + 70 \\ \underline{+ 5x^2 + 25x + 70} \\ 0 \end{array}$$

Q.10. $B = 3x^2 + 14x + 8$, $H = 3x + 2$, $L = 6x^3 + 25x^2 + 2x - 8$,
 $A = ?$

Sol: Formula: $A = \frac{H \times L}{B}$

$$\begin{aligned} & (2x - 1) \\ &= \frac{(3x + 2)(6x^3 + 25x^2 + 2x - 8)}{(3x^2 + 14x + 8)} \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

96

$$= (3x + 2)(2x - 1)$$

$$= 6x^2 + x - 2$$

$$\begin{array}{r}
 \text{Working} \\
 2x - 1 \\
 3x^2 + 14x + 18 \quad \overline{) \quad 6x^3 + 25x^2 + 2x - 8} \\
 \underline{\pm 6x^3 \pm 28x^2 \pm 16x} \\
 -3x^2 - 14x - 8 \\
 \underline{+ 3x^2 + 14x + 8} \\
 0
 \end{array}$$

Q.11. The product of two polynomials and their L.C.M. are $x^4 + 6x^3 - 3x^2 - 56x - 48$ and $x^3 + 2x^2 - 11x - 12$ respectively. Find their H.C.F.

Sol:

$$A \times B = x^4 + 6x^3 - 3x^2 - 56x - 48$$

$$H = x^3 + 2x^2 - 11x - 12$$

$$L = ?$$

$$\text{Formula: } L = \frac{A \times B}{H}$$

$$\begin{aligned}
 &= \frac{x^4 + 6x^3 - 3x^2 - 56x - 48}{x^3 + 2x^2 - 11x - 12} \\
 &= x + 4
 \end{aligned}$$

$$\begin{array}{r}
 \text{Working} \\
 x + 4 \\
 x^3 + 2x^2 - 11x - 12 \quad \overline{) \quad x^4 + 6x^3 - 3x^2 - 56x - 48} \\
 \underline{\pm x^4 \pm 2x^3 \mp 11x^2 \mp 12x} \\
 + 4x^3 + 8x^2 - 44x - 48 \\
 \underline{\pm 4x^3 \pm 8x^2 \mp 44x \mp 48} \\
 0
 \end{array}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

97

Q.12. The product of two polynomials and their L.C.M. are
 $x^4 + 5x^3 - x^2 - 17x + 12$ and $x^3 + 6x^2 + 5x - 12$
 respectively.

Find their H.C.F.

Sol: $A \times B = x^4 + 5x^3 - x^2 - 17x + 12$

$L = x^3 + 6x^2 + 5x - 12$

$H = ?$

$$\text{Formula } H = \frac{A \times B}{L}$$

$$= \frac{x^4 + 5x^3 - x^2 - 17x + 12}{x^3 + 6x^2 + 5x - 12}$$

$H = x - 1$

Working

$$\begin{array}{r} x - 1 \\ x^3 + 6x^2 + 5x - 12 \overline{) x^4 + 5x^3 - x^2 - 17x + 12} \\ \underline{+ x^4 + 6x^3 + 5x^2 - 12x} \\ -x^3 - 6x^2 - 5x + 12 \\ \underline{+ x^3 + 6x^2 + 5x - 12} \\ 0 \end{array}$$

Q.13. The product of two polynomials and their H.C.F. are
 $x^4 - 12x^3 + 53x^2 - 102x + 72$ and $x - 3$ respectively. Find
 L.C.M.

Sol:

$A \times B = x^4 - 12x^3 + 53x^2 - 102x + 72$

$L = x - 3$

$H = ?$

$$\text{Formula } H = \frac{A \times B}{L}$$

$$= \frac{x^4 - 12x^3 + 53x^2 - 102x + 72}{(x - 3)}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

98

$$H = x^3 - 9x^2 + 26x - 24$$

Working

$$\begin{array}{r}
 x^3 - 9x^2 + 26x - 24 \\
 x - 3 \overline{) \begin{array}{l} x^4 - 12x^3 + 53x^2 - 102x + 72 \\ \underline{+ x^4 + 3x^3} \\ - 9x^3 + 53x^2 - 102x + 72 \\ \underline{+ 9x^3 + 27x^2} \\ 26x^2 - 102x + 72 \\ \underline{+ 26x^2 + 78x} \\ - 24x + 72 \\ \underline{+ 24x + 72} \\ 0 \end{array}}
 \end{array}$$

Q.14. The product of two polynomials and their H.C.F. is $x^4 - 5x^3 + 2x^2 + 20x - 24$ and $x + 2$ respectively. Find their L.C.M.

Sol:

$$A \times B = x^4 - 5x^3 + 2x^2 + 20x - 24$$

$$H = x + 2$$

$$L = ?$$

$$\text{Formula } L = \frac{A \times B}{H}$$

$$= \frac{x^4 - 5x^3 + 2x^2 + 20x - 24}{(x + 2)}$$

$$H = x^3 - 7x^2 + 16x - 12$$

Working

$$\begin{array}{r}
 x^3 - 7x^2 + 16x - 12 \\
 x + 2 \overline{) \begin{array}{l} x^4 - 5x^3 + 2x^2 + 20x - 24 \\ \underline{+ x^4 + 2x^3} \\ - 7x^3 + 2x^2 + 20x - 24 \end{array}}
 \end{array}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

99

$$\begin{array}{r}
 \mp 7x^3 \mp 14x^2 \\
 \hline
 + 16x^2 + 20x - 24 \\
 \pm 16x^3 \pm 32x \\
 \hline
 \pm 12x - 24 \\
 \mp 12x \mp 24 \\
 \hline
 0
 \end{array}$$

Q.15. One algebraic expression is $x^3 + 3x^2 - 4x - 12$ and $x^3 + 5x^2 - 4x - 20$ other one is . Their H.C.F is $x^2 - 4$. Find their L.C.M.

Sol:

$$A = x^3 + 3x^2 - 4x - 12$$

$$B = x^3 + 5x^2 - 4x - 20$$

$$H = x^2 - 4$$

$$L = ?$$

$$\text{Formula: } L = \frac{A \times B}{H}$$

$$\frac{(x^3 + 3x^2 - 4x - 12)(x^3 + 5x^2 - 4x - 20)}{(x^2 - 4)}$$

$$\begin{aligned}
 &= (x + 3)(x^2 + 5x^2 - 4x - 20) \\
 &= x^4 + 8x^3 + 11x^2 - 32x - 60
 \end{aligned}$$

Working
 $x + 3$

$$\begin{array}{r}
 x^2 - 4 \overline{) \begin{array}{l} x^3 + 3x^2 - 4x - 12 \\ \pm x^3 \quad \mp 4x \\ \hline 3x^2 \quad - 12 \\ \pm 3x^2 \quad \mp 12 \\ \hline 0 \end{array}}
 \end{array}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

100

Q.16. One algebraic expression is $x^3 - x^2 + 2x - 2$ and other one is $x^3 - x^2 - 2x + 2$. Their H.C.F is $x - 1$. Find their L.C.M.

Sol:

$$A = x^3 - x^2 + 2x - 2$$

$$B = x^3 - x^2 - 2x + 2$$

$$H = x - 1$$

$$L = ?$$

$$\text{Formula: } L = \frac{A \times B}{H}$$

$$(x^2 + 2)$$

$$= \frac{(x^3 - x^2 + 2x - 2)(x^3 - x^2 - 2x + 2)}{(x - 1)}$$

$$= (x^2 + 2)(x^3 - x^2 - 2x + 2)$$

$$= x^5 - x^4 - 2x + 4$$

Working

$$x^2 + 2$$

$$\begin{array}{r} x-1 \overline{) \begin{array}{l} x^3 - x^2 + 2x - 2 \\ \underline{+ x^3 - x^2} \\ 2x^2 - 2 \\ \underline{+ 2x^2 + 2} \\ 0 \end{array}} \end{array}$$

Q.17. Prove that $H^3 + L^3 = A^3 + B^3$ where $H + L = A + B$ 'H' and 'L' stand for H.C.F and L.C.M respectively and 'A, B' represent two polynomials.

Sol:

Proof: We know that _____

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

101

$$H + L = A + B$$

$$(H + L)^3 = (A + B)^3 \quad \text{Taking cube}$$

$$H^3 + L^3 + 3HL(H + L) = A^3 + B^3 + 3AB(A + B)$$

$$H^3 + L^3 = A^3 + B^3 + 3AB(A + B) - 3HL(H + L) \quad (i)$$

Now $H + L = A + B$

and $H \times L = A \times B$

Putting values in (i) $H + L$ and HL

$$H^3 + L^3 = A^3 + B^3 + 3AB(A + B) - 3AB(A + B)$$

$$H^3 + L^3 = A^3 + B^3 \quad \text{Proved}$$

Exercise 3.5

Simplify

Q.1 $\frac{1}{a} + \frac{2}{a+1} - \frac{3}{a+2}$

Sol: $= \frac{1(a+1)(a+2) + 2(a)(a+2) - 3(a)(a+1)}{(a)(a+1)(a+2)}$

$$= \frac{a^2 + 3a + 2 + 2a^2 + 4a - 3a^2 - 3a}{(a)(a+1)(a+2)}$$

$$= \frac{4a + 2}{(a)(a+1)(a+2)}$$

$$= \frac{2(2a+1)}{(a)(a+1)(a+2)}$$

Q.2 $\frac{2a}{(x-2a)} - \frac{x-a}{x^2-5ax+6a^2} + \frac{2}{x-3a}$

Sol: $= \frac{2a}{(x-2a)} - \frac{x-a}{x^2-2ax-3ax+6a^2} + \frac{2}{x-3a}$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

102

$$= \frac{2a}{(x-2a)} - \frac{x-a}{x(x-2a)-3a(ax-2a)} + \frac{2}{x-3a}$$

$$= \frac{2a}{(x-2a)} - \frac{x-a}{(x-2a)(x-3a)} + \frac{2}{x-3a}$$

$$= \frac{2a(x-3a)-(x-a)+2(x-2a)}{(x-2a)(x-3a)}$$

$$= \frac{2ax-6a^2-x+a+2x-4a}{(x-2a)(x-3a)}$$

$$= \frac{2ax+x-3a-6a^2}{(x-2a)(x-3a)}$$

Q.3: $\frac{1}{a^2+1} - \frac{a^4}{a^2+1} + \frac{a^6}{a^2-1} - \frac{1}{a^2-1}$

Sol: $= \frac{1(a^2-1)-a^4(a^2-1)+a^6(a^2+1)-1(a^2+1)}{(a^2+1)(a^2-1)}$

$$= \frac{a^2-1-a^4+a^4+a^6-a^2-1}{(a^2+1)(a^2-1)}$$

$$= \frac{a^6+a^4-2}{(a^2+1)(a^2-1)}$$

$$= \frac{a^4+2-a^4-2}{(a^2+1)(a^2-1)}$$

$$= \frac{a^4(a^2+2)-1(a^2+2)}{(a^2+1)(a^2-1)}$$

$$= \frac{(a^4+2)(a^2-1)}{(a^2+1)(a^2-1)}$$

$$= a^4+2$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

103

$$Q.4 \quad \frac{1}{x^2+x+1} - \frac{1}{x^2-x+1} + \frac{2x+1}{x^4+x^2+1}$$

$$\begin{aligned} \text{Sol:} &= \frac{1}{x^2+x+1} - \frac{1}{x^2-x+1} + \frac{2x+1}{(x^2+x+1)(x^2-x+1)} \\ &= \frac{1(x^2-x+1) - 1(x^2+x+1) + 2x+1}{(x^2+x+1)(x^2-x+1)} \\ &= \frac{x^2-x+1-x^2-x-1+2x+1}{(x^2+x+1)(x^2-x+1)} \end{aligned}$$

$$= \frac{1}{(x^2+x+1)(x^2-x+1)}$$

$$= \frac{1}{(x^4+x^2+1)}$$

$$Q.5 \quad \frac{a^2(b-c)}{(a+b)(a+c)} - \frac{b^2(c-a)}{(b+c)(b+a)} + \frac{c^2(a-b)}{(c+a)(c+b)}$$

$$\text{Sol:} = \frac{a^2(b-c)(b+c) - b^2(c-a)(c+a) + c^2(a-b)(a+b)}{(a+b)(a+c)(b+c)}$$

$$= \frac{a^2(b^2-c^2) - b^2(c^2-a^2) + c^2(a^2-b^2)}{(a+b)(a+c)(b+c)}$$

$$= \frac{a^2b^2 - a^2c^2 - b^2c^2 + b^2a^2 + a^2c^2 - b^2c^2}{(a+b)(a+c)(b+c)}$$

$$= \frac{2a^2b^2 - 2b^2c^2}{(a+b)(a+c)(b+c)}$$

$$= \frac{2b^2(a^2-c^2)}{(a+b)(a+c)(b+c)}$$

$$= \frac{2b^2(a+c)(a-c)}{(a+b)(a+c)(b+c)}$$

$$= \frac{2b^2(a-c)}{(a+b)(b+c)}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

104

Q.6 $\frac{1}{x-1} + \frac{1}{x+1} - \frac{x+2}{x^2+x+1} - \frac{x-2}{x^2-x+1}$

Sol: Changing order

$$\begin{aligned} &= \left(\frac{1}{x-1} - \frac{x+2}{x^2+x+1} \right) + \left(\frac{1}{x+1} - \frac{x-2}{x^2-x+1} \right) \\ &= \frac{(x^2+x+1) - (x-1)(x+2)}{(x-1)(x^2+x+1)} + \frac{(x^2-x+1) - (x-2)(x+1)}{(x+1)(x^2-x+1)} \\ &= \frac{(x^2+x+1) - (x^2+x-2)}{x^3-1} + \frac{(x^2-x+1) - (x^2-x-2)}{x^3+1} \\ &= \frac{x^2+x+1-x^2-x+2}{x^3-1} + \frac{x^2-x+1-x^2+x+2}{x^3+1} \\ &= \frac{3}{x^3-1} + \frac{3}{x^3+1} \\ &= \frac{3(x^3+1) + 3(x^3-1)}{(x^3-1)(x^3+1)} \\ &= \frac{3x^3+3+3x^3-3}{x^6-1} \\ &= \frac{6x^3}{x^6-1} \end{aligned}$$

Q.7 $\frac{a^3+ab+b^2}{a+b} + \frac{a^2-ab+b^2}{a-b}$

Sol:
$$\begin{aligned} &= \frac{(a-b)(a^3+ab+b^2) + (a+b)(a^2-ab+b^2)}{(a+b)(a-b)} \\ &= \frac{(a^3-b^3) + (a^3+b^3)}{a^2-b^2} \\ &= \frac{a^3-b^3+a^3+b^3}{a^2-b^2} \\ &= \frac{2a^3}{a^2-b^2} \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

105

$$Q.8 \quad \frac{x^2 - y^2}{x^2 - 2xy + y^2} \times \frac{x - y}{x(x + y)} + \frac{x^2 + y^2}{x}$$

$$\begin{aligned} \text{Sol:} &= \frac{(x^2 + y^2)(x^2 - y^2)}{(x - y)^2} \times \frac{x - y}{x(x + y)} \times \frac{x}{(x^2 + y^2)} \\ &= \frac{(x + y)(x - y)}{(x - y)(x - y)} \times \frac{(x - y)}{x(x + y)} \times x \\ &= 1 \end{aligned}$$

$$Q.9 \quad \frac{x^2 - 1}{x^2 + x - 2} \times \frac{x^3 + 8}{x^4 + 4x^2 + 16} + \frac{x^2 + x}{x^3 + 2x^2 + 4x}$$

$$\begin{aligned} \text{Sol:} &= \frac{(x + 1)(x - 1)}{x^2 + 2x - x - 2} \times \frac{(x)^3 + (2)^3}{x^4 + 4x^2 + 16} \times \frac{x^3 + 2x^2 + 4x}{x^2 + x} \\ &= \frac{(x + 1)(x - 1)}{x(x + 2) - 1(x + 2)} \times \frac{(x + 2)(x^2 - 2x + 4)}{x^4 + 8x^2 + 16 - 4x^2} \times \frac{x(x^2 + 2x + 4)}{x(x + 1)} \end{aligned}$$

(Completing square)

$$\begin{aligned} &= \frac{(x + 1)(x - 1)}{(x + 2)(x - 1)} \times \frac{(x + 2)(x^2 - 2x + 4)}{(x^2 + 4)^2 - (2x)^2} \times \frac{x^2 + 2x + 4}{x + 1} \\ &= \frac{(x^2 - 2x + 4)(x^2 + 2x + 4)}{(x^2 + 4 - 2x)(x^2 + 4 + 2x)} \\ &= \frac{(x^2 - 2x + 4)(x^2 + 2x + 4)}{(x^2 - 2x + 4)(x^2 + 2x + 4)} \\ &= 1 \end{aligned}$$

$$Q.10 \quad \frac{a^3 + 64b^3}{a^2 + 20ab + 64b^2} + \frac{a^2 - 4ab + 16b^2}{a^2 + 4ab + 16b^2} \times \frac{a^2 + 12ab - 64b^2}{a^3 - 64b^3}$$

$$\begin{aligned} \text{Sol:} &= \frac{(a)^3 + (4b)^3}{a^2 + 4ab + 16ab + 64b^2} \times \frac{a^2 + 4ab + 16b^2}{a^2 - 4ab + 16b^2} \times \frac{a^2 + 12ab - 64b^2}{(a)^3 - (4b)^3} \\ &= \frac{(a + 4b)(a^2 - 4ab + 16b^2)}{a(a + 4b) + 16b(a + 4b)} \times \frac{a^2 + 4ab + 16b^2}{a^2 - 4ab + 16b^2} \times \frac{a(a + 16b) - 4b(a + 16b)}{(a - 4b)(a^3 + 4ab + 16b^3)} \\ &= \frac{(a + 4b)(a^2 - 4ab + 16b^2)}{(a + 4b)(a + 16b)} \times \frac{a^2 + 4ab + 16b^2}{a^2 - 4ab + 16b^2} \times \frac{(a + 16b)(a - 4b)}{(a - 4b)(a^3 + 4ab + 16b^3)} \\ &= 1 \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

106

$$Q.11 \quad \frac{a}{(a+b)^2 - 2ab} \times \frac{a^4 - b^4}{(a+b)^3 - 3ab(a+b)} + \frac{(a+b)^2 - 4ab}{(a+b)^2 - 3ab}$$

$$\text{Sol: } = \frac{a}{a^2 + b^2 + 2ab - 2ab} \times \frac{(a^2)^2 - (b^2)^2}{a^3 + b^3 + 3ab(a+b) - 3ab(a+b)} + \frac{(a+b)^2 - 4ab}{(a+b)^2 - 3ab}$$

$$= \frac{a}{(a^2 + b^2)} \times \frac{(a^2 + b^2)(a^2 - b^2)}{a^3 + b^3} \times \frac{a^2 + b^2 + 2ab - 3ab}{a^2 + b^2 + 2ab - 4ab}$$

$$= a \times \frac{(a+b)(a-b)}{(a+b)(a^2 - ab + b^2)} \times \frac{(a^2 + b^2 - ab)}{a^2 + b^2 - 2ab}$$

$$= \frac{a \times (a-b)}{a^2 - 2ab + b^2} = \frac{a(a-b)}{(a-b)^2}$$

$$= \frac{a(a-b)}{(a-b)(a-b)}$$

$$= \frac{a}{a-b}$$

$$Q.12 \quad \frac{a^2 - 1}{a^2 - a - 2} + \frac{a^2 + 5a + 6}{a^2 - 5a + 6} + \frac{a^2 - 4a + 3}{a^2 + 4a + 3}$$

$$\text{Sol: } = \frac{(a)^2 - (1)^2}{a^2 - 2a + a - 2} \times \frac{a^2 - 5a + 6}{a^2 + 5a + 6} \times \frac{a^2 + 4a + 3}{a^2 - 4a + 3}$$

$$= \frac{(a-1)(a+1)}{a(a-2)+1(a-2)} \times \frac{a^2 - 2a - 3a + 6}{a^2 + 2a + 3a + 6} \times \frac{a^2 + a + 3a + 3}{a^2 - a - 3a + 3}$$

$$= \frac{(a-1)(a+1)}{(a-2)(a+1)} \times \frac{a(a-2)-3(a-2)}{a(a+2)+3(a+2)} \times \frac{a(a+1)+3(a+1)}{a(a-1)-3(a-1)}$$

$$= \frac{(a-1)(a+1)}{(a-2)(a+1)} \times \frac{(a-2)(a-3)}{(a+2)(a+3)} \times \frac{(a+1)(a+3)}{(a-1)(a-3)}$$

$$= \frac{a+1}{a+2}$$

Exercise 5.6

Find the Square Root of the Following.

Q.1 $16x^2 + 24xy + 9y^2$

Sol:
$$\begin{aligned} &= 16x^2 + 12xy + 12xy + 9y^2 \\ &= (16x^2 + 12xy) + (12xy + 9y^2) \\ &= 4x(4x + 3y) + 3y(4x + 3y) \\ &= (4x + 3y)(4x + 3y) \\ &= (4x + 3y)^2 \\ \sqrt{16x^2 + 24xy + 9y^2} &= \sqrt{(4x + 3y)^2} \\ &= \pm(4x + 3y) \end{aligned}$$

Q.2 $(x^2 - 7x + 12)(x^2 - 9x + 20)(x^2 - 8x + 15)$

Sol:
$$\begin{aligned} &= [x^2 - 3x - 4x + 12][x^2 - 4x - 5x + 20][x^2 - 3x - 5x + 15] \\ &= [(x^2 - 3x) - (4x - 12)][(x^2 - 4x) - (5x - 20)][(x^2 - 3x) - (5x - 15)] \\ &= [x(x - 3) - 4(x - 3)][x(x - 4) - 5(x - 4)][x(x - 3) - 5(x - 5)] \\ &= (x - 3)(x - 4)(x - 4)(x - 5)(x - 3)(x - 5) \\ &= (x - 3)^2(x - 4)^2(x - 5)^2 \end{aligned}$$

Now we will take square root.

$$\begin{aligned} &= \sqrt{(x - 3)^2(x - 4)^2(x - 5)^2} \\ &= \pm(x - 3)(x - 4)(x - 5) \end{aligned}$$

Q.3 $(x^2 + 8x + 7)(2x^2 - x - 3)(2x^2 + 11x - 21)$

Sol:
$$\begin{aligned} &= (x^2 + x + 7x + 7)(2x^2 - 3x + 2x - 3)(2x^2 + 14x - 3x - 21) \\ &= [(x^2 + x) + (7x + 7)][(2x^2 - 3x) + (2x - 3)][(2x^2 + 14x) - (3x + 21)] \\ &= [x(x + 1) + 7(x + 1)][x(2x - 3) + 1(2x - 3)][2x(x + 7) - 3(x + 7)] \\ &= [(x + 1)(x + 7)][(2x - 3)(x + 1)][(x + 7)(2x - 3)] \\ &= (x + 1)^2(x + 7)^2(2x - 3)^2 \\ &= \sqrt{(x + 1)^2(x + 7)^2(2x - 3)^2} \text{ (Taking square root)} \\ &= \pm(x + 1)(x + 7)(2x - 3) \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

108

Q.4 $x(x+2)(x+4)(x+6) + 16$

Sol: $= x(x+6)(x+2)(x+4) + 16$ (Rearranging)
 $= [x(x+6)][(x+2)(x+4)] + 16$
 $= (x^2 + 6x)(x^2 + 6x - 8) + 16 \dots\dots\dots(i)$

Let $x^2 + 6x = y$
 $= (y)(y + 8) + 16$ from (i)
 $= y^2 + 8y + 16$
 $= y^2 + 4y + 4y + 16$
 $= (y^2 + 4y) + (4y + 16)$
 $= y(y + 4) + 4(y + 4)$
 $= (y + 4)(y + 4)$
 $= (y + 4)^2$

Putting values of y in $x^2 + 6x$

$= (x^2 + 6x + 4)^2$
 $= \sqrt{(x^2 + 6x + 4)^2}$ (Taking square root)
 $= \pm(x^2 + 6x + 4)$

Q.5 $(2x+1)(2x+3)(2x+5)(2x+7) + 16$

Sol: (Rearranging)
 $= (2x+1)(2x+7)(2x+3)(2x+5) + 16$
 $= [(2x+1)(2x+7)][(2x+3)(2x+5)] + 16$
 $= [4x^2 + 16x + 7][(4x^2 + 16x + 15)] + 16$

Let $4x^2 + 16x = y$
 $= (y + 7)(y + 15) + 16$
 $= y^2 + 22y + 105 + 16$
 $= y^2 + 22y + 121$
 $= y^2 + 11y + 11y + 121$
 $= (y^2 + 11y) + (11y + 121)$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

109

$$= x(y + 11) + 11(y + 11)$$

$$= (y + 11)(y + 11)$$

$$= (y + 11)^2$$

Putting value of y

$$= (4x^2 + 16x + 11)^2$$

$$= \sqrt{(4x^2 + 16x + 11)^2} \quad (\text{Taking square root})$$

$$= \pm(4x^2 + 16x + 11)$$

Q.6 $\left(x^2 + \frac{1}{x^2}\right) - 10\left(x + \frac{1}{x}\right) + 27; x \neq 0$

Sol: $\left(x^2 + \frac{1}{x^2}\right) - 10\left(x + \frac{1}{x}\right) + 27 \quad (i)$

Let $x + \frac{1}{x} = y$

$$x^2 + \frac{1}{x^2} + 2 = y^2 \quad (\text{Squaring both sides.})$$

$$x^2 + \frac{1}{x^2} = y^2 - 2$$

Putting values $x + \frac{1}{x} = y$ and $x^2 + \frac{1}{x^2} = y^2 - 2$ in (i).

$$= y^2 - 2 - 10y + 27$$

$$= y^2 - 10y + 25$$

$$= y^2 - 5y - 5y + 25$$

$$= y(y - 5) - 5(y - 5)$$

$$= (y - 5)(y - 5)$$

$$= (y - 5)^2$$

$$= \left(x + \frac{1}{x} - 5\right)^2$$

$$= \sqrt{\left(x + \frac{1}{x} - 5\right)^2} \quad (\text{Taking square root})$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

110

$$= \pm \left(x + \frac{1}{x} - 5 \right)$$

Q.7 $\left(t - \frac{1}{t} \right)^2 - 4 \left(t + \frac{1}{t} \right) + 8, (t \neq 0)$

Sol: Let $t + \frac{1}{t} = y$

We know that

$$\begin{aligned} \left(t - \frac{1}{t} \right)^2 &= \left(t + \frac{1}{t} \right)^2 - 4t \times \frac{1}{t} \\ &= y^2 - 4 \end{aligned}$$

Now $t + \frac{1}{t} = y$ and $\left(t + \frac{1}{t} \right)^2 = y^2 - 4$ putting in given values

$$= y^2 - 4 - 4y + 8$$

$$= y^2 - 4y + 4$$

$$= y^2 - 2y - 2y + 4$$

$$= y(y - 2) - 2(y - 2)$$

$$= (y - 2)(y - 2)$$

$$= (y - 2)^2$$

Putting value of y.

$$= \left(t + \frac{1}{t} - 2 \right)^2$$

$$= \sqrt{\left(t + \frac{1}{t} - 2 \right)^2} \quad \text{(Taking square root)}$$

$$= \pm \left(t + \frac{1}{t} - 2 \right)$$

Q.8 $\left(x^2 + \frac{1}{x^2} \right)^2 - 4 \left(x + \frac{1}{x} \right)^2 + 12; x \neq 0$

Sol: Let $x + \frac{1}{x} = y$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

111

$$x + \frac{1}{x} = y$$

Squaring both sides

$$\left(x + \frac{1}{x}\right)^2 = y^2$$

$$x^2 + \frac{1}{x^2} + 2 = y^2 \dots\dots\dots (i)$$

$$x^2 + \frac{1}{x^2} = y^2 - 2$$

Putting $x + \frac{1}{x} = y$ and $x^2 + \frac{1}{x^2} = y^2 - 2$ in the given statement.

$$\begin{aligned} &= (y^2 - 2)^2 - 4y^2 + 12 \\ &= y^4 - 4y^2 + 4 - 4y^2 + 12 \\ &= y^4 - 8y^2 + 16 \\ &= y^4 - 4y^2 - 4y^2 + 16 \\ &= y^2(y^2 - 4) - 4(y^2 - 4) \\ &= (y^2 - 4)(y^2 - 4) \\ &= (y^2 - 4)^2 \end{aligned}$$

From (i), putting values $y^2 = x^2 + \frac{1}{x^2} + 2$

$$\begin{aligned} &= \left(x^2 + \frac{1}{x^2} + 2 - 4\right)^2 \\ &= \left(x^2 + \frac{1}{x^2} - 2\right)^2 \\ &= \sqrt{\left(x^2 + \frac{1}{x^2} - 2\right)^2} \quad \text{(Taking square root)} \\ &= \pm \left(x^2 + \frac{1}{x^2} - 2\right) \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

112

Q.9 $4x^4 + 12x^3 + 25x^2 + 24x + 16$

Sol: $2x^2 + 3x + 4$

$2x^2$	$4x^4 + 12x^3 + 25x^2 + 24x + 16$ $\pm 4x^4$
$4x^2 + 3x$	$12x^3 + 25x^2 + 24x + 16$ $\pm 12x^3 \pm 9x^2$
$4x^2 + 6x + 4$	$+ 16x^2 + 24x + 16$ $\pm 16x^2 \pm 24x \pm 16$
	0

Square root = $\pm(2x^2 + 3x + 4)$

Q.10 $\frac{9x^3}{4y^2} - \frac{3x}{2y} - \frac{7}{4} + \frac{2y}{3x} + \frac{4y^2}{9x^2} (x \neq 0, y \neq 0)$

Sol:

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

113

$$\text{Square root} = \pm \left[\frac{3x}{2y} - \frac{1}{2} - \frac{2y}{3x} \right]$$

Q.11. For what value of x , $x^4 + 4x^2 + x + \frac{8}{x^2} + \frac{4}{x^4}$ is a complete square, where $x \neq 0$

Sol: First we will find square root

$$\begin{array}{r} x^2 \overline{) x^4 + 4x^2 + x + \frac{8}{x^2} + \frac{4}{x^4}} \\ \underline{\pm x^4} \phantom{+ x + \frac{8}{x^2} + \frac{4}{x^4}} \\ 4x^2 + x + \frac{8}{x^2} + \frac{4}{x^4} \\ 2x^2 + 2 \overline{) 4x^2 + x + \frac{8}{x^2} + \frac{4}{x^4}} \\ \underline{\pm 4x^2 \pm 4} \phantom{+ \frac{8}{x^2} + \frac{4}{x^4}} \\ -4 + x + \frac{8}{x^2} + \frac{4}{x^4} \\ 2x^2 + 4 + \frac{2}{x^2} \overline{) -4 + x + \frac{8}{x^2} + \frac{4}{x^4}} \\ \underline{\pm 4 \phantom{+ x + \frac{8}{x^2} + \frac{4}{x^4}}} \phantom{+ \frac{8}{x^2} + \frac{4}{x^4}} \\ -8 + x \phantom{+ \frac{4}{x^4}} \end{array}$$

For completing square remainder must be zero.

Therefore $-8 + x = 0$

$\Rightarrow x = 8$

Q.12. If $x^4 + lx^3 + mx^2 + 12x + 9$ is a complete square then find the values of l and m .

Sol:

$$\begin{array}{r} x^2 \overline{) x^4 + lx^3 + mx^2 + 12x + 9} \\ \underline{\pm x^4} \\ lx^3 + mx^2 + 12x + 9 \\ 2x^2 + 2x \overline{) lx^3 + mx^2 + 12x + 9} \\ \underline{\pm 4x^3 \pm 4x^2} \\ (l-4)x^3 + mx^2 - 4x^2 + 12x + 9 \\ 2x^2 + 4x + 3 \overline{) (l-4)x^3 + mx^2 - 4x^2 + 12x + 9} \\ \underline{\pm 6x^2 \pm 12x \pm 9} \\ (l-4)x^3 + mx^2 - 10x^2 \end{array}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 3)

Pilot Super One "General Math" 10th

114

For completing square remainder must be zero

$$(l - 4)x^3 + (m - 10)x^2 = 0$$

Therefore, $l - 4 = 0$ and $m - 10 = 0$

$$l = 4, m = 10$$

0

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 4)

Pilot Super One "General Math" 10th

118

EXERCISE 14

Solve:

Q.1. (i) $3x + 20 = 44$

Sol. $3x = 44 - 20$

$$3x = 24$$

$$\frac{3x}{3} = \frac{24}{3}$$

(Dividing by 3)

$$x = 8$$

(ii) $\frac{4x}{5} - \frac{3x}{4} = 4$

Sol. Multiplied by 4, 5 L.C.M. 20.

$$\frac{4x}{5}(20) - \frac{3x}{4}(20) = 4(20)$$

$$16x - 15x = 80$$

$$x = 80$$

(iii) $3x + 3(x + 1) = 69$

Sol. $3x + 3x + 3 = 69$

$$6x + 3 = 69$$

$$6x = 69 - 3$$

$$6x = 66$$

$$\frac{6x}{6} = \frac{66}{6} \quad (\text{Dividing by 6})$$

$$x = 11$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 4)

Pilot Super One "General Math" 10th

119

(iv) $(90 - 9x) + 27 = 90 + 9$

Sol. $90 - 9x + 27 = 99$

$$-9x = 99 - 90 - 27$$

$$-9x = -18$$

$$\frac{-9x}{-9} = \frac{-18}{-9} \text{ (Dividing by -9)}$$

$$x = 2$$

Q.2. $3(x + 3) = 14 + x$

Sol. $3x + 9 = 14 + x$

$$3x - x = 14 - 9$$

$$2x = 5$$

$$\frac{2x}{2} = \frac{5}{2} \text{ (Dividing by 2)}$$

$$x = \frac{5}{2}$$

Q.3. $3(2x + 5) = 25 + x$

Sol. $6x + 15 = 25 + x$

$$6x - x = 25 - 15$$

$$5x = 10$$

$$\frac{5x}{5} = \frac{10}{5} \text{ (Dividing by 5)}$$

$$x = 2$$

Q.4. $9x - 3 = 3(2x - 8)$

Sol. $9x - 3 = 6x - 24$

$$9x - 6x = -24 + 3$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 4)

=====

Pilot Super One "General Math" 10th

120

$$3x = -21$$

$$\frac{3x}{3} = \frac{-21}{3} \text{ (Dividing by 3)}$$

$$x = -7$$

Q.5. $3(2x - 1) = 5(x - 1)$

Sol. $6x - 3 = 5x - 5$

$$6x - 5x = -5 + 3 \text{ (By Transposing)}$$

$$x = -2$$

Q.6. $2(7x - 6) = 3(1 + 3x)$

Sol. $14x - 12 = 3 + 9x$

$$14x - 9x = 3 + 12$$

$$5x = 15$$

$$\frac{x}{5} = \frac{15}{5} \text{ (Dividing by 5)}$$

$$x = 3$$

Q.7. $\frac{10x - 1}{2x + 5} = 3$

Sol. $10x - 1 = 3(2x + 5)$

$$10x - 1 = 6x + 15$$

$$10x - 6x = 15 + 1$$

$$4x = 16$$

$$\frac{x}{4} = \frac{16}{4} \text{ (Dividing by 4)}$$

$$x = 4$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 4)

Pilot Super One "General Math" 10th

121

Q.8. $\frac{2x+1}{x+5} = 1$

Sol. $2x + 1 = 1(x + 5)$
 $2x + 1 = x + 5$
 $2x - x = 5 - 1$
 $x = 4$

Q.9. $\frac{5x+3}{x+6} = 2$

Sol. $5x + 3 = 2(x + 6)$
 $5x + 3 = 2x + 12$
 $5x - 2x = 12 - 3$
 $3x = 9$
 $\frac{x}{3} = \frac{9}{3}$ (Dividing by 3)
 $x = 3$

Q.10. $y - 6 + \sqrt{y} = 0$

Sol. $\sqrt{y} = 6 - y$
 $(\sqrt{y})^2 = (6 - y)^2$ (Squaring on both sides)
 $y = 36 + y^2 - 12y$
 $0 = 36 + y^2 - 12y - y$
 $0 = 36 + y^2 - 13y$
 $\Rightarrow y^2 - 13y + 36 = 0$
 $y^2 - 4y - 9y + 36 = 0$
 $y(y - 4) - 9(y - 4) = 0$
 $(y - 4)(y - 9) = 0$
 If $y - 4 = 0$
 Then, $y = 4$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 4)

Pilot Super One "General Math" 10th

122

and if $y - 9 = 0$

Then, $y = 9$

Check to eliminate extraneous root $y = 9$.

Q.11

$$x = 15 - 2\sqrt{x}$$

Sol.

$$2\sqrt{x} = 15 - x$$

$$(2\sqrt{x})^2 = (15 - x)^2 \text{ (Squaring on both sides)}$$

$$4x = 225 + x^2 - 30x$$

$$0 = 225 + x^2 - 30x - 4x$$

$$0 = x^2 + 225 - 34x$$

$$\Rightarrow x^2 - 34x + 225 = 0$$

$$x^2 - 9x - 25x + 225 = 0$$

$$x(x - 9) - 25(x - 9) = 0$$

$$(x - 9)(x - 25) = 0$$

$$\text{If } x - 9 = 0$$

$$\text{then } x = 9$$

$$\text{And if } x - 25 = 0$$

$$\text{then } x = 25$$

Check to eliminate extraneous roots $x = 25$

Q.12.

$$m - 13 = \sqrt{m + 7}$$

Sol.

$$(m - 13)^2 = (\sqrt{m + 7})^2 \text{ Squaring both sides}$$

$$m^2 - 26m + 169 = m + 7$$

$$m^2 - 26m + 169 - m - 7 = 0$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 4)

Pink Super One "General Math" 10th

123

$$m^2 - 27m + 162 = 0$$

$$m^2 - 18m - 9m + 162 = 0$$

$$m(m - 18) - 9(m - 18) = 0$$

$$(m - 18)(m - 9) = 0$$

$$\text{If } m - 18 = 0$$

$$\text{then } m = 18$$

$$\text{If any } m - 9 = 0$$

$$\text{Then } m = 9$$

Check to eliminate extraneous roots $m = 9$

Q.13 $\sqrt{5n+9} = n-1$

Sol. $(\sqrt{5n+9})^2 = (n-1)^2$ (Squaring both sides)

$$5n+9 = n^2 - 2n + 1$$

$$0 = n^2 - 2n + 1 - 5n - 9$$

$$0 = n^2 - 7n - 8$$

$$\Rightarrow n^2 - 7n - 8 = 0$$

$$n^2 - 8n + n - 8 = 0$$

$$n(n-8) + 1(n-8) = 0$$

$$(n-8)(n+1) = 0$$

$$\text{If } n - 8 = 0$$

$$\text{Then } n = 8$$

$$\text{If any } n + 1 = 0$$

$$\text{then } n = -1$$

Check to eliminate extraneous roots $n = -1$

Q.14. $3 + \sqrt{2x-1} = 0$

Sol. $\sqrt{2x-1} = 0 - 3$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 4)

Pilot Super One "General Math" 10th

124

$$\sqrt{2x-1} = -3$$

Square root is always positive.

Therefore = { } S.S.

Let solve

$$\sqrt{2x-1} = -3$$

$$(\sqrt{2x-1})^2 = (-3)^2 \text{ (Squaring both sides)}$$

$$2x - 1 = 9$$

$$2x = 9 + 1$$

$$2x = 10$$

$$\frac{x}{2} = \frac{10}{2} \text{ (Dividing by 2)}$$

$$x = 5$$

L. H. S. = $\sqrt{2x-1}$ To check

$$= \sqrt{2 \times 5 - 1} \text{ (Putting } x = 5)$$

$$= \sqrt{10-1}$$

$$= \sqrt{9}$$

$$= 3$$

Square root is always positive.

Hence, S.S. = { }

$$\sqrt{x+5} + 7 = 0$$

$$\sqrt{x+5} = 0 - 7$$

$$\sqrt{x+5} = -7$$

Square root is always taken positive.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 4)

Pilot Super One "General Math" 10th

125

Hence, S.S. = { }

Q.16. $\sqrt{2x-1} - \sqrt{x-4} = 2$

Sol. $\sqrt{2x-1} = 2 + \sqrt{x-4}$

Squaring root on both sides

$$(\sqrt{2x-1})^2 = (2 + \sqrt{x-4})^2$$

$$2x-1 = 4 + x - 4 + 4\sqrt{x-4}$$

$$2x-1-x = 4\sqrt{x-4}$$

$$x-1 = 4\sqrt{x-4}$$

(Again squaring) $(x-1)^2 = (4\sqrt{x-4})^2$

$$x^2 - 2x + 1 = 16(x-4)$$

$$x^2 - 2x + 1 = 16x - 64$$

$$x^2 - 2x - 16x + 1 + 64 = 0$$

$$x^2 - 18x + 65 = 0$$

$$x^2 - 13x - 5x + 65 = 0$$

$$x(x-13) - 5(x-13) = 0$$

$$(x-13)(x-5) = 0$$

If $x-13 = 0$

Then $x = 13$

If any $x-5 = 0$

then $x = 5$

$$\text{S.S.} = \{13, 5\}$$

Q.17. $\sqrt{x+1} = 3$

Sol. $(\sqrt{x+1})^2 = (3)^2$ (Squaring both sides)

$$x+1 = 9$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 4)

Pilot Super One "General Math" 10th

126

$$x = 9 - 1$$

$$x = 8$$

Q.18. $\sqrt{2x-1} = 5$

Sol. $(\sqrt{2x-1})^2 = (5)^2$ (Squaring both sides)

$$2x - 1 = 25$$

$$2x = 25 + 1$$

$$2x = 26$$

$$x = \frac{26}{2}$$

$$x = 13$$

Q.19. $\sqrt{x-1} = 10$

Sol. $(\sqrt{x-1})^2 = (10)^2$ (Squaring both sides)

$$x - 1 = 100$$

$$x = 100 + 1$$

$$x = 101$$

Q.20. $\sqrt{3x+4} = 7$

Sol. $(\sqrt{3x+4})^2 = (7)^2$ (Squaring both sides)

$$3x + 4 = 49$$

$$3x = 49 - 4$$

$$3x = 45$$

$$x = \frac{45}{3}$$

$$x = 15$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 4)

Pilot Super One "General Math" 10th

127

Exercise 1.2

Solve and Check:

Q.1. $|x| = 9$

Sol. $x = 9$ And $-x = 9$
 $x = 9$

Hence, $x = \pm 9$

Check, $|\pm 9| = 9$

Check

Q.2. $|x - 3| = 4$

<p>Sol. $x - 3 = 4$ $x = 4 + 3$ $x = 7$</p>	<p>$-(x - 3) = 4$ $-x + 3 = 4$ $-x = 4 - 3$ $-x = 1$ $x = -1$</p>
---	--

Hence: $x = -1, 7$

Check: $|-1 - 3| = |-4| = 4$ and $|7 - 3| = |4| = 4$

Q.3. $|x + 1| = 5$

<p>Sol. $x + 1 = 5$ $x = 5 - 1$ $x = 4$</p>	<p>$-(x + 1) = 5$ $-x - 1 = 5$ $-x = 5 + 1$ $-x = 6$ $x = -6$</p>
---	--

Hence, $x = 4, -6$

Check: $|4 + 1| = |5| = 5$ and $|-6 + 1| = |-5| = 5$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 4)

Pilot Super One "General Math" 10th

128

Q.4. $|2x - 3| = 5$

<p>Sol. $2x - 3 = 5$ $2x = 5 + 3$ $2x = 8$ $\frac{x}{2} = \frac{8}{2}$ $x = 4$</p>	<p>$-(2x - 3) = 5$ $-2x + 3 = 5$ $-2x = 5 - 3$ $\frac{-2x}{2} = \frac{2}{-2}$ $x = -1$</p>
---	---

S.S. = $\{4, -1\}$

Check: $|2(4) - 3| = |8 - 3| = |5| = 5$ and $|2(-1) - 3|$
 $= |-2 - 3| = |-5| = 5$

Q.5. $|3x + 4| = 9$

<p>Sol. $3x + 4 = 9$ $3x = 9 - 4$ $3x = 5$ $x = \frac{5}{3}$</p>	<p>$-(3x + 4) = 9$ $-3x - 4 = 9$ $-3x = 9 + 4$ $-3x = 13$ $x = -\frac{13}{3}$</p>
---	--

S. S. = $\left\{\frac{5}{3}, -\frac{13}{3}\right\}$

Check: Putting $\frac{5}{3}$

$\left|3\left(\frac{5}{3}\right) + 4\right| = |5 + 4| = |9| = 9$

Putting $-\frac{13}{3}$

$\left|3\left(-\frac{13}{3}\right) + 4\right| = |-13 + 4| = |-9| = 9$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 4)

Pilot Super One "General Math" 10th

129

Q.6. $3(x - 2) < 2x + 1$

Sol. $3x - 6 < 2x + 1$

$$3x - 2x < 1 + 6$$

$$x < 7$$

Check: Let $x = 6$

These putting values

$$3(6 - 2) < 2(6) + 1$$

$$3(4) < 12 + 1$$

$$12 < 13$$

Which is true

Therefore, $x < 7$

Q.7. $3(x + 5) > 2(x + 2) + 8$

Sol. $3x + 15 > 2x + 4 + 8$

$$3x - 2x > 4 + 8 - 15$$

$$x > -3$$

Check: Suppose that $x = -2$

$$3(-2 + 5) > 2(-2 + 2) + 8$$

$$3(3) > 2(0) + 8$$

$$9 > 8$$

Which is true

Hence, $x > -3$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 4)

Pilot Super One "General Math" 10th

130

Q.8. $\frac{1}{2}(2-x) > \frac{1}{4}(3-x) + \frac{1}{2}$

Sol. $\frac{2-x}{2} > \frac{3-x}{4} + \frac{1}{2}$

$$1 - \frac{x}{2} > \frac{3-x}{4} + \frac{1}{2}$$

$$-\frac{x}{2} + \frac{x}{4} > \frac{3}{4} + \frac{1}{2} - 1$$

$$\frac{-2x+x}{4} > \frac{3+2-4}{4}$$

$$-\frac{x}{4} > \frac{1}{4}$$

$$-x > 1 \quad (\text{Multiplying by 4})$$

Therefore, $x < -1$

Now, Suppose that $x = -2$

$$x$$

Putting value $x = -2$

$$\frac{1}{2}(2+2) > \frac{1}{4}(3+2) + \frac{1}{2}$$

$$\frac{4}{2} > \frac{5}{4} + \frac{1}{2}$$

$$2 > \frac{5+2}{4}$$

$$2 > \frac{7}{4} \quad (\text{Which is true})$$

Hence, $x < -1$

Q.9. $\frac{x-2}{4} + \frac{2}{3} < \frac{x-4}{6}$

Sol. Multiply by L.C.M. of 4, 3, 6 by 12.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 4)

Pilot Super One "General Math" 10th

131

$$12 \frac{(x-2)}{4} + 12 \left(\frac{2}{3} \right) < 12 \frac{(x-4)}{6}$$

$$3(x-2) + 4(2) < 2(x-4)$$

$$3x - 6 + 8 < 2x - 8$$

$$3x - 2x < -8 - 8 + 6$$

$$x < -10$$

Q.10. $\frac{3x+4}{5} - \frac{x+1}{3} > 1 - \frac{x+5}{3}$

Sol. Multiply by L.C.M of 5, 3 by 15.

$$15 \left(\frac{3x+4}{5} \right) - 15 \left(\frac{x+1}{3} \right) > 15 \times 1 - 15 \left(\frac{x+5}{3} \right)$$

$$3(3x+4) - 5(x+1) > 15 - 5(x+5)$$

$$9x + 12 - 5x - 5 > 15 - 5x - 25$$

$$9x - 5x + 5x > 15 - 25 - 12 + 5$$

$$9x > -17$$

$$x > -\frac{17}{9}$$

Q.11. $\frac{x+1}{2} - \frac{x+3}{3} > \frac{x+1}{4} + 1$

Sol. Multiply by 12, L.C.M of 4, 3, 2.

$$12 \left(\frac{x+1}{2} \right) - 12 \left(\frac{x+3}{3} \right) > 12 \left(\frac{x+1}{4} \right) + 12 \times 1$$

$$6(x+1) - 4(x+3) > 3(x+1) + 12$$

$$6x + 6 - 4x - 12 > 3x + 3 + 12$$

$$6x - 4x - 3x > 3 + 12 - 6 + 12$$

$$-x > 21$$

Therefore, $x < -21$ N.T.S

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 4)

Pilot Super One "General Math" 10th

132

Q.12. $\frac{x+3}{4} - \frac{x+2}{5} < 1 + \frac{x+5}{6}$

Sol. Multiply by 60, L.C.M. of 4, 5, 6.

$$60\left(\frac{x+3}{4}\right) - 60\left(\frac{x+2}{5}\right) < 60 \times 1 + 60\left(\frac{x+5}{6}\right)$$

$$15(x+3) - 12(x+2) < 60 + 10(x+5)$$

$$15x + 45 - 12x - 24 < 60 + 10x + 50$$

$$15x - 12x - 10x < 60 + 50 - 45 + 24$$

$$-7x < 89$$

$$7x < -89 \text{ (Changing symbol)}$$

$$x > -\frac{89}{7}$$

$$x > -12\frac{5}{7}$$

Q.13. $\frac{1}{2}x \geq 1 + \frac{1}{3}x$

Sol. Multiply by 6, L.C.M. of 2, 3.

$$6\left(\frac{1}{2}x\right) \geq 6 \times 1 + 6 \times \frac{1}{3}x$$

$$3x \geq 6 + 2x$$

$$3x - 2x \geq 6$$

$$x \geq 6$$

Q.14. $\frac{1}{4}(2x+3) \leq (7-4x)$

Sol. $4 \times \frac{1}{4}(2x+3) \leq 4(7-4x) \text{ (Multiply by 4)}$

$$2x+3 \leq 28-16x$$

$$+16x \leq 28-3$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 4)

Pilot Super One "General Math" 10th

133

$$18x \leq 25$$

$$\frac{18x}{18} \leq \frac{25}{18} \quad (\text{Dividing by 18})$$

$$x \leq \frac{7}{18}$$

Q.15.

$$\frac{4}{3}(2x+3) \geq 10 - \frac{4x}{3}$$

Sol

$$3 \times \frac{4}{3}(2x+3) \geq 3 \times 10 - 3 \times \frac{4x}{3} \quad (\text{Multiply by 3})$$

$$4(2x+3) \geq 30 - 4x$$

$$8x + 12 \geq 30 - 4x$$

$$8x + 4x \geq 30 - 12$$

$$12x \geq 18$$

$$x \geq \frac{18}{12}$$

$$x \geq \frac{3}{2}$$

$$x \geq 1\frac{1}{2}$$

Q.16.

$$\frac{x-2}{4} - \frac{x-5}{6} \geq \frac{1}{3}$$

Sol.

Multiply by 12, L.C.M. of 4, 6, 3.

$$12\left(\frac{x-2}{4}\right) - 12\left(\frac{x-5}{6}\right) \geq 12 \times \frac{1}{3}$$

$$3(x-2) - 2(x-5) \geq 4$$

$$3x - 6 - 2x + 10 \geq 4$$

$$3x - 2x \geq 4 + 6 - 10$$

$$x \geq 0$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

138



-I Solve by using factrization method.

Q.1. $x^2 - 4x - 12 = 0$

Sol. $x^2 - 6x + 2x - 12 = 0$

$$(x^2 - 6x) + (2x - 12) = 0$$

$$x(x - 6) + 2(x - 6) = 0$$

$$(x - 6)(x + 2) = 0$$

If $x - 6 = 0$

then $x = 6$

and if $x + 2 = 0$

then $x = -2$

$$\text{S. S.} = \{6, -2\}$$

Q.2. $x^2 - 6x + 5 = 0$

Sol. $x^2 - x - 5x + 5 = 0$

$$(x^2 - x) - (5x - 5) = 0$$

$$x(x - 1) - 5(x - 1) = 0$$

$$(x - 1)(x - 5) = 0$$

If $x - 1 = 0$

then $x = 1$

and if $x - 5 = 0$

then $x = 5$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

139

$$S.S. = \{1, 5\}$$

Q.3.

$$x^2 = 8 - 7x$$

Sol.

$$x^2 + 7x - 8 = 0$$

$$x^2 + 8x - x - 8 = 0$$

$$(x^2 + 8x) - (x + 8) = 0$$

$$x(x + 8) - 1(x + 8) = 0$$

$$(x + 8)(x - 1) = 0$$

If $x + 8 = 0$

then $x = -8$

and if $x - 1 = 0$

then $x = 1$

$$\text{Solution set} = \{-8, 1\}$$

Q.4.

$$5x = x^2 + 6$$

Sol.

$$0 = x^2 + 6 - 5x$$

$$x^2 + 6 - 5x = 0$$

$$x^2 - 5x + 6 = 0$$

$$x^2 - 2x - 3x + 6 = 0$$

$$(x^2 - 2x) - (3x - 6) = 0$$

$$x(x - 2) - 3(x - 2) = 0$$

$$(x - 2)(x - 3) = 0$$

If $x - 2 = 0$

then $x = 2$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

140

and if $x - 3 = 0$

then $x = 3$

Solution set = $\{2, 3\}$

Q.5. $3x^2 - 10x + 8 = 0$

Sol. $3x^2 - 6x - 4x + 8 = 0$

$(3x^2 - 6x) - (4x - 8) = 0$

$3x(x - 2) - 4(x - 2) = 0$

$(x - 2)(3x - 4) = 0$

If $x - 2 = 0$

then $x = 2$

and if $3x - 4 = 0$

$3x = 4$

then $x = \frac{4}{3}$

Solution set = $\left\{2, \frac{4}{3}\right\}$

Q.6. $2x^2 + 15x - 8 = 0$

Sol. $2x^2 - x + 16x - 8 = 0$

$(2x^2 - x) + (16x - 8) = 0$

$x(2x - 1) + 8(2x - 1) = 0$

$(2x - 1)(x + 8) = 0$

If $2x - 1 = 0$

$2x = 1$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

141

then

$$x = \frac{1}{2}$$

and if

$$x + 8 = 0$$

then

$$x = -8$$

$$\text{Solution set} = \left\{ \frac{1}{2}, -8 \right\}$$

Q.7.

$$\frac{x}{4}(x+1) = 3$$

Sol.

$$4\left(\frac{x}{4}\right)(x+1) = 3 \times 4 \text{ (Multiply by 4)}$$

$$x(x+1) = 12$$

$$x^2 + x = 12$$

$$x^2 + x - 12 = 0$$

$$x^2 + 4x - 3x - 12 = 0$$

$$(x^2 + 4x) - (3x + 12) = 0$$

$$x(x+4) - 3(x+4) = 0$$

$$(x+4)(x-3) = 0$$

$$\text{If } x+4 = 0$$

$$\text{then } x = -4$$

$$\text{and if } x-3 = 0$$

$$\text{then } x = 3$$

$$\text{Solution set} = \{-4, 3\}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

142

Q.8. $3x^2 - 8x - 3 = 0$

Sol. $3x^2 - 9x + x - 3 = 0$

$$(3x^2 - 9x) + (x - 3) = 0$$

$$3x(x - 3) + 1(x - 3) = 0$$

$$(x - 3)(3x + 1) = 0$$

If $x - 3 = 0$

then $x = 3$

and if $3x + 1 = 0$

$$3x = -1$$

then $x = -\frac{1}{3}$

$$\text{Solution set} = \left\{ 3, -\frac{1}{3} \right\}$$

Q.9. $2x = \frac{2}{x} + 3$

Sol. (Multiply by x)



$$x(2x) = x\left(\frac{2}{x}\right) + 3(x)$$

$$2x^2 = 2 + 3x$$

$$2x^2 - 3x - 2 = 0$$

$$2x^2 - 4x + x - 2 = 0$$

$$(2x^2 - 4x) + (x - 2) = 0$$

$$2x(x - 2) + 1(x - 2) = 0$$

$$(x - 2)(2x + 1) = 0$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

143

If $x - 2 = 0$

then $x = 2$

and if $2x + 1 = 0$

$$2x = -1$$

then $x = -\frac{1}{2}$

$$\text{Solution set} = \left\{ 2, -\frac{1}{2} \right\}$$

Q.10. $5x^2 - 6x - 8 = 0$

Sol. $5x^2 - 10x + 4x - 8 = 0$

$$(5x^2 - 10x) + (4x - 8) = 0$$

$$5x(x - 2) + 4(x - 2) = 0$$

$$(x - 2)(5x + 4) = 0$$

If $x - 2 = 0$

then $x = 2$

and if $5x + 4 = 0$

$$5x = -4$$

then $x = -\frac{4}{5}$

$$\text{Solution set} = \left\{ 2, -\frac{4}{5} \right\}$$

Q.11. $(2x + 3)(x - 2) = 0$

Sol. If $x - 2 = 0$

then $x = 2$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

=====

Pilot Super One "General Math" 10th

144

and if $2x + 3 = 0$

$$2x = -3$$

then $\boxed{x = -\frac{3}{2}}$

$$\text{Solution set} = \left\{ 2, -\frac{3}{2} \right\}$$

Q.12. $(2x + 1)(5x - 4) = 0$

Sol. If $2x + 1 = 0$

$$2x = -1$$

then $\boxed{x = -\frac{1}{2}}$

and if $5x - 4 = 0$

$$5x = 4$$

then $\boxed{x = \frac{4}{5}}$

$$\text{Solution set} = \left\{ -\frac{1}{2}, \frac{4}{5} \right\}$$

Q.13. $4x(3x - 1) - 2 = (2x - 1)(5x + 1)$

Sol. $12x^2 - 4x - 2 = 10x^2 - 3x - 1$

$$12x^2 - 10x^2 - 4x + 3x - 2 + 1 = 0$$

$$2x^2 - x - 1 = 0$$

$$2x^2 - 2x + x - 1 = 0$$

$$(2x^2 - 2x) + (x - 1) = 0$$

$$2x(x - 1) + 1(x - 1) = 0$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

145

$$(x - 1)(2x + 1) = 0$$

If $x - 1 = 0$

then $x = 1$

and if $2x + 1 = 0$

$$2x = -1$$

then $x = -\frac{1}{2}$

$$\text{Solution set} = \left\{1, -\frac{1}{2}\right\}$$

II- Solve by completing the square method.

Q.14 $x^2 - 10x - 3 = 0$

Sol. $x^2 - 10x = 3$

$\frac{10}{2} = 5$ then add $(5)^2$ on both sides

$$x^2 - 10x + (5)^2 = 3 + (5)^2$$

$$(x - 5)^2 = 3 + 25$$

$$(x - 5)^2 = 28$$

(By taking square root on both sides)

$$x - 5 = \sqrt{28}$$

$$x - 5 = \pm 2\sqrt{7}$$

$$x = 5 \pm 2\sqrt{7}$$

Q.15 $x^2 - 6x - 3 = 0$

Sol. $x^2 - 6x = 3$

$\frac{6}{2} = 3$ then add $(3)^2$ on both sides

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

148

$$x^2 - 6x + (3)^2 = 3 + (3)^2$$

$$(x-3)^2 = 3 + 9$$

$$(x-3)^2 = 12$$

(By taking square root on both sides)

$$x-3 = \pm\sqrt{12}$$

$$x = \pm 2\sqrt{3}$$

$$x = 3 \pm 2\sqrt{3}$$

Q.16

$$x^2 + x - 1 = 0$$

Sol.

$$x^2 + x = 1$$

Adding $\left(\frac{1}{2}\right)^2$ on both sides

$$x^2 + x + \left(\frac{1}{2}\right)^2 = 1 + \left(\frac{1}{2}\right)^2$$

$$\left(x + \frac{1}{2}\right)^2 = 1 + \frac{1}{4}$$

$$\left(x + \frac{1}{2}\right)^2 = \frac{4+1}{4}$$

$$\left(x + \frac{1}{2}\right)^2 = \frac{5}{4}$$

$$x + \frac{1}{2} = \pm\sqrt{\frac{5}{4}} \text{ (Taking square root)}$$

$$x = -\frac{1}{2} \pm \sqrt{\frac{5}{4}}$$

$$= -\frac{1}{2} \pm \frac{\sqrt{5}}{2}$$

$$= \frac{-1 \pm \sqrt{5}}{2}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

147

Q.17. $x^2 + 6x - 3 = 0$

Sol. $x^2 + 6x = 3$

$\frac{6}{2} = 3$, then $(3)^2$ adding on both sides

$$x^2 + 6x + (3)^2 = 3 + (3)^2$$

$$(x+3)^2 = 3 + 9$$

$$(x+3)^2 = 12$$

$$x+3 = \pm\sqrt{12} \text{ (Taking square root)}$$

$$x+3 = \pm 2\sqrt{3}$$

$$x = -3 \pm 2\sqrt{3}$$

Q.18. $2x^2 - 4x + 1 = 0$

Sol. $2x^2 - 4x = -1$

Divided by co-efficient 2 of x .

$$x^2 - 2x = -\frac{1}{2}$$

$\frac{2}{2} = 1$, add $(1)^2$ on both sides

$$x^2 - 2x + (1)^2 = -\frac{1}{2} + (1)^2$$

$$(x-1)^2 = -\frac{1}{2} + 1$$

$$(x-1)^2 = \frac{-1+2}{2}$$

$$(x-1)^2 = \frac{1}{2}$$

$$x-1 = \pm\sqrt{\frac{1}{2}} \text{ (Taking square root)}$$

thus, $x = 1 \pm \sqrt{\frac{1}{2}}$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

=====

Pilot Super One "General Math" 10th

148

Q19. $2x^2 - 6x + 3 = 0$

Sol. $2x^2 - 6x = -3$

Divided by co-efficient 2 of x.

$$x^2 - 3x = -\frac{3}{2}$$

Adding $\left(\frac{3}{2}\right)^2$ on both sides

$$x^2 - 3x + \left(\frac{3}{2}\right)^2 = -\frac{3}{2} + \left(\frac{3}{2}\right)^2$$

$$\left(x - \frac{3}{2}\right)^2 = -\frac{3}{2} + \frac{9}{2}$$

$$\left(x - \frac{3}{2}\right)^2 = \frac{-3+9}{2}$$

$$\left(x - \frac{3}{2}\right)^2 = \frac{6}{2}$$

$$\left(x - \frac{3}{2}\right)^2 = 3$$

$$x - \frac{3}{2} = \pm\sqrt{3} \text{ (Taking square root)}$$

$$x = \frac{3}{2} \pm \sqrt{3}$$

$$x = \frac{3 \pm 2\sqrt{3}}{2}$$

Q20. $3x^2 + 5x - 4 = 0$

Sol. $3x^2 + 5x = 4$

Divided by 3 co-efficient of x.

$$x^2 + \frac{5}{3}x = \frac{4}{3}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

149

$$\frac{1}{2}\left(\frac{5}{3}\right) = \frac{5}{6}, \text{ Adding } \left(\frac{5}{6}\right)^2 \text{ on both sides}$$

$$x^2 + \frac{5}{3}x + \left(\frac{5}{6}\right)^2 = \left(\frac{5}{6}\right)^2 + \frac{4}{3}$$

$$\left(x + \frac{5}{6}\right)^2 = \frac{25}{36} + \frac{4}{3}$$

$$\left(x + \frac{5}{6}\right)^2 = \frac{25 + 48}{36}$$

$$\left(x + \frac{5}{6}\right)^2 = \frac{73}{36}$$

$$x + \frac{5}{6} = \pm \frac{\sqrt{73}}{6} \text{ (Taking square root)}$$

$$x = -\frac{5}{6} \pm \frac{\sqrt{73}}{6}$$

$$x = \frac{-5 \pm \sqrt{73}}{6}$$

Q21.

$$x^2 + mx + n = 0$$

Sol.

$$x^2 + mx = -n$$

Adding $\left(\frac{m}{2}\right)^2$ on both sides.

$$x^2 + mx + \left(\frac{m}{2}\right)^2 = \left(\frac{m}{2}\right)^2 - n$$

$$\left(x + \frac{m}{2}\right)^2 = \frac{m^2}{4} - n$$

$$\left(x + \frac{m}{2}\right)^2 = \frac{m^2 - 4n}{4}$$

$$x + \frac{m}{2} = \pm \sqrt{\frac{m^2 - 4n}{4}} \text{ (Taking square root)}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

150

$$x = \frac{-m}{2} \pm \sqrt{\frac{m^2 - 4n}{4}}$$

$$x = \frac{-m}{2} \pm \frac{\sqrt{m^2 - 4n}}{2}$$

$$x = \frac{-m \pm \sqrt{m^2 - 4n}}{2}$$

Q22

$$11x^2 = 6x + 21$$

Sol.

$$11x^2 - 6x = 21$$

$$x^2 - \frac{6}{11}x = \frac{21}{11} \text{ (Divided by 11)}$$

$$\left(\frac{6}{11} \times \frac{1}{2}\right)^2 = \frac{3}{11} \text{ adding } \left(\frac{3}{11}\right)^2 \text{ on both sides}$$

$$x^2 - \frac{6}{11}x + \left(\frac{3}{11}\right)^2 = \left(\frac{3}{11}\right)^2 + \frac{21}{11}$$

$$\left(x - \frac{3}{11}\right)^2 = \frac{9}{121} + \frac{21}{11}$$

$$\left(x - \frac{3}{11}\right)^2 = \frac{9 + 231}{121}$$

$$\left(x - \frac{3}{11}\right)^2 = \frac{240}{121}$$

$$x - \frac{3}{11} = \pm \sqrt{\frac{240}{121}} \text{ (Taking square root)}$$

$$x - \frac{3}{11} = \pm \sqrt{\frac{16 \times 15}{11 \times 11}}$$

$$x - \frac{3}{11} = \pm \frac{4\sqrt{15}}{11}$$

$$x = \frac{3}{11} \pm \frac{4\sqrt{15}}{11}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

151

$$x = \frac{3 \pm 4\sqrt{15}}{11}$$

Q23. $2x^2 + 8x - 26 = 0$

Sol. $2x^2 + 8x = 26$

$$x^2 + 4x = 13 \text{ (Dividing by 2)}$$

$$\frac{4}{2} = 2 \text{ adding } (2)^2 \text{ on both sides.}$$

$$x^2 + 4x + (2)^2 = (2)^2 + 13$$

$$(x+2)^2 = 4 + 13$$

$$(x+2)^2 = 17$$

$$x+2 = \pm\sqrt{17} \text{ (Taking square root)}$$

$$x = -2 \pm \sqrt{17}$$

Q24. $5x^2 - 20x - 28 = 0$

Sol. $5x^2 - 20x = 28$

$$x^2 - 4x = \frac{28}{5} \text{ (Divided by 5)}$$

$$\frac{4}{2} = 2, \text{ Adding } (2)^2 \text{ on both sides.}$$

$$x^2 - 4x + (2)^2 = (2)^2 + \frac{28}{5}$$

$$(x-2)^2 = 4 + \frac{28}{5}$$

$$(x-2)^2 = \frac{20+28}{5}$$

$$(x-2)^2 = \frac{48}{5}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

152

$$x - 2 = \pm \sqrt{\frac{48}{5}} \text{ (Taking square root)}$$

$$x - 2 = \pm \frac{4\sqrt{3}}{\sqrt{5}}$$

$$x = 2 \pm \frac{4\sqrt{3}}{\sqrt{5}} \text{ (Rationalize)}$$

$$x = 2 \pm 4 \frac{\sqrt{3}}{\sqrt{5}} \times \frac{\sqrt{5}}{\sqrt{5}}$$

$$x = 2 \pm \frac{4\sqrt{15}}{5}$$

$$x = \frac{10 \pm 4\sqrt{15}}{5} \text{ Ans.}$$

Q25.

$$x^2 - 11x - 26 = 0$$

Sol.

$$x^2 - 11x = 26$$

Adding $\left(\frac{11}{2}\right)^2$ on both sides.

$$x^2 - 11x + \left(\frac{11}{2}\right)^2 = \left(\frac{11}{2}\right)^2 + 26$$

$$\left(x - \frac{11}{2}\right)^2 = \frac{121}{4} + 26$$

$$\left(x - \frac{11}{2}\right)^2 = \frac{121 + 104}{4}$$

$$\left(x - \frac{11}{2}\right)^2 = \frac{225}{4}$$

$$x - \frac{11}{2} = \pm \sqrt{\frac{225}{4}} \text{ (Taking square root)}$$

$$x - \frac{11}{2} = \pm \frac{15}{2}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

153

$$x = \frac{11 \pm 15}{2}$$

$$x = \frac{11 \pm 15}{2}$$

$$x = \frac{11+15}{2}, \frac{11-15}{2}$$

$$x = \frac{26}{2}, \frac{-4}{2}$$

$$x = 13, -2$$

$$\text{S.S.} = \{13, -2\}$$

5.3 The Quadratic Formula

If $ax^2 + bx + c = 0$ $a \neq 0$

then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$



Solve using quadratic formula:

Q.1. $x^2 - 5x + 6 = 0$

Sol. Here $a = 1$

$$b = -5$$

$$c = 6$$

(Formula) $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

Putting values of a, b, c

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

154

$$x = \frac{-(-5) \pm \sqrt{(-5)^2 - 4(1)(6)}}{2(1)}$$

$$x = \frac{5 \pm \sqrt{25 - 24}}{2}$$

$$x = \frac{5 \pm \sqrt{1}}{2}$$

$$x = \frac{5 \pm 1}{2}$$

$$x = \frac{5+1}{2}, \frac{5-1}{2}$$

$$x = \frac{6}{2}, \frac{4}{2}$$

$$x = 3, 2$$

Solution set = {3, 2}

Q.2.

$$(3 - 4x) = (4x - 3)^2$$

Sol.

$$3 - 4x = 16x^2 - 24x + 9$$

$$0 = 16x^2 - 24x + 9 - 3 + 4x$$

$$0 = 16x^2 - 20x + 6$$

$$\text{or } 16x^2 - 20x + 6 = 0$$

$$8x^2 - 10x + 3 = 0 \quad (\text{Divided by 2})$$

$$\text{Here } a = 8$$

$$b = -10$$

$$c = 3$$

(Formula)

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Putting the values of a, b, c

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

155

$$x = \frac{-(-10) \pm \sqrt{(-10)^2 - 4(8)(3)}}{2(8)}$$

$$x = \frac{10 \pm \sqrt{100 - 96}}{16}$$

$$x = \frac{10 \pm \sqrt{4}}{16}$$

$$x = \frac{10 \pm 2}{16}$$

$$x = \frac{10+2}{16}, \frac{10-2}{16}$$

$$x = \frac{12}{16}, \frac{8}{16}$$

$$x = \frac{3}{4}, \frac{1}{2}$$

$$\text{Solution set} = \left\{ \frac{3}{4}, \frac{1}{2} \right\}$$

Q.3. $3x^2 + x - 2 = 0$

Sol. Here $a = 3$
 $b = 1$

$$c = -2$$

(Formula) $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

By putting the values of a, b, c

$$x = \frac{-1 \pm \sqrt{(1)^2 - 4(3)(-2)}}{2(3)}$$

$$x = \frac{-1 \pm \sqrt{1+24}}{6}$$

$$x = \frac{-1 \pm \sqrt{25}}{6}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

156

$$x = \frac{-1 \pm 5}{6}$$

$$x = \frac{-1+5}{6}, \frac{-1-5}{6}$$

$$x = \frac{4}{6}, \frac{-6}{6}$$

$$x = \frac{2}{3}, -1$$

$$\text{Solution set} = \left\{ \frac{2}{3}, -1 \right\}$$

0

Q.4. $10x^2 - 5x = 15$

Sol. $10x^2 - 5x - 15 = 0$

$$2x^2 - x - 3 = 0 \quad (\text{Divided by 5})$$

Here $a = 2$

$$b = -1$$

$$c = -3$$

Formula: $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

By putting the values of a, b, c

$$x = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(2)(-3)}}{2(2)}$$

$$x = \frac{1 \pm \sqrt{1+24}}{4}$$

$$x = \frac{1 \pm \sqrt{25}}{4}$$

$$x = \frac{1 \pm 5}{4}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

157

$$x = \frac{1+5}{4}, \frac{1-5}{4}$$

$$x = \frac{6}{4}, \frac{-4}{4}$$

$$x = \frac{3}{2}, -1$$

$$\text{Solution set} = \left\{ \frac{3}{2}, -1 \right\}$$

Q.5. $(x-1)(x+3) - 12 = 0$

Sol. $x^2 + 2x - 3 - 12 = 0$

$$x^2 + 2x - 15 = 0$$

Here $a = 1$

$$b = 2$$

$$c = -15$$

(Formula) $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

By putting the values of a, b, c

$$x = \frac{-2 \pm \sqrt{(2)^2 - 4(1)(-15)}}{2(1)}$$

$$x = \frac{-2 \pm \sqrt{4+60}}{2}$$

$$x = \frac{-2 \pm \sqrt{64}}{2}$$

$$x = \frac{-2 \pm 8}{2}$$

$$x = \frac{-2+8}{2}, \frac{-2-8}{2}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

158

$$x = \frac{6}{2}, \frac{-10}{2}$$

$$x = 3, -5$$

$$\text{Solution set} = \{3, -5\}$$

Q.6. $x(2x + 7) - 3(2x + 7) = 0$

Sol. $2x^2 + 7x - 6x - 21 = 0$

$$2x^2 + x - 21 = 0$$

Here $a = 2$

$$b = 1$$

$$c = -21$$

(Formula) $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

By putting the values of a, b, c

$$x = \frac{-1 \pm \sqrt{(1)^2 - 4(2)(-21)}}{2(2)}$$

$$x = \frac{-1 \pm \sqrt{1 + 168}}{4}$$

$$x = \frac{-1 \pm \sqrt{169}}{4}$$

$$x = \frac{-1 \pm 13}{4}$$

$$x = \frac{-1 + 13}{4}, \frac{-1 - 13}{4}$$

$$x = \frac{12}{4}, \frac{-14}{4}$$

$$x = 3, -\frac{7}{2}$$

$$\text{Solution set} = \left\{3, -\frac{7}{2}\right\}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

159

Q.7. $\frac{x+1}{x+4} = \frac{2x-1}{x+6}$, where $x \neq -4, -6$

Sol. $(x+1)(x+6) = (2x-1)(x+4)$

$$x^2 + 7x + 6 = 2x^2 + 7x - 4$$

$$x^2 - 2x^2 + 7x - 7x + 6 + 4 = 0$$

$$-x + 10 = 0$$

$$x - 10 = 0$$

Here $a = 1$

$$b = 0$$

$$c = -10$$

(Formula) $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

By putting the values of a, b, c

$$x = \frac{-(0) \pm \sqrt{(0)^2 - 4(1)(-10)}}{2(1)}$$

$$x = \frac{0 \pm \sqrt{0+40}}{2}$$

$$x = \frac{0 \pm \sqrt{40}}{2}$$

$$x = \frac{0 \pm 2\sqrt{10}}{2}$$

$$x = \pm \frac{2\sqrt{10}}{2}$$

$$x = \pm \sqrt{10}$$

Solution set = $\{\pm \sqrt{10}\}$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

160

Q.8. $\frac{x}{6} + \frac{6}{x} = \frac{4}{x} + \frac{x}{4}$, where $x \neq 0$

Sol. Multiplying by L.C.M. $12x$

$$12x \left(\frac{x}{6} \right) + 12x \left(\frac{6}{x} \right) = 12x \left(\frac{4}{x} \right) + 12x \left(\frac{x}{4} \right)$$

$$2x^2 + 72 = 48 + 3x^2$$

$$2x^2 - 3x^2 + 72 - 48 = 0$$

$$-x^2 + 24 = 0$$

$$x^2 - 24 = 0$$

Here $a = 1$

$$b = 0$$

$$c = -24$$

(Formula) $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

By putting the values of a, b, c

$$x = \frac{-(0) \pm \sqrt{(0)^2 - 4(1)(-24)}}{2(1)}$$

$$x = \frac{\pm \sqrt{0 + 96}}{2}$$

$$x = \frac{\pm \sqrt{96}}{2}$$

$$x = \frac{\pm \sqrt{16 \times 6}}{2}$$

$$x = \frac{\pm 4\sqrt{6}}{2}$$

$$x = \pm 2\sqrt{6}$$

$$\text{Solution set} = \{\pm 2\sqrt{6}\}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

161

Q.9. $\frac{x+4}{x-4} + \frac{x-4}{x+4} = \frac{10}{3}$ where $x \neq -4$

Sol. Multiplying by $3(x-4)(x+4)$

$$3(x+4)(x+4) + 3(x-4)(x-4) = 10(x-4)(x+4)$$

$$3(x^2 + 8x + 16) + 3(x^2 - 8x + 16) = 10(x^2 - 16)$$

$$3x^2 + 24x + 48 + 3x^2 - 24x + 48 = 10x^2 - 160$$

$$3x^2 + 3x^2 - 10x^2 + 24x - 24x + 48 + 48 + 160 = 0$$

$$-4x^2 + 256 = 0$$

$$4x^2 - 256 = 0$$

$$x^2 - 64 = 0 \quad (\text{Divided by 4})$$

Here

$$a = 1$$

$$b = 0$$

$$c = -64$$

(Formula)

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

By putting the values of a, b, c

$$x = \frac{-0 \pm \sqrt{(0)^2 - 4(1)(-64)}}{2(1)}$$

$$x = \frac{\pm \sqrt{0 + 256}}{2}$$

$$x = \frac{\pm \sqrt{256}}{2}$$

$$x = \frac{\pm 16}{2}$$

$$x = \pm 8$$

$$\text{Solution set} = \{8, -8\}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

162

Q.10. $\frac{1}{x-1} + \frac{1}{x-2} = \frac{2}{x-3}$ where $x \neq 1, 2, 3$

Sol. Multiplying by $(x-1)(x-2)(x-3)$

$$(x-1)(x-2)(x-3) \times \frac{1}{(x-1)} + (x-1)(x-2)(x-3) \times \frac{1}{(x-2)} = (x-1)(x-2)(x-3) \times \frac{2}{(x-3)}$$

$$(x-2)(x-3) + (x-1)(x-3) = (x-1)(x-2)(2)$$

$$x^2 - 5x + 6 + x^2 - 4x + 3 = 2(x^2 - 3x + 2)$$

$$2x^2 - 9x + 9 = 2x^2 - 6x + 4$$

$$-9x + 6x = 4 - 9$$

$$-3x = -5$$

$$x = \frac{-5}{-3}$$

$$x = \frac{5}{3}$$

Q.11. $(x+4)(x-1) + (x+5)(x+2) = 6$

Sol. $x^2 + 3x - 4 + x^2 + 7x + 10 = 6$

$$2x^2 + 10x - 4 + 10 - 6 = 0$$

$$2x^2 + 10x = 0$$

Here $a = 2$

$$b = 10$$

$$c = 0$$

(Formula) $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

By putting the values of a, b, c

$$x = \frac{-10 \pm \sqrt{(10)^2 - 4(2)(0)}}{2(2)}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

163

$$x = \frac{-10 \pm \sqrt{100}}{4}$$

$$x = \frac{-10 \pm 10}{4}$$

$$x = \frac{-10+10}{4}, \frac{-10-10}{4}$$

$$x = \frac{0}{4}, \frac{-20}{4}$$

$$x = 0, -5$$

Solution set = {0, -5}

Q.12 $(2x+4)^2 - (4x-6)^2 = 0$

Sol.

$$(4x^2 + 16x + 16) - (16x^2 - 48x + 36) = 0$$

$$-12x^2 + 64x - 20 = 0$$

(Dividing by -4)

$$3x^2 - 16x + 5 = 0$$

Here $a = 3$

$$b = -16$$

$$c = 5$$

$$\text{(Formula)} \quad x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

By putting the values of a, b, c

$$x = \frac{-(-16) \pm \sqrt{(-16)^2 - 4(3)(5)}}{2(3)}$$

$$x = \frac{16 \pm \sqrt{256 - 60}}{6}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

=====

Pilot Super One "General Math" 10th

164

$$x = \frac{16 \pm \sqrt{196}}{6}$$

$$x = \frac{16 \pm 14}{6}$$

$$x = \frac{16+14}{6}, \frac{16-14}{6}$$

$$x = \frac{30}{6}, \frac{2}{6}$$

$$x = 5, \frac{1}{3}$$

$$\text{Solution set} = \left\{ 5, \frac{1}{3} \right\}$$

Exercise 53

Q.1. Find two consecutive positive odd numbers such that the sum of their squares is 74.

Sol. Let 1st odd number = $2x + 1$

2nd odd number = $2x + 3$

According to statement

$$(2x+1)^2 + (2x+3)^2 = 74$$

$$4x^2 + 4x + 1 + 4x^2 + 12x + 9 = 74$$

$$8x^2 + 16x + 10 - 74 = 0$$

$$8x^2 + 16x - 64 = 0$$

$$x^2 + 2x - 8 = 0 \text{ (Divided by 8)}$$

$$x^2 + 4x - 2x - 8 = 0$$

$$(x^2 + 4x) - (2x + 8) = 0$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

165

$$x(x + 4) - 2(x + 4) = 0$$

$$x + 4(x - 2) = 0$$

$$\text{If } x - 2 = 0$$

$$\text{then } \boxed{x = 2}$$

$$\text{and if } x + 4 = 0$$

$$\text{then } \boxed{x = -4}$$

$$\text{when } x = 2 \text{ then 1st number} = 2x + 1$$

$$= 2(2) + 1$$

$$= 4 + 1$$

$$= 5$$

$$\text{and } 2^{\text{nd}} \text{ number} = 2x + 3$$

$$= 2(2) + 3$$

$$= 4 + 3$$

$$= 7$$

$$\text{Required odd number} = 5, 7$$

Q.2. Find two consecutive positive even numbers such that the sum of their squares is 164.

$$\text{Sol } \text{Let 1st number} = 2x$$

$$2^{\text{nd}} \text{ number} = 2x + 2$$

According to the statement

$$(2x)^2 + (2x + 2)^2 = 164$$

$$4x^2 + 4x^2 + 8x + 4 = 164$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

166

$$8x^2 + 8x + 4 - 164 = 0$$

$$8x^2 + 8x - 160 = 0$$

$$x^2 + x - 20 = 0 \quad (\text{Divided by } 8)$$

$$x^2 + 5x - 4x - 20 = 0$$

$$(x^2 + 5x) - (4x + 20) = 0$$

$$x(x + 5) - 4(x + 5) = 0$$

$$(x + 5)(x - 4) = 0$$

$$\text{If } x - 4 = 0$$

$$\text{then } x = 4$$

$$\text{and if } x + 5 = 0$$

$$\text{then } x = -5$$

when $x = 4$, then 1st number = $2x$

$$= 2(4) = 8$$

And 2nd number = $2x + 2$

$$= 2(4) + 2 = 8 + 2 = 10$$

Required even number = 8, 10

Q.3. The difference of two numbers is 9 and the product of the numbers is 162. Find the two numbers.

Sol. Let 1st number = x

2nd number = $x + 9$

According to the statement

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

=====

Pilot Super One "General Math" 10th

167

$$x^2 + 9x = 162$$

$$x^2 + 9x = 162$$

$$x^2 + 9x - 162 = 0$$

$$x^2 + 18x - 9x - 162 = 0$$

$$(x^2 + 18x) - (9x + 162) = 0$$

$$x(x + 18) - 9(x + 18) = 0$$

$$(x - 9)(x + 18) = 0$$

$$\text{If } x - 9 = 0$$

$$\text{then } \boxed{x = 9}$$

$$\text{and if } x + 18 = 0$$

$$\boxed{x = -18}$$

If $x = 9$, then 1st number = $x = 9$

$$2^{\text{nd}} \text{ number} = x + 9$$

$$= 9 + 9 = 18$$

$$\text{Required numbers} = \boxed{9, 18}$$

$$\text{If } x = -18$$

$$1^{\text{st}} \text{ number} = x = -18$$

$$2^{\text{nd}} \text{ number} = x + 9$$

$$= -18 + 9 = -9$$

$$\text{Required numbers} = \boxed{-18, -9}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

168

Q.4. The base and height of a triangle are $(x + 3)cm$ and $(2x - 5)cm$ respectively. If the area of the triangle is $20cm^2$, find x .

Sol. Length of base = $x + 3cm$

Length of altitude = $2x - 5cm$

Area of $\Delta = 20cm^2$

According to the statement

$$\frac{(x + 3)(2x - 5)}{2} = 20$$

$$\text{Area of } \Delta = \frac{B \times A}{2}$$

$$\frac{2x^2 + x - 15}{2} = 20$$

$$2x^2 + x - 15 = 2 \times 20$$

$$2x^2 + x - 15 = 40$$

$$2x^2 + x - 15 - 40 = 0$$

$$2x^2 + x - 55 = 0$$

$$2x^2 + 11x - 10x - 55 = 0$$

$$(2x^2 + 11x) - (10x + 55) = 0$$

$$x(2x + 11) - 5(2x + 11) = 0$$

$$(2x + 11)(x - 5) = 0$$

$$\text{If } x - 5 = 0$$

$$\text{then } \boxed{x = 5}$$

Therefore, Length of base = $x + 3$

$$= 5 + 3$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

169

$$= 8\text{cm}$$

$$\text{Length of altitude} = 2x - 5$$

$$= 2(5) - 5$$

$$= 10 - 5$$

$$= 5\text{cm}$$

$$\text{and if } 2x + 11 = 0$$

$$2x = -11$$

then

$$x = -\frac{11}{2}$$

Cancelling due to negativity.

Q.5. The perimeter and area of a rectangle are 22cm and 30cm^2 respectively. Find the length and breadth of the rectangle.

Sol. Rectangle's perimeter = 22cm

Rectangle's area = 30cm^2

Lct. Length of rectangle = x cm

Breadth of rectangle = $\frac{30}{x}$ cm

$$\therefore \text{Perimeter} = (\text{length} + \text{breadth}) \times 2$$

$$22 = \left(\frac{30}{x} + x \right) \times 2$$

$$11 = x + \frac{30}{x} \quad (\text{Divided by 2})$$

$$11x = 30 + x^2 \quad (\text{Multiplying by } x)$$

$$0 = 30 + x^2 - 11x$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

170

$$x^2 - 11x + 30 = 0$$

$$x^2 - 5x - 6x + 30 = 0$$

$$(x^2 - 5x) - (6x - 30) = 0$$

$$x(x - 5) - 6(x - 5) = 0$$

$$(x - 5)(x - 6) = 0$$

$$\text{If } x - 5 = 0$$

$$\text{then } \boxed{x = 5}$$

$$\text{Length of rectangle} = x$$

$$= 5\text{cm}$$

$$\text{and Breadth of rectangle} = \frac{30}{x}$$

$$= \frac{30}{5}$$

$$= 6\text{ cm}$$

$$\text{Length of rectangle} = 5\text{cm}$$

$$\text{Breadth of rectangle} = 6\text{cm}$$

$$\text{and if } x - 6 = 0$$

$$\text{then } \boxed{x = 6}$$

$$\text{Length of rectangle} = x$$

$$= 6\text{cm}$$

$$\text{and Breadth of rectangle} = \frac{30}{x}$$

$$= \frac{30}{6}$$

$$= 5\text{cm}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

171

Length of rectangle = 6cm

So, Breadth of rectangle = 5cm

Q.6. The product of two consecutive positive numbers is 156. Find the numbers.

Sol. Let, 1st number = x

2nd number = $x + 1$

According to the statement

$$(x)(x + 1) = 156$$

$$x^2 + x = 156$$

$$x^2 + x - 156 = 0$$

$$x^2 + 13x - 12x - 156 = 0$$

$$(x^2 + 13x) - (12x + 156) = 0$$

$$x(x + 13) - 12(x + 13) = 0$$

$$(x - 12)(x + 13) = 0$$

if $x - 12 = 0$

then

$$x = 12$$

Required numbers = $x, x + 1$

$$= 12, 12 + 1$$

$$= \boxed{12, 13}$$

and if $x + 13 = 0$

then

$$x = -13$$

Required numbers = $x, x + 1$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

172

$$= -13, -13 + 1$$

$$= \boxed{-13, -12}$$

Q.7. Find two consecutive positive odd numbers given
 that the difference between their reciprocals is $\frac{2}{63}$.

Sol. Suppose that 1st number = $2x + 1$

$$2^{\text{nd}} \text{ number} = 2x + 3$$

According to given condition

$$\frac{1}{2x+1} - \frac{1}{2x+3} = \frac{2}{63}$$

Multiplying by $(2x+1)(2x+3)(63)$

$$\begin{aligned} 63(2x+1)(2x+3) \times \frac{1}{(2x+1)} - 63(2x+1)(2x+3) \times \frac{1}{(2x+3)} \\ = \frac{2}{63} \times 63(2x+1)(2x+3) \end{aligned}$$

$$63(2x+3) - 63(2x+1) = 2(2x+1)(2x+3)$$

$$126x + 189 - 126x - 63 = 2(4x^2 + 8x + 3)$$

$$126 = 8x^2 + 16x + 6$$

$$8x^2 + 16x + 6 - 126 = 0$$

$$8x^2 + 16x - 120 = 0$$

$$x^2 + 2x - 15 = 0 \quad (\text{Dividing by 8})$$

$$x^2 + 5x - 3x - 15 = 0$$

$$(x^2 + 5x) - (3x + 15) = 0$$

$$x(x+5) - 3(x+5) = 0$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 5)

Pilot Super One "General Math" 10th

173

$$(x + 5)(x - 3) = 0$$

If $x - 3 = 0$

then $x = 3$

Required numbers: 1st number = $2x + 1$ required numbers

$$= 2(3) + 1$$

$$= 6 + 1$$

$$\boxed{1st\ number = 7}$$

$$2nd\ number = 2x + 3$$

$$= 2(3) + 3$$

$$= 6 + 3$$

$$\boxed{2nd\ number = 9}$$

and if $x + 5 = 0$

then $x = -5$

Cancelling due to negativity.

Q.8. The sum of the two positive number is 12 and the sum of whose squares is 80. Find the numbers.

Sol. Suppose that 1st number = x

$$2nd\ number = 12 - x$$

According to given condition

$$(x)^2 + (12 - x)^2 = 80$$

$$x^2 + 144 + x^2 - 24x = 80$$

$$2x^2 - 24x + 144 - 80 = 0$$

$$2x^2 - 24x + 64 = 0$$

$$x^2 - 12x + 32 = 0 \quad (\text{Dividing by 2})$$

$$x^2 - 4x - 8x + 32 = 0$$

$$x(x - 4) - 8(x - 4) = 0$$

$$(x - 4)(x - 8) = 0$$

$$\text{If } x - 4 = 0$$

$$\text{then } \boxed{x = 4}$$

$$\text{Required numbers: 1st number} = \boxed{x = 4}$$

$$\text{2nd number} = 12 - x$$

$$\boxed{\text{2nd number} = 12 - 4 = 8}$$

$$\text{and if } x - 8 = 0$$

$$x = 8$$

$$\text{Hence } \boxed{\text{1st number} = 8}$$

$$\text{2nd number} = 12 - x$$

$$= 12 - 8$$

$$\boxed{\text{2nd number} = 4}$$

$$\text{Required numbers} = 8, 4$$

$$\text{or Required numbers} = 4, 8$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

180

Exercise 6.1

With the help of the given matrices answer the questions from 1 to 3.

$$A = \begin{bmatrix} 2 & -2 \\ -5 & 0 \end{bmatrix}, B = \begin{bmatrix} -3 & -2 \\ 0 & 4 \end{bmatrix}, C = \begin{bmatrix} 3 \\ -1 \\ 0 \end{bmatrix},$$
$$D = \begin{bmatrix} -3 & 2 & 0 \\ 0 & 1 & 5 \\ 4 & -2 & 2 \end{bmatrix}, E = [-3 \ 2 \ 0], F = \begin{bmatrix} -3 & 4 \\ 0 & 5 \\ 3 & -1 \end{bmatrix}$$

- 1- What are the orders of matrices A, C and F?
- 2- What are the orders of matrices B, D and E?
- 3- What element is in the second row and third column of matrix D?

Answers:

- 1.(i) (R) Number of rows in matrix A = 2
(C) Number of columns in matrix B = 2
Order of matrix A (R × C) = 2-by-2
- (ii) (R) Number of rows in matrix C = 3
(C) Number of columns in matrix C = 1
Order of matrix C (R × C) = 3-by-1
- (iii) (R) Number of rows in matrix F = 3
(C) Number of columns in matrix F = 2

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

181

Order of matrix $F(R \times C) = 3\text{-by-}2$

- 2.(i) (R) Number of rows in matrix $B = 2$
 (C) Number of columns in matrix $B = 2$
 Order of matrix $B(R \times C) = 2\text{-by-}2$

- (ii) (R) Number of rows in matrix $D = 3$
 (C) Number of columns in matrix $D = 3$
 Order of matrix $D(R \times C) = 3\text{-by-}3$

- (iii) (R) Number of rows in matrix $E = 1$
 (C) Number of columns in matrix $E = 3$
 Order of matrix $E(R \times C) = 1\text{-by-}3$

3. The element is in the second row and third column of matrix D is 5.

- Q.4. Which of the following matrices are equal and which of them are not?

$$A = \begin{bmatrix} 4 \end{bmatrix}, B = \begin{bmatrix} 1 & 2 \end{bmatrix}, C = \begin{bmatrix} 6 \\ 9 \end{bmatrix}, D = \begin{bmatrix} 2+2 \end{bmatrix}$$

$$E = \begin{bmatrix} 3+3 \\ 8+1 \end{bmatrix}, F = \begin{bmatrix} 5 & 4 \\ 5 & 2 \end{bmatrix}, G = \begin{bmatrix} 1 & 3 \\ 6 & 8 \end{bmatrix}$$

$$H = \begin{bmatrix} 1 & 2 & 5 \\ 0 & 3 & 4 \\ 2 & 6 & 3 \end{bmatrix}, I = \begin{bmatrix} 1 & 3 \\ 6 & 7 \end{bmatrix}, J = \begin{bmatrix} 1 & 3 \\ 6 & \frac{16}{2} \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

182

$$K = \begin{bmatrix} 1 & 2 & 3+2 \\ 0 & 3 & 4 \\ 2 & 4+2 & 3 \end{bmatrix}, L = \begin{bmatrix} 1 & 3 & 5 \\ 0 & 3 & 4 \\ 2 & 6 & 3 \end{bmatrix}$$

Answers:

(i) $B = F, G = J, H = K, C = E, A = D$

(ii) I and L are not equal to any matrix.

TYPES OF MATRICES

(i) **Row Matrix:**

A matrix with only one row is called a row matrix.

For example: $B = [2 \ 3 \ 4]$ is of order 1 - by - 3.

(ii) **Column Matrix:**

A matrix with only one column is called a column matrix.

For example: $D = \begin{bmatrix} 3 \\ 2 \\ 7 \end{bmatrix}$ is of order 3 - by - 1.

(iii) **Rectangular Matrix:**

If in a matrix, the number of rows and the number of columns are not equal, then the matrix is called a rectangular matrix.

For example: $C = \begin{bmatrix} 1 & 2 & 7 \\ 3 & 4 & 5 \end{bmatrix}$

(iv) **Square Matrix:**

If a matrix has equal number of rows and columns, it is called a square matrix.

For example: $Q = \begin{bmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \\ 3 & 5 & 7 \end{bmatrix}$

(v) **Zero or Null Matrix:**

If all the elements in a matrix are zeros, it is called a zero matrix or null matrix. A null matrix is denoted by the letter O .

For example: $O = [0]$ is of order 1 - by - 1.

$O = [0 \ 0]$ is of order 1 - by - 2.

(vi) **Diagonal Matrix:**

A square matrix in which all the elements except at least the one element in the diagonal are zeros is called a diagonal matrix.

For example: $A = \begin{bmatrix} 3 & 0 \\ 0 & 4 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 0 \\ 0 & 3 \end{bmatrix}$

(vii) **Scalar Matrix:**

A diagonal matrix having equal elements is called a scalar matrix.

For example: $B = \begin{bmatrix} \sqrt{2} & 0 \\ 0 & \sqrt{2} \end{bmatrix}$

(viii) **Unit Matrix or Identity Matrix:**

A scalar matrix having each element equal to 1 is called a unit or identity matrix. Identity or unit matrix is generally denoted by I .

○ For example: $I = [1]$ $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

(ix) **Transpose of a Matrix:**

If A is a matrix of order (m - by - n), then a matrix (n - by - m) obtained by interchanging the rows and columns of A is called the transpose of A . It is denoted by A' .

For example: $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then $A' = \begin{bmatrix} a & c \\ b & d \end{bmatrix}$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

184

(x) **Symmetric Matrix:**

A square matrix A is called symmetric if $A' = A$

For example: $A = \begin{bmatrix} p & q \\ q & r \end{bmatrix}$, and $A' = \begin{bmatrix} p & q \\ q & r \end{bmatrix}$

(xi) **Skew-Symmetric Matrix:**

A square matrix A is called skew symmetric (or anti-symmetric) if $A' = -A$

For example: $A = \begin{bmatrix} 0 & -2 & 3 \\ 2 & 0 & 4 \\ -3 & -4 & 0 \end{bmatrix}$

$$A' = \begin{bmatrix} 0 & 2 & -3 \\ -2 & 0 & -4 \\ 3 & 4 & 0 \end{bmatrix} = - \begin{bmatrix} 0 & -2 & 3 \\ 2 & 0 & 4 \\ -3 & -4 & 0 \end{bmatrix} = -A$$

$A' = -A$ Hence A is skew symmetric.

EXERCISE 6.2

- 1- Identify row matrices, column matrices, square matrices, and rectangular matrices in the following matrices.

$$A = [3 \ 1 \ 1 \ 1], B = \begin{bmatrix} 5+2 & 4 \\ 2 & 6 \end{bmatrix}, C = \begin{bmatrix} a+x \\ b+y \end{bmatrix},$$

$$D = \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, E = \begin{bmatrix} x & -2 \\ b & 5 \end{bmatrix}, F = \begin{bmatrix} 1 & 3 & 2 \\ 2 & 4 & 5 \\ 1 & -5 & 0 \end{bmatrix},$$

$$G = \begin{bmatrix} 1 & 2 & 4 \\ 5 & 7 & 8 \end{bmatrix}, H = [0]$$

Solution:

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

185

$$A = \begin{bmatrix} 3 & 1 & 1 & 1 \end{bmatrix} \quad \text{Row Matrix}$$

$$C = \begin{bmatrix} (a+x) \\ (b+y) \end{bmatrix} \quad \text{Column Matrix}$$

$$B = \begin{bmatrix} (5+2) & 4 \\ 2 & 6 \end{bmatrix}, D = \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix} \quad \text{Square Matrix}$$

$$E = \begin{bmatrix} x & -2 \\ h & 5 \end{bmatrix}, F = \begin{bmatrix} 1 & 3 & 2 \\ 2 & 4 & 5 \\ 1 & -5 & 0 \end{bmatrix}, H = [0]$$

$$C = \begin{bmatrix} (a+x) \\ (b+y) \end{bmatrix}, G = \begin{bmatrix} 1 & 2 & 4 \\ 5 & 7 & 8 \end{bmatrix} \quad \text{Rectangle Matrix}$$

2. Identify, diagonal matrices, scalar matrices, identity matrices.

$$A = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 7 & 0 \\ 0 & 0 & 7 \end{bmatrix}, C = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 4 \end{bmatrix},$$

$$D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, E = \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix}, F = \begin{bmatrix} 8 & 0 \\ 0 & 0 \end{bmatrix},$$

$$G = \begin{bmatrix} k & 0 & 0 \\ 0 & k & 0 \\ 0 & 0 & k \end{bmatrix}$$

Solution:

$$G = \begin{bmatrix} k & 0 & 0 \\ 0 & k & 0 \\ 0 & 0 & k \end{bmatrix} \quad \text{Diagonal Matrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

186

$$A = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, C = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 4 \end{bmatrix}, F = \begin{bmatrix} 8 & 0 \\ 0 & 0 \end{bmatrix}$$

Scalar Matrix

$$B = \begin{bmatrix} 7 & 0 & 0 \\ 0 & 7 & 0 \\ 0 & 0 & 7 \end{bmatrix}, E = \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix}, G = \begin{bmatrix} k & 0 & 0 \\ 0 & k & 0 \\ 0 & 0 & k \end{bmatrix}$$

$$D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Identity Matrix

3. Find transpose of the following matrices.

$$A = \begin{bmatrix} 3 & 4 \\ -1 & 4 \end{bmatrix}, B = \begin{bmatrix} -3 & -2 \\ -1 & 4 \end{bmatrix},$$

$$C = \begin{bmatrix} a & -b \\ c & d \end{bmatrix}, D = \begin{bmatrix} l & m & n \\ p & q & r \\ a & b & c \end{bmatrix}$$

Solution:

$$A = \begin{bmatrix} 3 & 4 \\ -1 & 4 \end{bmatrix}, A' = \begin{bmatrix} 3 & -1 \\ 4 & 4 \end{bmatrix}$$

$$B = \begin{bmatrix} -3 & -2 \\ -1 & 4 \end{bmatrix}, B' = \begin{bmatrix} -3 & -1 \\ -2 & 4 \end{bmatrix}$$

$$C = \begin{bmatrix} a & -b \\ c & d \end{bmatrix}, C' = \begin{bmatrix} a & c \\ -b & d \end{bmatrix}$$

..

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

187

$$D = \begin{bmatrix} l & m & n \\ p & q & r \\ a & b & c \end{bmatrix}, D' = \begin{bmatrix} l & p & a \\ m & q & b \\ n & r & c \end{bmatrix}$$

4. Identify all row matrices, if:

$$A = [3 \ 4 \ 5], B = \begin{bmatrix} -1 & 3 \\ 4 & 6 \end{bmatrix}, C = [e \ f \ g],$$

$$D = \begin{bmatrix} 3 & 7 & 5 \\ 4 & 6 & 2 \\ 1 & 9 & 8 \end{bmatrix}, F = \begin{bmatrix} 1 & 4 & 6 \\ 3 & 7 & 3 \end{bmatrix}$$

Solution: Row Matrix

$$A = [3 \ 4 \ 5], C = [e \ f \ g]$$

5. Identify all column matrices, if:

$$A = \begin{bmatrix} 3 \\ 6 \\ 10 \end{bmatrix}, B = \begin{bmatrix} 2 & 3 \\ 6 & 5 \\ 4 & 7 \end{bmatrix}, C = \begin{bmatrix} a \\ b \\ c \end{bmatrix},$$

$$D = \begin{bmatrix} 2 & -1 & 3 \\ 4 & 6 & 5 \\ -2 & 3 & 4 \end{bmatrix}, E = \begin{bmatrix} 5 \\ 7 \\ -4 \end{bmatrix}, F = [9 \ 7 \ 1]$$

Solution: Column Matrix

$$A = \begin{bmatrix} 3 \\ 6 \\ 10 \end{bmatrix}, C = \begin{bmatrix} a \\ b \\ c \end{bmatrix}, E = \begin{bmatrix} 5 \\ 7 \\ -4 \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

188

6. Identify all column matrices, if:

$$A = \begin{bmatrix} 1 & 3 \\ 2 & 2 \end{bmatrix}, B = \begin{bmatrix} 3 & -2 & 4 \\ 1 & 6 & 5 \\ 7 & 3 & 4 \end{bmatrix}, C = \begin{bmatrix} 1 \\ 2 \\ 9 \end{bmatrix},$$
$$D = \begin{bmatrix} 7 & 8 \\ 6 & 5 \end{bmatrix}, E = [3 \ 7 \ 5], F = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

Solution: Column Matrix

$$C = \begin{bmatrix} 1 \\ 2 \\ 9 \end{bmatrix}$$

7. Identify all 3-by-3 square matrices, if:

$$A = \begin{bmatrix} 2 & -3 & 6 \\ 1 & 5 & 4 \\ 3 & 6 & -3 \end{bmatrix}, B = \begin{bmatrix} 2 \\ 4 \\ 7 \end{bmatrix}, C = [7 \ 3 \ 4]$$

Solution:

$$A = \begin{bmatrix} 2 & -3 & 6 \\ 1 & 5 & 4 \\ 3 & 6 & -3 \end{bmatrix}$$

A Scalar Multiplication

Any element from the set of real numbers is also called a scalar. We define the product of a matrix A and a scalar k , denoted by kA , to be the matrix formed by multiplying each element of A by k .

$$\text{For example: } A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, \text{ then } kA = \begin{bmatrix} ka & kb \\ kc & kd \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

189

Laws of Addition of Matrices

Commutative Law:

For any two matrices A and B of the same order

$$A + B = B + A$$

This law is called commutative law of matrices with respect to addition.

Associative Law:

For three matrices A, B and C of same order,

$$(A + B) + C = A + (B + C)$$

This law is called associative law of matrices with respect to addition.

Additive Identity of Matrices

$$A + O = O + A = A$$

Additive Inverse of a Matrix

If two matrices A and B are such that their sum (A + B) is a zero matrix, then A and B are called additive inverse of each other.

For example: $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, and $B = \begin{bmatrix} -a & -b \\ -c & -d \end{bmatrix}$

Exercise 6.3

1- If $A = \begin{bmatrix} 2 & 3 & 4 \\ 1 & 5 & 5 \\ 4 & 9 & 3 \end{bmatrix}$ $B = \begin{bmatrix} 0 & 1 & 5 \\ 2 & 3 & 6 \\ 1 & 4 & -2 \end{bmatrix}$

Find

- (i) $A + B$ (ii) $A - B$ (iii) $B - A$
 (iv) $2A + 3B$ (v) $3A - 4B$ (vi) $A - 2B$

Solutions:

(i) $A + B = \begin{bmatrix} 2 & 3 & 4 \\ 1 & 5 & 5 \\ 4 & 9 & 3 \end{bmatrix} + \begin{bmatrix} 0 & 1 & 5 \\ 2 & 3 & 6 \\ 1 & 4 & -2 \end{bmatrix}$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

190

$$= \begin{bmatrix} 2+0 & 3+1 & 4+5 \\ 1+2 & 5+3 & 5+6 \\ 4+1 & 9+4 & 3-2 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 4 & 9 \\ 3 & 8 & 11 \\ 5 & 13 & 1 \end{bmatrix}$$

(ii)

$$A-B = \begin{bmatrix} 2 & 3 & 4 \\ 1 & 5 & 5 \\ 4 & 9 & 3 \end{bmatrix} - \begin{bmatrix} 0 & 1 & 5 \\ 2 & 3 & 6 \\ 1 & 4 & -2 \end{bmatrix}$$

$$= \begin{bmatrix} 2-0 & 3-1 & 4-5 \\ 1-2 & 5-3 & 5-6 \\ 4-1 & 9-4 & 3+2 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 2 & -1 \\ -1 & 2 & -1 \\ 3 & 5 & 5 \end{bmatrix}$$

(iii)

$$B-A = \begin{bmatrix} 0 & 1 & 5 \\ 2 & 3 & 6 \\ 1 & 4 & -2 \end{bmatrix} - \begin{bmatrix} 2 & 3 & 4 \\ 1 & 5 & 5 \\ 4 & 9 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 0-2 & 1-3 & 5-4 \\ 2-1 & 3-5 & 6-5 \\ 1-4 & 4-9 & -2-3 \end{bmatrix}$$

$$= \begin{bmatrix} -2 & -2 & 1 \\ 1 & -2 & 1 \\ -3 & -5 & -5 \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pink Super One "General Math" 10th

191

$$\begin{aligned}
 \text{(iv)} \quad 2A + 3B &= 2 \begin{bmatrix} 2 & 3 & 4 \\ 1 & 5 & 5 \\ 4 & 9 & 3 \end{bmatrix} + 3 \begin{bmatrix} 0 & 1 & 5 \\ 2 & 3 & 6 \\ 1 & 4 & -2 \end{bmatrix} \\
 &= \begin{bmatrix} 4 & 6 & 8 \\ 2 & 10 & 10 \\ 8 & 18 & 6 \end{bmatrix} + \begin{bmatrix} 0 & 3 & 15 \\ 6 & 9 & 18 \\ 3 & 12 & -6 \end{bmatrix} \\
 &= \begin{bmatrix} 4+0 & 6+3 & 8+15 \\ 2+6 & 10+9 & 10+18 \\ 8+3 & 18+12 & 6-6 \end{bmatrix} \\
 &= \begin{bmatrix} 4 & 9 & 23 \\ 8 & 19 & 28 \\ 11 & 30 & 0 \end{bmatrix}
 \end{aligned}$$

$$\begin{aligned}
 \text{(v)} \quad 3A - 4B &= 3 \begin{bmatrix} 2 & 3 & 4 \\ 1 & 5 & 5 \\ 4 & 9 & 3 \end{bmatrix} - 4 \begin{bmatrix} 0 & 1 & 5 \\ 2 & 3 & 6 \\ 1 & 4 & -2 \end{bmatrix} \\
 &= \begin{bmatrix} 6 & 9 & 12 \\ 3 & 15 & 15 \\ 12 & 27 & 9 \end{bmatrix} - \begin{bmatrix} 0 & 4 & 20 \\ 8 & 12 & 24 \\ 4 & 16 & -8 \end{bmatrix} \\
 &= \begin{bmatrix} 6-0 & 9-4 & 12-20 \\ 3-8 & 15-12 & 15-24 \\ 12-4 & 27-16 & 9+8 \end{bmatrix} \\
 &= \begin{bmatrix} 6 & 5 & -8 \\ -5 & 3 & -9 \\ 8 & 11 & 17 \end{bmatrix}
 \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

192

$$\begin{aligned}
 \text{(vi)} \quad A-2B &= \begin{bmatrix} 2 & 3 & 4 \\ 1 & 5 & 5 \\ 4 & 9 & 3 \end{bmatrix} - 2 \begin{bmatrix} 0 & 1 & 5 \\ 2 & 3 & 6 \\ 1 & 4 & -2 \end{bmatrix} \\
 &= \begin{bmatrix} 2 & 3 & 4 \\ 1 & 5 & 5 \\ 4 & 9 & 3 \end{bmatrix} - \begin{bmatrix} 0 & 2 & 10 \\ 4 & 6 & 12 \\ 2 & 8 & -4 \end{bmatrix} \\
 &= \begin{bmatrix} 2-0 & 3-2 & 4-10 \\ 1-4 & 5-6 & 5-12 \\ 4-2 & 9-8 & 3+4 \end{bmatrix} \\
 &= \begin{bmatrix} 2 & 1 & -6 \\ -3 & -1 & -7 \\ 2 & 1 & 7 \end{bmatrix}
 \end{aligned}$$

2- Find the additive inverse of the following matrices.

$$A = \begin{bmatrix} 4 & 3 \\ 2 & 6 \end{bmatrix}, B = \begin{bmatrix} \sqrt{2} & 3 \\ 4 & \sqrt{3} \end{bmatrix}, C = \begin{bmatrix} 1 \\ -7 \\ 4 \end{bmatrix}$$

$$D = \begin{bmatrix} 1 & 0 & -2 \\ 0 & 3 & 4 \\ 2 & -1 & -3 \end{bmatrix}, E = [2 \ 5 \ -3]$$

Sol:

Matrix	Additive Inverse
$A = \begin{bmatrix} 4 & 3 \\ 2 & 6 \end{bmatrix},$	$-A = \begin{bmatrix} -4 & -3 \\ -2 & -6 \end{bmatrix},$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

193

$$B = \begin{bmatrix} \sqrt{2} & 3 \\ 4 & \sqrt{3} \end{bmatrix}, \quad -B = \begin{bmatrix} -\sqrt{2} & -3 \\ -4 & -\sqrt{3} \end{bmatrix},$$

$$C = \begin{bmatrix} 1 \\ -7 \\ 4 \end{bmatrix}, \quad -C = \begin{bmatrix} -1 \\ 7 \\ -4 \end{bmatrix},$$

$$D = \begin{bmatrix} 1 & 0 & -2 \\ 0 & 3 & 4 \\ 2 & -1 & -3 \end{bmatrix}, \quad -D = \begin{bmatrix} -1 & 0 & 2 \\ 0 & -3 & -4 \\ -2 & 1 & 3 \end{bmatrix},$$

$$E = [2 \ 5 \ -3] \quad -E = [-2 \ -5 \ 3]$$

3. If $A = \begin{bmatrix} 2 & 3 \\ 1 & 5 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 7 \\ 4 & 6 \end{bmatrix}$ then show that

(i) $4A - 3A = A$ (ii) $3B - 3A = 3(B - A)$

Sol:

$$\begin{aligned} \text{(i)} \quad 4A - 3A &= 4 \begin{bmatrix} 2 & 3 \\ 1 & 5 \end{bmatrix} - 3 \begin{bmatrix} 2 & 3 \\ 1 & 5 \end{bmatrix} \\ &= \begin{bmatrix} 8 & 12 \\ 4 & 20 \end{bmatrix} - \begin{bmatrix} 6 & 9 \\ 3 & 15 \end{bmatrix} \\ &= \begin{bmatrix} 8-6 & 12-9 \\ 4-3 & 20-15 \end{bmatrix} \\ 4A - 3A &= \begin{bmatrix} 2 & 3 \\ 1 & 5 \end{bmatrix} = A \quad (\text{Proved}) \end{aligned}$$

(ii) $3B - 3A = 3(B - A)$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pakistan Super One "General Math" 10th

194

$$\begin{aligned} \text{L.H.S. } 3B - 3A &= 3 \begin{bmatrix} 1 & 7 \\ 4 & 6 \end{bmatrix} - 3 \begin{bmatrix} 2 & 3 \\ 1 & 5 \end{bmatrix} \\ &= \begin{bmatrix} 3 & 21 \\ 12 & 18 \end{bmatrix} - \begin{bmatrix} 6 & 9 \\ 3 & 15 \end{bmatrix} \\ &= \begin{bmatrix} 3-6 & 21-9 \\ 12-3 & 18-15 \end{bmatrix} \\ &= \begin{bmatrix} -3 & 12 \\ 9 & 3 \end{bmatrix} \dots\dots\dots (i) \end{aligned}$$

$$\begin{aligned} \text{R.H.S. } 3(B - A) &= 3 \left(\begin{bmatrix} 1 & 7 \\ 4 & 6 \end{bmatrix} - \begin{bmatrix} 2 & 3 \\ 1 & 5 \end{bmatrix} \right) \\ &= 3 \left(\begin{bmatrix} 1-2 & 7-3 \\ 4-1 & 6-5 \end{bmatrix} \right) \\ &= 3 \begin{bmatrix} -1 & 4 \\ 3 & 1 \end{bmatrix} = \begin{bmatrix} -3 & 12 \\ 9 & 3 \end{bmatrix} \dots\dots\dots (ii) \end{aligned}$$

$$3B - 3A = 3(B - A) \text{ From (i) and (ii)}$$

4. Find x and y if $\begin{bmatrix} x+3 & 1 \\ -3 & 3y-4 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ -3 & 2 \end{bmatrix}$

Sol: 

$$x + 3 = 2$$

$$x = 2 - 3$$

$$\boxed{x = -1}$$

and

$$3y - 4 = 2$$

$$3y = 2 + 4$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

195

$$3y = 6$$

$$y = \frac{6}{3}$$

$$\boxed{y = 2}$$

5. If $A = \begin{bmatrix} 1 & 3 \\ 4 & 5 \end{bmatrix}$, $B = \begin{bmatrix} 4 & 7 \\ 6 & 5 \end{bmatrix}$, $C = \begin{bmatrix} 2 & 6 \\ 3 & -2 \end{bmatrix}$ then

prove that,

(i) $A + B = B + A$ (ii) $A + (B + C) = (A + B) + C$

(i) $A + B = B + A$

Sol:
$$A + B = \begin{bmatrix} 1 & 3 \\ 4 & 5 \end{bmatrix} + \begin{bmatrix} 4 & 7 \\ 6 & 5 \end{bmatrix}$$

$$= \begin{bmatrix} 1+4 & 3+7 \\ 4+6 & 5+5 \end{bmatrix}$$

$$= \begin{bmatrix} 5 & 10 \\ 10 & 10 \end{bmatrix} \dots\dots (i)$$

and $B + A = \begin{bmatrix} 4 & 7 \\ 6 & 5 \end{bmatrix} + \begin{bmatrix} 1 & 3 \\ 4 & 5 \end{bmatrix}$

$$= \begin{bmatrix} 4+1 & 7+3 \\ 6+4 & 5+5 \end{bmatrix}$$

$$= \begin{bmatrix} 5 & 10 \\ 10 & 10 \end{bmatrix} \dots\dots (ii)$$

From (i) and (ii) $A + B = B + A$

(ii) $A + (B + C) = (A + B) + C$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

196

$$\begin{aligned} A + (B + C) &= \begin{bmatrix} 1 & 3 \\ 4 & 5 \end{bmatrix} + \left(\begin{bmatrix} 4 & 7 \\ 6 & 5 \end{bmatrix} + \begin{bmatrix} 2 & 6 \\ 3 & -2 \end{bmatrix} \right) \\ &= \begin{bmatrix} 1 & 3 \\ 4 & 5 \end{bmatrix} + \begin{bmatrix} 4+2 & 7+6 \\ 6+3 & 5-2 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 3 \\ 4 & 5 \end{bmatrix} + \begin{bmatrix} 6 & 13 \\ 9 & 3 \end{bmatrix} \\ &= \begin{bmatrix} 1+6 & 3+13 \\ 4+9 & 5+3 \end{bmatrix} = \begin{bmatrix} 7 & 16 \\ 13 & 8 \end{bmatrix} \dots\dots (i) \end{aligned}$$

and

$$\begin{aligned} (A + B) + C &= \left(\begin{bmatrix} 1 & 3 \\ 4 & 5 \end{bmatrix} + \begin{bmatrix} 4 & 7 \\ 6 & 5 \end{bmatrix} \right) + \begin{bmatrix} 2 & 6 \\ 3 & -2 \end{bmatrix} \\ &= \begin{bmatrix} 1+4 & 3+7 \\ 4+6 & 5+5 \end{bmatrix} + \begin{bmatrix} 2 & 6 \\ 3 & -2 \end{bmatrix} \\ &= \begin{bmatrix} 5 & 10 \\ 10 & 10 \end{bmatrix} + \begin{bmatrix} 2 & 6 \\ 3 & -2 \end{bmatrix} \\ &= \begin{bmatrix} 5+2 & 10+6 \\ 10+3 & 10-2 \end{bmatrix} = \begin{bmatrix} 7 & 16 \\ 13 & 8 \end{bmatrix} \dots\dots (ii) \end{aligned}$$

From (i) and (ii)

$$A + (B + C) = (A + B) + C$$

6- Solve the matrix equation for X.

$$3X - 2A = B \text{ if } A = \begin{bmatrix} 2 & 3 \\ -4 & 1 \end{bmatrix} \text{ and } B = \begin{bmatrix} 2 & -3 \\ 4 & 4 \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

197

Solution:

$$3X - 2A = B$$

$$3X = B + 2A$$

$$3X = \begin{bmatrix} 2 & -3 \\ 4 & 4 \end{bmatrix} + 2 \begin{bmatrix} 2 & 3 \\ -4 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & -3 \\ 4 & 4 \end{bmatrix} + \begin{bmatrix} 4 & 6 \\ -8 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 2+4 & -3+6 \\ 4-8 & 4+2 \end{bmatrix}$$

$$3X = \begin{bmatrix} 6 & 3 \\ -4 & 6 \end{bmatrix}$$

$$X = \frac{1}{3} \begin{bmatrix} 6 & 3 \\ -4 & 6 \end{bmatrix}$$

$$= \begin{bmatrix} 6 \times \frac{1}{3} & 3 \times \frac{1}{3} \\ -4 \times \frac{1}{3} & 6 \times \frac{1}{3} \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 1 \\ -\frac{4}{3} & 2 \end{bmatrix}$$

7- Find a, b, c, d, e and f such that

$$\begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix} - \begin{bmatrix} 3 & -2 & 1 \\ 5 & 0 & -4 \end{bmatrix} = \begin{bmatrix} -1 & -2 & 3 \\ -2 & 4 & 6 \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

198

Sol:
$$\begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix} \cdot \begin{bmatrix} 3 & -2 & 1 \\ 5 & 0 & -4 \end{bmatrix} = \begin{bmatrix} -1 & -2 & 3 \\ -2 & 4 & 6 \end{bmatrix}$$

$$\begin{bmatrix} a+3 & b-2 & c+1 \\ d+5 & e-0 & f-4 \end{bmatrix} = \begin{bmatrix} -1 & -2 & 3 \\ -2 & 4 & 6 \end{bmatrix}$$

Hence

$$a+3 = -1, \quad b-2 = -2, \quad c+1 = 3$$

$$a = -1 - 3 \quad b = -2 + 2 \quad c = 3 - 1$$

$$\boxed{a = -4}$$

$$\boxed{b = 0}$$

$$\boxed{c = 2}$$

and

$$d+5 = -2, \quad e-0 = 4, \quad f-4 = 6$$

$$d = -2 - 5$$

$$\boxed{e = 4}$$

$$f = 6 + 4$$

$$\boxed{d = -7}$$

$$\boxed{f = 10}$$

8- Find w, x, y, z such that

$$\begin{bmatrix} w & x \\ y & z \end{bmatrix} + \begin{bmatrix} 3 & 0 \\ -1 & 5 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 6 & -3 \end{bmatrix}$$

Sol:
$$\begin{bmatrix} w+3 & x+0 \\ y-1 & z+5 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 6 & -3 \end{bmatrix}$$

Hence $w+3 = 2, \quad x+0 = 1$

$$w = 2 - 3$$

$$\boxed{x = 1}$$

$$\boxed{w = -1}$$

$$y-1 = 6,$$

$$z+5 = -3$$

$$y = 6 + 1$$

$$z = -3 - 5$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

199

$$y = 7$$

$$z = -8$$

9. If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ then what is the additive inverse of A ?

Sol: $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$

$$-A = -\begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

(Additive Inverse of A)

$$= \begin{bmatrix} -a & -b \\ -c & -d \end{bmatrix}$$

10. Given that $A = \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix}$ verify that $A^2 - 4A + 5I = 0$.

Sol: $A = \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix}$

$$L.H.S = A^2 - 4A + 5I$$

$$= \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix} - 4 \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix} + 5 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1-2 & -1-3 \\ 2+6 & -2+9 \end{bmatrix} - \begin{bmatrix} 4 & -4 \\ 8 & 12 \end{bmatrix} + \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix}$$

$$= \begin{bmatrix} -1 & -4 \\ 8 & 7 \end{bmatrix} - \begin{bmatrix} 4 & -4 \\ 8 & 12 \end{bmatrix} + \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix}$$

$$= \begin{bmatrix} -1-4 & -4+4 \\ 8-8 & 7-12 \end{bmatrix} + \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

200

$$= \begin{bmatrix} -5 & 0 \\ 0 & -5 \end{bmatrix} + \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix}$$

$$= \begin{bmatrix} -5+5 & 0+0 \\ 0+0 & -5+5 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 0$$

Hence proved

$$A^2 - 4A + 5I = 0$$

11- If $A = \begin{bmatrix} 2 & 4 \\ 1 & 5 \end{bmatrix}$, $B = \begin{bmatrix} 3 & -2 \\ 4 & 6 \end{bmatrix}$, then verify that

$$(A + B)' = A' + B'$$

Sol: $A + B = \begin{bmatrix} 2 & 4 \\ 1 & 5 \end{bmatrix} + \begin{bmatrix} 3 & -2 \\ 4 & 6 \end{bmatrix}$

$$= \begin{bmatrix} 2+3 & 4-2 \\ 1+4 & 5+6 \end{bmatrix}$$

$$= \begin{bmatrix} 5 & 2 \\ 5 & 11 \end{bmatrix}$$

$$(A + B)' = \begin{bmatrix} 5 & 5 \\ 2 & 11 \end{bmatrix} \dots\dots (i)$$

Now $A = \begin{bmatrix} 2 & 4 \\ 1 & 5 \end{bmatrix}$

$$A' = \begin{bmatrix} 2 & 1 \\ 4 & 5 \end{bmatrix} \dots\dots (ii)$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

201

and $B = \begin{bmatrix} 3 & -2 \\ 4 & 6 \end{bmatrix}$

$$B' = \begin{bmatrix} 3 & 4 \\ -2 & 6 \end{bmatrix} \dots\dots (iii)$$

$$A' + B' = \begin{bmatrix} 2 & 1 \\ 4 & 5 \end{bmatrix} + \begin{bmatrix} 3 & 4 \\ -2 & 6 \end{bmatrix} \quad \text{From (i) and (ii)}$$

$$= \begin{bmatrix} 2+3 & 1+4 \\ 4-2 & 5+6 \end{bmatrix}$$

$$= \begin{bmatrix} 5 & 5 \\ 2 & 11 \end{bmatrix} \dots\dots (iv)$$

$$(A + B)' = A' + B' \quad \text{From (i) and (iv)}$$

12- If $A = \begin{bmatrix} 1 & 2 \\ 3 & -4 \end{bmatrix}$, $B = \begin{bmatrix} 2 & -7 \\ 5 & 8 \end{bmatrix}$, $C = \begin{bmatrix} 1 & 5 \\ 0 & 2 \end{bmatrix}$ then

show that $A + B - C = \begin{bmatrix} 2 & -10 \\ 8 & 2 \end{bmatrix}$

Solution:

$$A + B - C = \begin{bmatrix} 1 & 2 \\ 3 & -4 \end{bmatrix} + \begin{bmatrix} 2 & -7 \\ 5 & 8 \end{bmatrix} - \begin{bmatrix} 1 & 5 \\ 0 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 1+2 & 2-7 \\ 3+5 & -4+8 \end{bmatrix} - \begin{bmatrix} 1 & 5 \\ 0 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 3 & -5 \\ 8 & 4 \end{bmatrix} - \begin{bmatrix} 1 & 5 \\ 0 & 2 \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

202

$$= \begin{bmatrix} 3-1 & -5-5 \\ 8-0 & 4-2 \end{bmatrix} = \begin{bmatrix} 2 & -10 \\ 8 & 2 \end{bmatrix}$$

Hence proved

$$A + B - C = \begin{bmatrix} 2 & -10 \\ 8 & 2 \end{bmatrix}$$

MULTIPLICATION OF MATRICES

Two matrices A and B are said to be conformable for the product AB, if the number of columns in A is equal to the number of rows in B.

Remember that:

For multiplication AB of two matrices A and B the following points should be kept in mind.

- (i) The number of columns in A = number of rows in B.
- (ii) The product of matrices A and B is denoted by $A \times B$ or AB.
- (iii) If A is a m -by- p matrix and B is a p -by- n matrix then AB is m -by- n matrix.

Associative Law of Matrices with respect to Multiplication

If three matrices A, B and C are conformable for multiplication, then

$$A(BC) = (AB)C$$

is called associative law with respect to multiplication.

Distributive Laws:

If the matrices A, B and C are conformable for addition and multiplication, then

- (i) $A(B + C) = AB + AC$ (left distributive law for matrices)
 - (ii) $(A + B)C = AC + BC$ (right distributive law for matrices)
- (i) and (ii) are called distributive laws.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

203

Exercise 6.4

In Problems 1 to 8 Verify Each Statement, Using

$$A = \begin{bmatrix} 4 & 2 \\ 0 & 0 \end{bmatrix}, B = \begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix}, C = \begin{bmatrix} -1 & 2 \\ 4 & 2 \end{bmatrix}$$

1. $(AB)C = A(BC)$

L.H.S = $(AB)C$

$$\begin{aligned} (AB)C &= \left(\begin{bmatrix} 4 & 2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix} \right) \times \begin{bmatrix} -1 & 2 \\ 4 & 2 \end{bmatrix} \\ &= \left(\begin{bmatrix} 8-4 & 4+8 \\ 0+0 & 0+0 \end{bmatrix} \right) \times \begin{bmatrix} -1 & 2 \\ 4 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 4 & 12 \\ 0 & 0 \end{bmatrix} \times \begin{bmatrix} -1 & 2 \\ 4 & 2 \end{bmatrix} \\ &= \begin{bmatrix} -4+48 & 8+24 \\ 0+0 & 0+0 \end{bmatrix} = \begin{bmatrix} 44 & 32 \\ 0 & 0 \end{bmatrix} \dots\dots (i) \end{aligned}$$

R.H.S = $A(BC)$

$$\begin{aligned} A(BC) &= \begin{bmatrix} 4 & 2 \\ 0 & 0 \end{bmatrix} \times \left(\begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix} \times \begin{bmatrix} -1 & 2 \\ 4 & 2 \end{bmatrix} \right) \\ &= \begin{bmatrix} 4 & 2 \\ 0 & 0 \end{bmatrix} \times \left(\begin{bmatrix} -2+4 & 4+2 \\ 2+16 & -4+8 \end{bmatrix} \right) \\ &= \begin{bmatrix} 4 & 2 \\ 0 & 0 \end{bmatrix} \times \begin{bmatrix} 2 & 6 \\ 18 & 4 \end{bmatrix} \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

204

$$= \begin{bmatrix} 8+36 & 24+8 \\ 0+0 & 0+0 \end{bmatrix} = \begin{bmatrix} 44 & 32 \\ 0 & 0 \end{bmatrix} \dots\dots (ii)$$

$(AB)C = A(BC)$ from (i) and (ii)

2. $AB \neq BA$

Sol: $AB = \begin{bmatrix} 4 & 2 \\ 0 & 0 \end{bmatrix} \times \begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix}$
 $= \begin{bmatrix} 8-4 & 4+8 \\ 0+0 & 0+0 \end{bmatrix} = \begin{bmatrix} 4 & 12 \\ 0 & 0 \end{bmatrix} \dots\dots (i)$

$$BA = \begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix} \times \begin{bmatrix} 4 & 2 \\ 0 & 0 \end{bmatrix} \quad \checkmark$$

$$= \begin{bmatrix} 8+0 & 4+0 \\ -8+0 & -4+0 \end{bmatrix} = \begin{bmatrix} 8 & 4 \\ -8 & -4 \end{bmatrix} \dots\dots (ii)$$

$AB \neq BA$ from (i) and (ii)

3. $A(B+C) = AB+AC$

Sol: L.H.S = $A(B+C)$

Putting values of A, B, C

$$\bigcirc A(B+C) = \begin{bmatrix} 4 & 2 \\ 0 & 0 \end{bmatrix} \left(\begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix} + \begin{bmatrix} -1 & 2 \\ 4 & 2 \end{bmatrix} \right)$$

$$= \begin{bmatrix} 4 & 2 \\ 0 & 0 \end{bmatrix} \left(\begin{bmatrix} 2-1 & 1+2 \\ -2+4 & 4+2 \end{bmatrix} \right)$$

$$= \begin{bmatrix} 4 & 2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 3 \\ 2 & 6 \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

205

$$= \begin{bmatrix} 4+4 & 12+12 \\ 0+0 & 0+0 \end{bmatrix} = \begin{bmatrix} 8 & 24 \\ 0 & 0 \end{bmatrix} \dots\dots (i)$$

$$\text{R.H.S} = AB + AC$$

Putting values of A, B, C

$$\begin{aligned} AB + AC &= \begin{bmatrix} 4 & 2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix} + \begin{bmatrix} 4 & 2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} -1 & 2 \\ 4 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 8-4 & 4+8 \\ 0+0 & 0+0 \end{bmatrix} + \begin{bmatrix} -4+8 & 8+4 \\ 0+0 & 0+0 \end{bmatrix} \\ &= \begin{bmatrix} 4 & 12 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 4 & 12 \\ 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 4+4 & 12+12 \\ 0+0 & 0+0 \end{bmatrix} = \begin{bmatrix} 8 & 24 \\ 0 & 0 \end{bmatrix} \dots\dots (ii) \end{aligned}$$

from (i) and (ii)

$$A(B + C) = AB + AC$$

4. $(B + C)A = BA + CA$

Sol: L.H.S = (B + C)A

$$\begin{aligned} (B+C)A &= \left(\begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix} + \begin{bmatrix} -1 & 2 \\ 4 & 2 \end{bmatrix} \right) \begin{bmatrix} 4 & 2 \\ 0 & 0 \end{bmatrix} \\ &= \left(\begin{bmatrix} 2-1 & 1+2 \\ -2+4 & 4+2 \end{bmatrix} \right) \begin{bmatrix} 4 & 2 \\ 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 3 \\ 2 & 6 \end{bmatrix} \begin{bmatrix} 4 & 2 \\ 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 4+0 & 2+0 \\ 8+0 & 4+0 \end{bmatrix} = \begin{bmatrix} 4 & 2 \\ 8 & 4 \end{bmatrix} \dots\dots (i) \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

206

$$R.H.S = BA + CA$$

$$BA + CA$$

$$\begin{aligned} BA + CA &= \begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} 4 & 2 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} -1 & 2 \\ 4 & 2 \end{bmatrix} \begin{bmatrix} 4 & 2 \\ 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 8+0 & 4+0 \\ -8+0 & -4+0 \end{bmatrix} + \begin{bmatrix} -4+0 & -2+0 \\ 16+0 & 8+0 \end{bmatrix} \\ &= \begin{bmatrix} 8 & 4 \\ -8 & -4 \end{bmatrix} + \begin{bmatrix} -4 & -2 \\ 16 & 8 \end{bmatrix} \\ &= \begin{bmatrix} 8-4 & 4-2 \\ -8+16 & -4+8 \end{bmatrix} = \begin{bmatrix} 4 & 2 \\ 8 & 4 \end{bmatrix} \dots\dots (ii) \end{aligned}$$

from (i) and (ii)

$$(B + C)A = BA + CA$$

5. $(B + C)(B - C) \neq B^2 - C^2$

Sol: L.H.S = $(B + C)(B - C)$

Putting values of B, C

$$\begin{aligned} (B + C)(B - C) &= \left(\begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix} + \begin{bmatrix} -1 & 2 \\ 4 & 2 \end{bmatrix} \right) \left(\begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix} - \begin{bmatrix} -1 & 2 \\ 4 & 2 \end{bmatrix} \right) \\ &= \begin{bmatrix} 2-1 & 1+2 \\ -2+4 & 4+2 \end{bmatrix} \begin{bmatrix} 2+1 & 1-2 \\ -2-4 & 4-2 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 3 \\ 2 & 6 \end{bmatrix} \begin{bmatrix} 3 & -1 \\ -6 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 3-18 & -1+6 \\ 6-36 & -2+12 \end{bmatrix} \\ &= \begin{bmatrix} -15 & 5 \\ -30 & 10 \end{bmatrix} \dots\dots (i) \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

207

$$\text{R.H.S } B^2 - C^2 = B \times B - C \times C$$

$$= \begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix} - \begin{bmatrix} -1 & 2 \\ 4 & 2 \end{bmatrix} \begin{bmatrix} -1 & 2 \\ 4 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 4-2 & 2+4 \\ -4-8 & -2+16 \end{bmatrix} - \begin{bmatrix} 1+8 & -2+4 \\ -4+8 & 8+4 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 6 \\ -12 & 14 \end{bmatrix} - \begin{bmatrix} 9 & 2 \\ 4 & 12 \end{bmatrix}$$

$$= \begin{bmatrix} 2-9 & 6-2 \\ -12-4 & 14-12 \end{bmatrix}$$

$$= \begin{bmatrix} -7 & 4 \\ -16 & 2 \end{bmatrix} \dots\dots\dots (ii)$$

$$(B + C)(B - C) \neq B^2 - C^2 \quad \text{from (i) and (ii)}$$

6. $(BC)^T = C^T B^T$

Sol: L.H.S = $(BC)^T$

$$BC = \begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} -1 & 2 \\ 4 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} -2+4 & 4+2 \\ 2+16 & -4+8 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 6 \\ 18 & 4 \end{bmatrix}$$

$$(BC)^T = \begin{bmatrix} 2 & 18 \\ 6 & 4 \end{bmatrix} \dots\dots\dots (i) \quad \text{and}$$

Now R.H.S = $C^T B^T$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

208

$$C = \begin{bmatrix} -1 & 2 \\ 4 & 2 \end{bmatrix}$$

$$C' = \begin{bmatrix} -1 & 4 \\ 2 & 2 \end{bmatrix}$$

and

$$B = \begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix}$$

$$B' = \begin{bmatrix} 2 & -2 \\ 1 & 4 \end{bmatrix}$$

$$\begin{aligned} \text{Now } C'B' &= \begin{bmatrix} -1 & 4 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} 2 & -2 \\ 1 & 4 \end{bmatrix} \\ &= \begin{bmatrix} -2+4 & 2+16 \\ 4+2 & -4+8 \end{bmatrix} \\ &= \begin{bmatrix} 2 & 18 \\ 6 & 4 \end{bmatrix} \dots\dots (ii) \end{aligned}$$

$$(BC)' = C'B' \quad \text{from (i) and (ii)}$$

7. $BI = B$

Sol: L.H.S = BI

$$BI = \begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$BI = \begin{bmatrix} 2+0 & 0+1 \\ -2+0 & 0+4 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix} = B$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

209

8. $BC \neq CB$

Sol: $BC = \begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} -1 & 2 \\ 4 & 2 \end{bmatrix}$
 $= \begin{bmatrix} -2+4 & 4+2 \\ 2+16 & -4+8 \end{bmatrix} = \begin{bmatrix} 2 & 6 \\ 18 & 4 \end{bmatrix} \dots\dots (i)$
 $CB = \begin{bmatrix} -1 & 2 \\ 4 & 2 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix}$
 $= \begin{bmatrix} -2-4 & -1+8 \\ 8-4 & 4+8 \end{bmatrix} = \begin{bmatrix} -6 & 7 \\ 4 & 12 \end{bmatrix} \dots\dots (ii)$

from (i) and (ii)

$BC \neq CB$

Find the Matrix Products.

9. $\begin{bmatrix} 2 & 5 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix}$

Sol: $= \begin{bmatrix} 2(1) + 5(2) & 2(-1) + (5)(3) \end{bmatrix}$
 $= \begin{bmatrix} 2+10 & -2+15 \end{bmatrix}$
 $= \begin{bmatrix} 12 & 13 \end{bmatrix}$

10. $\begin{bmatrix} 3 & 4 \\ -1 & -2 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix}$

Sol: $= \begin{bmatrix} 3(-1) + 4(2) \\ -1(-1) + (-2)(2) \end{bmatrix}$
 $= \begin{bmatrix} -3+8 \\ +1-4 \end{bmatrix} = \begin{bmatrix} 5 \\ -3 \end{bmatrix}$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

210

$$11. \begin{bmatrix} 2 & -3 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & -2 \end{bmatrix}$$

$$\begin{aligned} \text{Sol: } &= \begin{bmatrix} 2(1) + (-3)(0) & 2(-1) + (-3)(-2) \\ 1(1) + 2(0) & 1(-1) + (2)(-2) \end{bmatrix} \\ &= \begin{bmatrix} 2+0 & -2+6 \\ 1+0 & -1-4 \end{bmatrix} \\ &= \begin{bmatrix} 2 & 4 \\ 1 & -5 \end{bmatrix} \end{aligned}$$

$$12. \begin{bmatrix} -3 & 2 \\ 4 & -1 \end{bmatrix} \begin{bmatrix} -1 & 5 \\ -1 & 3 \end{bmatrix}$$

$$\begin{aligned} \text{Sol: } &= \begin{bmatrix} -3(-1) + (2)(-1) & -3(5) + (2)(3) \\ 4(-1) + (-1)(-1) & 4(5) + (-1)(3) \end{bmatrix} \\ &= \begin{bmatrix} 3-2 & -15+6 \\ -4+1 & 20-3 \end{bmatrix} \\ &= \begin{bmatrix} 1 & -9 \\ -3 & 17 \end{bmatrix} \end{aligned}$$

$$13. \begin{bmatrix} -5 & -2 \\ 1 & -3 \end{bmatrix} \begin{bmatrix} -2 & 1 \\ 0 & -3 \end{bmatrix}$$

$$\begin{aligned} \text{Sol: } &= \begin{bmatrix} (-5)(-2) + (-2)(0) & (-5)(1) + (-2)(-3) \\ 1(-2) + (-3)(0) & (1)(1) + (-3)(-3) \end{bmatrix} \\ &= \begin{bmatrix} 10+0 & -5+6 \\ -2+0 & 1+9 \end{bmatrix} \\ &= \begin{bmatrix} 10 & 1 \\ -2 & 10 \end{bmatrix} \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

211

$$14. \begin{bmatrix} -2 & 4 \\ 0 & -3 \end{bmatrix} \begin{bmatrix} -5 & -5 \\ 1 & -3 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 2 & -0 \end{bmatrix}$$

$$\begin{aligned} \text{Sol: } &= \left(\begin{bmatrix} -2 & 4 \\ 0 & -3 \end{bmatrix} \begin{bmatrix} -5 & -5 \\ 1 & -3 \end{bmatrix} \right) \begin{bmatrix} 1 & -1 \\ 2 & -0 \end{bmatrix} \\ &= \begin{bmatrix} (-2)(-5) + (4)(1) & (-2)(-5) + (4)(-3) \\ (0)(-5) + (-3)(1) & (0)(-5) + (-3)(-3) \end{bmatrix} \\ &= \begin{bmatrix} 10 + 4 & 10 - 12 \\ 0 - 3 & 0 + 9 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 2 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 14 & -2 \\ -3 & 9 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 2 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 14(1) + (-2)(2) & 14(-1) + (-2)(0) \\ -3(1) + 9(2) & -3(-1) + 9(0) \end{bmatrix} \\ &= \begin{bmatrix} 14 - 4 & -14 + 0 \\ -3 + 18 & 3 + 0 \end{bmatrix} \\ &= \begin{bmatrix} 10 & -14 \\ 15 & 3 \end{bmatrix} \end{aligned}$$

$$15. \text{ If } \begin{bmatrix} 1 & 5 \\ 3 & a \end{bmatrix} \begin{bmatrix} b \\ 7 \end{bmatrix} = \begin{bmatrix} 35 \\ 10 \end{bmatrix}, \text{ then find the values of}$$

a and b .

$$\begin{aligned} \text{Sol: } &\begin{bmatrix} 1 & 5 \\ 3 & a \end{bmatrix} \begin{bmatrix} b \\ 7 \end{bmatrix} = \begin{bmatrix} 35 \\ 10 \end{bmatrix} \\ &\begin{bmatrix} 1(b) + 5(7) \\ 3(b) + (a)(7) \end{bmatrix} = \begin{bmatrix} 35 \\ 10 \end{bmatrix} \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

212

$$\begin{bmatrix} b + 35 \\ 3b + 7a \end{bmatrix} = \begin{bmatrix} 35 \\ 10 \end{bmatrix}$$

Now $b + 35 = 35$

Hence $b = 0$

and $3b + 7a = 10$

$$3(0) + 7a = 10 \quad (\text{Putting values of } b)$$

$$7a = 10$$

$$a = \frac{10}{7}$$

16. If $A = \begin{bmatrix} 2 & 6 \\ 7 & 8 \end{bmatrix}$, $B = \begin{bmatrix} -1 & -3 \\ 2 & 0 \end{bmatrix}$, then verify

$$(AB)^T = B^T A^T$$

Sol:

$$\text{L.H.S} = (AB)^T$$

$$\begin{aligned} AB &= \begin{bmatrix} 2 & 6 \\ 7 & 8 \end{bmatrix} \begin{bmatrix} -1 & -3 \\ 2 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 2(-1) + 6(2) & 2(-3) + 6(0) \\ 7(-1) + 8(2) & 7(-3) + 8(0) \end{bmatrix} \\ &= \begin{bmatrix} -2 + 12 & -6 + 0 \\ -7 + 16 & -21 + 0 \end{bmatrix} \\ &= \begin{bmatrix} 10 & -6 \\ 9 & -21 \end{bmatrix} \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

213

$$(AB)' = \begin{bmatrix} 10 & 9 \\ -6 & -21 \end{bmatrix} \dots\dots (i)$$

$$R.I.L.S = B' A'$$

$$B = \begin{bmatrix} -1 & -3 \\ 2 & 0 \end{bmatrix}$$

$$B' = \begin{bmatrix} -1 & 2 \\ -3 & 0 \end{bmatrix}$$

and $A = \begin{bmatrix} 2 & 6 \\ 7 & 8 \end{bmatrix}$

$$A' = \begin{bmatrix} 2 & 7 \\ 6 & 8 \end{bmatrix}$$

Now $B' A' = \begin{bmatrix} -1 & 2 \\ -3 & 0 \end{bmatrix} \begin{bmatrix} 2 & 7 \\ 6 & 8 \end{bmatrix}$

$$= \begin{bmatrix} -1(2) + 2(6) & -1(7) + 2(8) \\ -3(2) + (0)(6) & -3(7) + 0(8) \end{bmatrix}$$

$$= \begin{bmatrix} -2 + 12 & -7 + 16 \\ -6 + 0 & -21 + 0 \end{bmatrix}$$

$$= \begin{bmatrix} 10 & 9 \\ -6 & -21 \end{bmatrix} \dots\dots (ii)$$

$$(AB)' = B' A' \quad \text{from (i) and (ii)}$$

MULTIPLICATIVE INVERSE OF A MATRIX

Determinant Function

If A is a square matrix, then $\det A$ or $|A|$ read "The determinant of A " is used to denote the unique real number.

Singular Matrix:

A square matrix A is called a singular matrix. If $\det A = 0$

Example : $A = \begin{bmatrix} 12 & 6 \\ 6 & 3 \end{bmatrix}$

$$\det A = \begin{vmatrix} 12 & 6 \\ 6 & 3 \end{vmatrix} = 36 - 36$$

$\det A = 0$. Hence matrix A is singular.

Non-Singular Matrix:

A square matrix A is called non-singular matrix, if $\det A \neq 0$.

Example

If $A = \begin{bmatrix} 2 & 5 \\ 6 & 8 \end{bmatrix}$

$$\det A = \begin{vmatrix} 2 & 5 \\ 6 & 8 \end{vmatrix} = 16 - 30$$

$\det A = -14$. Hence matrix A is non-singular.

Adjoint of a Matrix:

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ be a square matrix of order 2-by-2.

Then the matrix obtained by interchanging the elements of the diagonal (i.e a and d) and by changing the signs of the other elements b and c is called the adjoint of the matrix A .

Multiplicative Inverse

In the set of real numbers, we know that for each real

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

215

number a (except zero) there exists a real number a^{-1} such that $aa^{-1} = 1$.

The number a^{-1} is called the multiplicative inverse of a .

$$A^{-1} = \frac{\text{adj } A}{|A|}, |A| \neq 0$$

If A is a singular matrix then the multiplicative inverse of A does not exist.

Remember that:

- (i) Inverse of square matrix A is denoted by A^{-1} .
- (ii) Only non-singular matrices have inverses.
- (iii) Inverse of square matrix A is always unique.
- (iv) Non-square matrices cannot possess inverses.
- (v) $A^{-1} = \frac{\text{adj } A}{|A|}$

Exercise 6.5

I- Find the determinants of the following matrices.

(i) $\begin{bmatrix} u & v \\ x & y \end{bmatrix}$

(ii) $\begin{bmatrix} -2 & 5 \\ 1 & 4 \end{bmatrix}$

(iii) $\begin{bmatrix} -8 & -4 \\ -4 & -2 \end{bmatrix}$

(iv) $\begin{bmatrix} 1 & 3 \\ 1 & 8 \\ 1 & 1 \\ 8 & 4 \end{bmatrix}$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

216

(i) $A = \begin{bmatrix} u & v \\ x & y \end{bmatrix}$

Sol: Let $|A| = \begin{vmatrix} u & v \\ x & y \end{vmatrix} = uy - vx$

(ii) $P = \begin{bmatrix} -2 & 5 \\ 1 & 4 \end{bmatrix}$

Sol: Let $|P| = \begin{vmatrix} -2 & 5 \\ 1 & 4 \end{vmatrix}$
 $= (-2)(4) - (5)(1)$
 $= -8 - 5 = -13$

(iii) $B = \begin{bmatrix} -8 & -4 \\ -4 & -2 \end{bmatrix}$

Sol: Let $|B| = \begin{vmatrix} -8 & -4 \\ -4 & -2 \end{vmatrix}$
 $= (-8)(-2) - (-4)(-4)$
 $= 16 - 16 = 0$

(iv) $A = \begin{bmatrix} 1 & 3 \\ 1 & 8 \\ 1 & 1 \\ 8 & 4 \end{bmatrix}$

Sol: Let $|A| = \begin{vmatrix} 1 & 3 \\ 1 & 8 \\ 1 & 1 \\ 8 & 4 \end{vmatrix}$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

217

$$= (1) \left(\frac{1}{4} \right) - \left(\frac{3}{8} \right) \left(\frac{1}{8} \right)$$

$$= \frac{1}{4} - \frac{3}{64}$$

$$= \frac{16-3}{64} = \frac{13}{64}$$

2. Identify the singular and non-singular matrices.

(i) $\begin{bmatrix} -1 & 3 \\ 1 & -3 \end{bmatrix}$

Sol: Let $A = \begin{bmatrix} -1 & 3 \\ 1 & -3 \end{bmatrix}$

$$|A| = \begin{vmatrix} -1 & 3 \\ 1 & -3 \end{vmatrix}$$

$$= (-1)(-3) - (3)(1)$$

$$= 3 - 3 = 0$$

Therefore, $\begin{bmatrix} -1 & 3 \\ 1 & -3 \end{bmatrix}$ is singular matrix

(ii) $\begin{bmatrix} 3 & 8 \\ 4 & 9 \end{bmatrix}$

Sol: Let $B = \begin{bmatrix} 3 & 8 \\ 4 & 9 \end{bmatrix}$

$$|B| = \begin{vmatrix} 3 & 8 \\ 4 & 9 \end{vmatrix}$$

$$= (3)(9) - (8)(4)$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

218

$$= 27 - 32 = -5 \neq 0$$

Therefore, $\begin{bmatrix} 3 & 8 \\ 4 & 9 \end{bmatrix}$ is non-singular matrix.

(iii) $\begin{bmatrix} -a & b \\ a & b \end{bmatrix}$

Sol Let $P = \begin{bmatrix} -a & b \\ a & b \end{bmatrix}$

$$|P| = \begin{vmatrix} -a & b \\ a & b \end{vmatrix}$$

$$= (-a)(b) - (a)(b)$$

$$= -ab - ab$$

$$= -2ab \neq 0$$

Therefore, $\begin{bmatrix} -a & b \\ a & b \end{bmatrix}$ is non-singular matrix.

3. Find the Inverse of each matrix A and show that

$A^{-1}A = I$. If the inverse does not exist, give

reason.

(i) $A = \begin{bmatrix} 1 & 2 \\ 1 & 3 \end{bmatrix}$

Sol: $|A| = \begin{vmatrix} 1 & 2 \\ 1 & 3 \end{vmatrix}$

$$= (1)(3) - (2)(1)$$

$$= 3 - 2 = 1$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

219

$$\begin{aligned} A^{-1} &= \frac{\text{adj } A}{|A|} \\ &= \frac{\begin{bmatrix} 3 & -2 \\ -1 & 1 \end{bmatrix}}{1} \\ &= \begin{bmatrix} 3 & -2 \\ -1 & 1 \end{bmatrix} \end{aligned}$$

Now $A^{-1}A = ?$

$$\begin{aligned} A^{-1}A &= \begin{bmatrix} 3 & -2 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 3 \end{bmatrix} \\ &= \begin{bmatrix} 3(1) + (-2)(1) & 3(2) + (-2)(3) \\ -1(1) + (1)(1) & -1(2) + (1)(3) \end{bmatrix} \\ &= \begin{bmatrix} 3-2 & 6-6 \\ -1+1 & -2+3 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I \end{aligned}$$

$$A^{-1}A = I \quad (\text{proved})$$

$$(ii) \quad A = \begin{bmatrix} 2 & 1 \\ 5 & 3 \end{bmatrix}$$

$$\text{Sol: } |A| = \begin{vmatrix} 2 & 1 \\ 5 & 3 \end{vmatrix} = (2)(3) - (1)(5) = 6 - 5 = 1$$

$$A^{-1} = \frac{\text{adj } A}{|A|}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

220

$$= \begin{bmatrix} 3 & -1 \\ -5 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 3 & -1 \\ -5 & 2 \end{bmatrix}$$

$$\begin{aligned} \text{and } A^{-1}A &= \begin{bmatrix} 3 & -1 \\ -5 & 2 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 5 & 3 \end{bmatrix} \\ &= \begin{bmatrix} 3(2) + (-1)(5) & 3(1) + (-1)(3) \\ -5(2) + (2)(5) & -5(1) + (2)(3) \end{bmatrix} \\ &= \begin{bmatrix} 6-5 & 3-3 \\ -10+10 & -5+6 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I \end{aligned}$$

$$A^{-1}A = I \quad (\text{proved})$$

$$(iii) \quad A = \begin{bmatrix} 2 & 0 \\ -1 & 3 \end{bmatrix}$$

$$\text{Sol: } |A| = (2)(3) - (-1)(0)$$

$$|A| = \begin{vmatrix} 2 & 0 \\ -1 & 3 \end{vmatrix} = (2)(3) - (-1)(0)$$

$$= 6 - 0 = 6$$

$$A^{-1} = \frac{\text{adj } A}{|A|}$$

$$A^{-1} = \frac{\begin{bmatrix} 3 & 0 \\ 1 & 2 \end{bmatrix}}{6}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

221

$$= \frac{1}{6} \begin{bmatrix} 3 & 0 \\ 1 & 2 \end{bmatrix}$$

$$\begin{aligned} \text{Now } A^{-1}A &= \frac{1}{6} \begin{bmatrix} 3 & 0 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ -1 & 3 \end{bmatrix} \\ &= \frac{1}{6} \begin{bmatrix} (3)(2) + (0)(-1) & (3)(0) + (0)(3) \\ (1)(2) + (2)(-1) & (1)(0) + (2)(3) \end{bmatrix} \\ &= \frac{1}{6} \begin{bmatrix} 6+0 & 0+0 \\ 2-2 & 0+6 \end{bmatrix} \\ &= \frac{1}{6} \begin{bmatrix} 6 & 0 \\ 0 & 6 \end{bmatrix} \\ &= \begin{bmatrix} \frac{6}{6} & \frac{0}{6} \\ \frac{0}{6} & \frac{6}{6} \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I \end{aligned}$$

$$A^{-1}A = I \quad (\text{proved})$$

$$(iv) \quad A = \begin{bmatrix} -6 & 4 \\ 3 & -2 \end{bmatrix}$$

$$\begin{aligned} \text{Sol: } |A| &= \begin{vmatrix} -6 & 4 \\ 3 & -2 \end{vmatrix} \\ &= (-6)(-2) - (4)(3) \\ &= 12 - 12 = 0 \end{aligned}$$

It is a singular matrix, its inverse is not possible.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

222

$$(v) \quad A = \begin{bmatrix} 1 & 3 \\ 2 & 8 \end{bmatrix}$$

$$\text{Sol: } |A| = \begin{vmatrix} 1 & 3 \\ 2 & 8 \end{vmatrix}$$

$$= (1)(8) - (3)(2)$$

$$= 8 - 6 = 2$$

$$A^{-1} = \frac{\text{adj } A}{|A|}$$

$$= \frac{\begin{bmatrix} 8 & -3 \\ -2 & 1 \end{bmatrix}}{2}$$

$$= \frac{1}{2} \begin{bmatrix} 8 & -3 \\ -2 & 1 \end{bmatrix}$$

$$\text{Now } A^{-1}A = \frac{1}{2} \begin{bmatrix} 8 & -3 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 \\ 2 & 8 \end{bmatrix}$$

$$= \frac{1}{2} \begin{bmatrix} 8(1) + (-3)(2) & 8(3) + (-3)(8) \\ -2(1) + 1(2) & -2(3) + 1(8) \end{bmatrix}$$

$$= \frac{1}{2} \begin{bmatrix} 8 - 6 & 24 - 24 \\ -2 + 2 & -6 + 8 \end{bmatrix}$$

$$= \frac{1}{2} \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} \frac{2}{2} & \frac{0}{2} \\ \frac{0}{2} & \frac{2}{2} \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

223

$$= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

$$A^{-1}A = I \quad (\text{proved})$$

(vi) $A = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$

Sol: $|A| = \begin{vmatrix} -1 & 0 \\ 0 & -1 \end{vmatrix}$
 $= (-1)(-1) - (0)(0)$
 $= 1 - 0 = 1$

$$A^{-1} = \frac{\text{adj } A}{|A|}$$

$$= \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$$

$$= \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$$

Now $A^{-1}A = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$

$$A^{-1}A = \begin{bmatrix} (-1)(-1) + (0)(0) & (-1)(0) + (0)(-1) \\ 0(-1) + (-1)(0) & (0)(0) + (-1)(-1) \end{bmatrix}$$

$$= \begin{bmatrix} 1+0 & 0+0 \\ 0+0 & 0+1 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

$$A^{-1}A = I \quad (\text{proved})$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

224

$$(vii) \quad A = \begin{bmatrix} \frac{3}{5} & \frac{-4}{5} \\ \frac{4}{5} & \frac{3}{5} \end{bmatrix}$$

$$\begin{aligned} \text{Sol:} \quad |A| &= \begin{vmatrix} \frac{3}{5} & \frac{-4}{5} \\ \frac{4}{5} & \frac{3}{5} \end{vmatrix} \\ &= \left(\frac{3}{5}\right)\left(\frac{3}{5}\right) - \left(\frac{-4}{5}\right)\left(\frac{4}{5}\right) \\ &= \frac{9}{25} + \frac{16}{25} \\ &= \frac{9+16}{25} \\ &= \frac{25}{25} = 1 \end{aligned}$$

$$A^{-1} = \frac{\text{adj } A}{|A|}$$

$$= \frac{\begin{bmatrix} 3 & 4 \\ 5 & 5 \end{bmatrix}}{1}$$

$$= \begin{bmatrix} \frac{3}{5} & \frac{4}{5} \\ \frac{4}{5} & \frac{3}{5} \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

225

$$\begin{aligned} \text{Now } A^{-1}A &= \begin{bmatrix} \frac{3}{5} & \frac{4}{5} \\ -\frac{4}{5} & \frac{3}{5} \end{bmatrix} \begin{bmatrix} \frac{3}{5} & -\frac{4}{5} \\ \frac{4}{5} & \frac{3}{5} \end{bmatrix} \\ &= \begin{bmatrix} \left(\frac{3}{5}\right)\left(\frac{3}{5}\right) + \left(\frac{4}{5}\right)\left(\frac{4}{5}\right) & \left(\frac{3}{5}\right)\left(-\frac{4}{5}\right) + \left(\frac{4}{5}\right)\left(\frac{3}{5}\right) \\ \left(-\frac{4}{5}\right)\left(\frac{3}{5}\right) + \left(\frac{3}{5}\right)\left(\frac{4}{5}\right) & \left(-\frac{4}{5}\right)\left(-\frac{4}{5}\right) + \left(\frac{3}{5}\right)\left(\frac{3}{5}\right) \end{bmatrix} \\ &= \begin{bmatrix} \frac{9}{25} + \frac{16}{25} & -\frac{12}{25} + \frac{12}{25} \\ -\frac{12}{25} + \frac{12}{25} & \frac{16}{25} + \frac{9}{25} \end{bmatrix} \\ A^{-1}A &= \begin{bmatrix} \frac{25}{25} & \frac{0}{25} \\ \frac{0}{25} & \frac{25}{25} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I \end{aligned}$$

$$A^{-1}A = I \quad \text{Hence proved}$$

4. Let $M = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$

(a) Find M^{-1}

(b) Verify that $M^{-1}M = MM^{-1}$

Sol: $M = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$

$$|M| = \begin{vmatrix} 1 & 2 \\ 3 & 4 \end{vmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

226

Pilot Super One "General Math" 10th

$$= (1 \times 4) - (2)(3)$$

$$= 4 - 6 = -2$$

$$M^{-1} = \frac{\text{adj } M}{|M|}$$

$$= \frac{\begin{bmatrix} 4 & -2 \\ -3 & 1 \end{bmatrix}}{-2}$$

$$= \frac{1}{-2} \begin{bmatrix} 4 & -2 \\ -3 & 1 \end{bmatrix} = \begin{bmatrix} 4\left(-\frac{1}{2}\right) & (-2)\left(-\frac{1}{2}\right) \\ -3\left(-\frac{1}{2}\right) & (1)\left(-\frac{1}{2}\right) \end{bmatrix}$$

$$= \begin{bmatrix} -2 & 1 \\ \frac{3}{2} & -\frac{1}{2} \end{bmatrix}$$

$$M^{-1}M = \begin{bmatrix} -2 & 1 \\ \frac{3}{2} & -\frac{1}{2} \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$

$$= \begin{bmatrix} (-2)(1) + (1)(3) & (-2)(2) + (1)(4) \\ \left(\frac{3}{2}\right)(1) + \left(-\frac{1}{2}\right)(3) & \left(\frac{3}{2}\right)(2) + \left(-\frac{1}{2}\right)(4) \end{bmatrix}$$

$$= \begin{bmatrix} -2 + 3 & -4 + 4 \\ \frac{3}{2} - \frac{3}{2} & 3 - 2 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \dots\dots\dots(i)$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

227

$$\begin{aligned} \text{and now } MM^{-1} &= \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} -2 & 1 \\ \frac{3}{2} & -\frac{1}{2} \end{bmatrix} \\ &= \begin{bmatrix} (1)(-2) + (2)\left(\frac{3}{2}\right) & (1)(1) + (2)\left(-\frac{1}{2}\right) \\ (3)(-2) + (4)\left(\frac{3}{2}\right) & (3)(1) + (4)\left(-\frac{1}{2}\right) \end{bmatrix} \\ &= \begin{bmatrix} -2 + 3 & 1 - 1 \\ -6 + 6 & 3 - 2 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \dots\dots (ii) \end{aligned}$$

$$M^{-1}M = MM^{-1} \quad \text{from (i) and (ii)}$$

5. If $A = \begin{bmatrix} 5 & 2 \\ 2 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 4 & 2 \\ 3 & -1 \end{bmatrix}$, verify that
 $(AB)^{-1} = B^{-1}A^{-1}$.

Sol: L.H.S = $(AB)^{-1}$

$$\begin{aligned} AB &= \begin{bmatrix} 5 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 2 \\ 3 & -1 \end{bmatrix} \\ &= \begin{bmatrix} (5)(4) + (2)(3) & (5)(2) + (2)(-1) \\ (2)(4) + (1)(3) & (2)(2) + (1)(-1) \end{bmatrix} \\ &= \begin{bmatrix} 20 + 6 & 10 - 2 \\ 8 + 3 & 4 - 1 \end{bmatrix} \\ &= \begin{bmatrix} 26 & 8 \\ 11 & 3 \end{bmatrix} \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

228

$$\text{Now } |AB| = \begin{vmatrix} 26 & 8 \\ 11 & 3 \end{vmatrix}$$

$$= (26)(3) - (11)(8)$$

$$= 78 - 88$$

$$= -10$$

$$(AB)^{-1} = \frac{\text{adj}(AB)}{|AB|}$$

$$= \frac{\begin{bmatrix} 3 & -8 \\ -11 & 26 \end{bmatrix}}{-10} = -\frac{1}{10} \begin{bmatrix} 3 & -8 \\ -11 & 26 \end{bmatrix}$$

$$= \begin{bmatrix} (3)\left(-\frac{1}{10}\right) & (-8)\left(-\frac{1}{10}\right) \\ (-11)\left(-\frac{1}{10}\right) & (26)\left(-\frac{1}{10}\right) \end{bmatrix}$$

$$= \begin{bmatrix} -\frac{3}{10} & \frac{4}{5} \\ \frac{11}{10} & -\frac{13}{5} \end{bmatrix}$$

$$\text{R.H.S} = B^{-1}A^{-1}$$

$$B^{-1}A^{-1} = ?$$

$$B = \begin{bmatrix} 4 & 2 \\ 3 & -1 \end{bmatrix}$$

$$|B| = \begin{vmatrix} 4 & 2 \\ 3 & -1 \end{vmatrix}$$

$$= (4)(-1) - (2)(3)$$

$$= -4 - 6$$

$$= -10$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

229

$$B^{-1} = \frac{\text{adj } B}{|B|}$$

$$B^{-1} = \frac{\begin{bmatrix} -1 & -2 \\ -3 & 4 \end{bmatrix}}{-10}$$

$$= -\frac{1}{10} \begin{bmatrix} -1 & -2 \\ -3 & 4 \end{bmatrix}$$

$$= \begin{bmatrix} (-1)\left(-\frac{1}{10}\right) & (-2)\left(-\frac{1}{10}\right) \\ (-3)\left(-\frac{1}{10}\right) & (4)\left(-\frac{1}{10}\right) \end{bmatrix}$$

$$= \begin{bmatrix} \frac{1}{10} & \frac{2}{10} \\ \frac{3}{10} & -\frac{4}{10} \end{bmatrix}$$

$$\text{and } A = \begin{bmatrix} 5 & 2 \\ 2 & 1 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 5 & 2 \\ 2 & 1 \end{vmatrix}$$

$$= (5)(1) - (2)(2)$$

$$= 5 - 4$$

$$= 1$$

$$A^{-1} = \frac{\text{adj } A}{|A|}$$

$$= \frac{\begin{bmatrix} 1 & -2 \\ -2 & 5 \end{bmatrix}}{1} = \begin{bmatrix} 1 & -2 \\ -2 & 5 \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

230

$$\begin{aligned}
 \text{Now } B^{-1}A^{-1} &= \begin{bmatrix} \frac{1}{10} & \frac{1}{5} \\ \frac{3}{10} & -\frac{2}{5} \end{bmatrix} \begin{bmatrix} 1 & -2 \\ -2 & 5 \end{bmatrix} \\
 &= \begin{bmatrix} \left(\frac{1}{10}\right)(1) + \left(\frac{1}{5}\right)(-2) & \left(\frac{1}{10}\right)(-2) + \left(\frac{1}{5}\right)(5) \\ \left(\frac{3}{10}\right)(1) + \left(-\frac{2}{5}\right)(-2) & \left(\frac{3}{10}\right)(-2) + \left(-\frac{2}{5}\right)(5) \end{bmatrix} \\
 &= \begin{bmatrix} \frac{1}{10} - \frac{2}{5} & -\frac{1}{5} + 1 \\ \frac{3}{10} + \frac{4}{5} & -\frac{3}{5} - 2 \end{bmatrix} \\
 &= \begin{bmatrix} \frac{1-4}{10} & \frac{-1+5}{5} \\ \frac{3+8}{10} & \frac{-3-10}{5} \end{bmatrix} \\
 &= \begin{bmatrix} -\frac{3}{10} & \frac{4}{5} \\ \frac{11}{10} & -\frac{13}{5} \end{bmatrix} \dots\dots (ii)
 \end{aligned}$$

from (i) and (ii)

$$(AB)^{-1} = B^{-1}A^{-1}$$

SOLUTION OF SIMULTANEOUS LINEAR EQUATIONS

- ★ To determine the value of two variables, we need a pair of equations. Such a pair of equations is called a system of simultaneous linear equations.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

231



- 1- Write the equation $2x + ky = 7$ and $4x - 9y = 4$ in matrix form. Also find the value of k if the matrix of the coefficients is singular.

Sol: $2x + ky = 7$
 $4x - 9y = 4$

In matrix form

$$\begin{bmatrix} 2 & k \\ 4 & -9 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 7 \\ 4 \end{bmatrix}$$

$$A \cdot X = B$$

$$A = \begin{bmatrix} 2 & k \\ 4 & -9 \end{bmatrix}, \quad X = \begin{bmatrix} x \\ y \end{bmatrix}, \quad B = \begin{bmatrix} 7 \\ 4 \end{bmatrix}$$

$$\Delta = 0$$

Now, if A is singular matrix.
 Therefore,

$$\begin{bmatrix} 2 & k \\ 4 & -9 \end{bmatrix} = 0$$

$$(2)(-9) - (k)(4) = 0$$

$$-18 - 4k = 0$$

$$-4k = 18$$

$$k = \frac{18}{-4}$$

$$= -\frac{9}{2}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

232

2. Solve the simultaneous equations by the matrix inversion method where possible. Where there is no solution, explain why this is so.

$$\begin{aligned} (i) \quad & 2x - 5y = 1 \\ & 3x - 7y = 2 \end{aligned}$$

Sol. In matrix form

$$\begin{bmatrix} 2 & -5 \\ 3 & -7 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

$$A \cdot X = B$$

$$A = \begin{bmatrix} 2 & -5 \\ 3 & -7 \end{bmatrix}, \quad X = \begin{bmatrix} x \\ y \end{bmatrix}, \quad B = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

$$\text{and } A^{-1} = \begin{bmatrix} 2 & -5 \\ 3 & -7 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 2 & -5 \\ 3 & -7 \end{vmatrix}$$

$$= (2)(-7) - (-5)(3)$$

$$= -14 + 15 = 1 \neq 0$$

A is non-singular matrix. Therefore, equations can be

solved

$$A^{-1} = \frac{\text{adj } A}{|A|}$$

$$= \frac{\begin{bmatrix} -7 & 5 \\ -3 & 2 \end{bmatrix}}{1}$$

$$= \begin{bmatrix} -7 & 5 \\ -3 & 2 \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

233

because $X = A^{-1}B$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -7 & 5 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -7(1) + (5)(2) \\ -3(1) + (2)(2) \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -7 + 10 \\ -3 + 4 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

Hence, $x = 3$
 $y = 1$

$$S.S = \{(3, 1)\}$$

(ii) $3x + 2y = 10$
 $2y - 3x = -4$

Sol: Write equations again

$$3x + 2y = 10$$

$$-3x + 2y = -4$$

In matrix form

$$\begin{bmatrix} 3 & 2 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 10 \\ -4 \end{bmatrix}$$

$$AX = B$$

$$A = \begin{bmatrix} 3 & 2 \\ -3 & 2 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix}, B = \begin{bmatrix} 10 \\ -4 \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

234

$$X = A^{-1}B$$

Now $A = \begin{bmatrix} 3 & 2 \\ -3 & 2 \end{bmatrix}$

$$|A| = \begin{vmatrix} 3 & 2 \\ -3 & 2 \end{vmatrix}$$

$$= (3 \times 2) - (2 \times -3)$$

$$= 6 + 6 = 12 \neq 0$$

A is non-singular matrix, therefore equations can be solve

$$A^{-1} = \frac{\text{adj } A}{|A|}$$

$$= \frac{\begin{bmatrix} 2 & -2 \\ 3 & 3 \end{bmatrix}}{12}$$

$$= \frac{1}{12} \begin{bmatrix} 2 & -2 \\ 3 & 3 \end{bmatrix}$$

Now $\begin{bmatrix} X \\ Y \end{bmatrix} = \frac{1}{12} \begin{bmatrix} 2 & -2 \\ 3 & 3 \end{bmatrix} \begin{bmatrix} 10 \\ -4 \end{bmatrix}$

$$\begin{bmatrix} X \\ Y \end{bmatrix} = \frac{1}{12} \begin{bmatrix} 2(10) + (-2)(-4) \\ 3(10) + (3)(-4) \end{bmatrix}$$

$$\begin{bmatrix} X \\ Y \end{bmatrix} = \frac{1}{12} \begin{bmatrix} 20 + 8 \\ 30 - 12 \end{bmatrix}$$

$$\begin{bmatrix} X \\ Y \end{bmatrix} = \frac{1}{12} \begin{bmatrix} 28 \\ 18 \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

235

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{28}{12} \\ \frac{18}{12} \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{7}{3} \\ \frac{3}{2} \end{bmatrix}$$

Hence, $x = \frac{7}{3}$

$$y = \frac{3}{2}$$

$$\text{S.S} = \left\{ \left(\frac{7}{3}, \frac{3}{2} \right) \right\}$$

(iii) $4x + 5y = 0$

$$2x + 5y = 1$$

Sol: $\begin{bmatrix} 4 & 5 \\ 2 & 5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$

In matrix form

Let $AX = B$

Now $A^{-1}A X = A^{-1}B$

$$X = A^{-1}B$$

Now $A = \begin{bmatrix} 4 & 5 \\ 2 & 5 \end{bmatrix}$

$$|A| = \begin{vmatrix} 4 & 5 \\ 2 & 5 \end{vmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

236

$$|A| = (4 \times 5) - (5 \times 2)$$

$$= 20 - 10 = 10 \neq 0$$

A is non-singular matrix, therefore equations can be solve.

$$|A| = 10$$

$$A^{-1} = \frac{\text{adj } A}{|A|}$$

$$= \frac{\begin{bmatrix} 5 & -5 \\ -2 & 4 \end{bmatrix}}{10} = \frac{1}{10} \begin{bmatrix} 5 & -5 \\ -2 & 4 \end{bmatrix}$$

$$\text{Now } A^{-1}B = \frac{1}{10} \begin{bmatrix} 5 & -5 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$= \frac{1}{10} \begin{bmatrix} (5 \times 0) + (-5 \times 1) \\ (-2 \times 0) + (4 \times 1) \end{bmatrix}$$

$$= \frac{1}{10} \begin{bmatrix} 0 - 5 \\ 0 + 4 \end{bmatrix}$$

$$= \frac{1}{10} \begin{bmatrix} -5 \\ 4 \end{bmatrix}$$

$$= \begin{bmatrix} \frac{-5}{10} \\ \frac{4}{10} \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

237

$$A^{-1}B = \begin{bmatrix} 1 \\ -\frac{1}{2} \\ \frac{2}{5} \end{bmatrix}$$

But $X = A^{-1}B$

Thus $X = \begin{bmatrix} 1 \\ -\frac{1}{2} \\ \frac{2}{5} \end{bmatrix}$

or $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ -\frac{1}{2} \\ \frac{2}{5} \end{bmatrix}$

Hence, $x = -\frac{1}{2}$

$$y = \frac{2}{5}$$

$$S.S = \left\{ \left(-\frac{1}{2}, \frac{2}{5} \right) \right\}$$

(iv) $5x + 6y = 25$

$$3x + 4y = 17$$

Sol: $\begin{bmatrix} 5 & 6 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 25 \\ 17 \end{bmatrix}$

In matrix form

Let $A \quad X = B$

$$A^{-1}AX = A^{-1}B$$

$$X = A^{-1}B$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

238

Now $A = \begin{bmatrix} 5 & 6 \\ 3 & 4 \end{bmatrix}$

$$|A| = \begin{vmatrix} 5 & 6 \\ 3 & 4 \end{vmatrix}$$

$$= (5)(4) - (6)(3)$$

$$= 20 - 18$$

$$|A| = 2 \neq 0$$

A is non-singular matrix. Therefore equations can be solve.

$$A^{-1} = \frac{\text{adj } A}{|A|}$$

$$= \frac{\begin{bmatrix} 4 & -6 \\ -3 & 5 \end{bmatrix}}{2}$$

$$= \frac{1}{2} \begin{bmatrix} 4 & -6 \\ -3 & 5 \end{bmatrix}$$

Now $A^{-1}B = \frac{1}{2} \begin{bmatrix} 4 & -6 \\ -3 & 5 \end{bmatrix} \begin{bmatrix} 25 \\ 17 \end{bmatrix}$

$$= \frac{1}{2} \begin{bmatrix} (4)(25) + (-6)(17) \\ (-3)(25) + (5)(17) \end{bmatrix}$$

$$= \frac{1}{2} \begin{bmatrix} 100 - 102 \\ -75 + 85 \end{bmatrix}$$

$$= \frac{1}{2} \begin{bmatrix} -2 \\ 10 \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

239

$$= \begin{bmatrix} \frac{-2}{2} \\ \frac{10}{2} \end{bmatrix}$$

$$A^{-1}B = \begin{bmatrix} -1 \\ 5 \end{bmatrix}$$

But $X = A^{-1}B$

Thus $X = \begin{bmatrix} -1 \\ 5 \end{bmatrix}$

or $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -1 \\ 5 \end{bmatrix}$

Hence, $x = -1$

$y = 5$

S.S = $\{(-1, 5)\}$

(v) $x + y = 2$

$y = 2 + x$

Sol: Write equations again

$x + y = 2$

$-x + y = 2$

In matrix form

$$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$$

Let $A X = B$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

240

$$A^{-1}AX = A^{-1}B$$

$$X = A^{-1}B$$

Now $A = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$

$$|A| = \begin{vmatrix} 1 & 1 \\ -1 & 1 \end{vmatrix}$$

$$= (1)(1) - (1)(-1)$$

$$= 1 + 1$$

$$|A| = 2 \neq 0$$

A is non-singular matrix. Therefore equations can be solve.

$$A^{-1} = \frac{\text{adj } A}{|A|}$$

$$= \frac{\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}}{2} = \frac{1}{2} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$$

and $A^{-1}B = \frac{1}{2} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 2 \end{bmatrix}$

Now

$$= \frac{1}{2} \begin{bmatrix} (1)(2) + (-1)(2) \\ (1)(2) + (1)(2) \end{bmatrix}$$

$$= \frac{1}{2} \begin{bmatrix} 2 - 2 \\ 2 + 2 \end{bmatrix}$$

$$= \frac{1}{2} \begin{bmatrix} 0 \\ 4 \end{bmatrix}$$

$$= \begin{bmatrix} 0 \\ 2 \end{bmatrix}$$

$$A^{-1}B = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$$

while $X = A^{-1}B$

$$X = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$$

or $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$

Hence, $x = 0$

$$y = 2$$

$$\text{s.s} = \{(0, 2)\}$$

(vi) $\frac{x}{2} + \frac{y}{3} = 1$ (i)

$$-4x + y = 14 \text{ (ii)}$$

Sol: Multiply (i) by 6.

$$6\left(\frac{x}{2}\right) + 6\left(\frac{y}{3}\right) = 6$$

$$3x + 2y = 6$$

(2nd condition)

$$-4x + y = 14$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

242

$$\begin{bmatrix} 3 & 2 \\ -4 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 6 \\ 14 \end{bmatrix} \quad (\text{In matrix form})$$

Let $A X = B$

$$A^{-1} A X = A^{-1} B$$

$$X = A^{-1} B$$

Now $A = \begin{bmatrix} 3 & 2 \\ -4 & 1 \end{bmatrix}$

$$|A| = \begin{vmatrix} 3 & 2 \\ -4 & 1 \end{vmatrix}$$

$$= (3)(1) - (2)(-4)$$

$$= 3 + 8$$

$$= 11 \neq 0$$

A is non-singular matrix. Therefore equations can be solve.

$$A^{-1} = \frac{\text{adj } A}{|A|}$$

$$Q = \frac{\begin{bmatrix} 1 & -2 \\ 4 & 3 \end{bmatrix}}{11} = \frac{1}{11} \begin{bmatrix} 1 & -2 \\ 4 & 3 \end{bmatrix}$$

and now $A^{-1} B = \frac{1}{11} \begin{bmatrix} 1 & -2 \\ 4 & 3 \end{bmatrix} \begin{bmatrix} 6 \\ 14 \end{bmatrix}$

$$= \frac{1}{11} \begin{bmatrix} (1)(6) + (-2)(14) \\ (4)(6) + (3)(14) \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

243

$$= \frac{1}{11} \begin{bmatrix} 6 - 28 \\ 24 + 42 \end{bmatrix}$$

$$= \frac{1}{11} \begin{bmatrix} -22 \\ 66 \end{bmatrix}$$

$$= \begin{bmatrix} \frac{-22}{11} \\ \frac{66}{11} \end{bmatrix}$$

$$A^{-1}B = \begin{bmatrix} -2 \\ 6 \end{bmatrix}$$

while $X = A^{-1}B$

$$X = \begin{bmatrix} -2 \\ 6 \end{bmatrix}$$

or $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -2 \\ 6 \end{bmatrix}$

Hence, $x = -2$

$$y = 6$$

$$S = \{(-2, 6)\}$$

Solve, using matrix inversion method

$$3x + y = 10$$

$$2x + 3y = 3$$

In matrix form

$$\begin{bmatrix} 3 & 1 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 10 \\ 3 \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

244

$$\text{Let } AX = B$$

$$\text{and } A^{-1}AX = A^{-1}B$$

$$X = A^{-1}B$$

$$\text{Now } A = \begin{bmatrix} 3 & -1 \\ 2 & 3 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 3 & -1 \\ 2 & 3 \end{vmatrix}$$

$$= (3)(3) - (-1)(2)$$

$$= 9 + 2$$

$$= 11 \neq 0$$

A is non-singular matrix. Therefore equations can be solve.

$$A^{-1} = \frac{\text{adj } A}{|A|}$$

$$= \frac{\begin{bmatrix} 3 & 1 \\ -2 & 3 \end{bmatrix}}{11} = \frac{1}{11} \begin{bmatrix} 3 & 1 \\ -2 & 3 \end{bmatrix}$$

and
now

$$A^{-1}B = \frac{1}{11} \begin{bmatrix} 3 & 1 \\ -2 & 3 \end{bmatrix} \begin{bmatrix} 10 \\ 3 \end{bmatrix}$$

$$= \frac{1}{11} \begin{bmatrix} (3)(10) + (1)(3) \\ (-2)(10) + (3)(3) \end{bmatrix}$$

$$= \frac{1}{11} \begin{bmatrix} 30 + 3 \\ -20 + 9 \end{bmatrix}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

245

$$= \frac{1}{11} \begin{bmatrix} 33 \\ -11 \end{bmatrix}$$

$$= \begin{bmatrix} \frac{33}{11} \\ \frac{-11}{11} \end{bmatrix}$$

$$A^{-1}B = \begin{bmatrix} 3 \\ -1 \end{bmatrix}$$

while $X = A^{-1}B$

$$X = \begin{bmatrix} 3 \\ -1 \end{bmatrix}$$

or $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3 \\ -1 \end{bmatrix}$

Hence, $x = 3$

$y = -1$

S.S = $\{(3, -1)\}$

4. Use Cramer's rule to solve the simultaneous equations. Give the reason where solution is not possible.

(i) $x + 2y = 3$

$x + 3y = 5$

Remember that:

Cramer's Rule

If $a_1x + a_2y = b_1$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

246

$$a_3x + a_4y = b_2$$

then

$$x = \frac{\begin{vmatrix} b_1 & a_2 \\ b_2 & a_4 \end{vmatrix}}{\begin{vmatrix} a_1 & a_2 \\ a_3 & a_4 \end{vmatrix}} = \frac{|D_1|}{|A|}$$

$$y = \frac{\begin{vmatrix} a_1 & b_1 \\ a_3 & b_2 \end{vmatrix}}{\begin{vmatrix} a_1 & a_2 \\ a_3 & a_4 \end{vmatrix}} = \frac{|D_2|}{|A|}$$

Sol: $x + 2y = 3$

$$x + 3y = 5$$

$$\begin{bmatrix} 1 & 2 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3 \\ 5 \end{bmatrix} \quad \text{In matrix form}$$

$$AX = B$$

$$A = \begin{bmatrix} 1 & 2 \\ 1 & 3 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 1 & 2 \\ 1 & 3 \end{vmatrix} = (1)(3) - (1)(2) = 3 - 2 = 1$$

$$|D_1| = \begin{vmatrix} 3 & 2 \\ 5 & 3 \end{vmatrix} = (3)(3) - (2)(5) = 9 - 10 = -1$$

$$|D_2| = \begin{vmatrix} 1 & 3 \\ 1 & 5 \end{vmatrix} = (1)(5) - (3)(1) = 5 - 3 = 2$$

$$x = \frac{|D_1|}{|A|} = \frac{-1}{1} = -1$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

247

$$y = \frac{|D_2|}{|A|} = \frac{2}{1} = 2$$

$$S.S = \{(-1, 2)\}$$

(ii) $2x + y = 1$
 $5x + 3y = 2$

Sol: $\begin{bmatrix} 2 & 1 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$
 $AX = E$

In matrix form

$$A = \begin{bmatrix} 2 & 1 \\ 5 & 3 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 2 & 1 \\ 5 & 3 \end{vmatrix}$$

$$= (2)(3) - (1)(5)$$

$$= 6 - 5 = 1$$

$$|D_1| = \begin{vmatrix} 1 & 1 \\ 2 & 3 \end{vmatrix}$$

$$= (1)(3) - (1)(2)$$

$$= 3 - 2 = 1$$

$$|D_2| = \begin{vmatrix} 2 & 1 \\ 5 & 2 \end{vmatrix} = (2)(2) - (1)(5)$$

$$= 4 - 5$$

$$= -1$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

=====

Pilot Super One "General Math" 10th

248

$$x = \frac{|D_1|}{|A|} = \frac{1}{1} = 1$$

$$y = \frac{|D_2|}{|A|} = \frac{-1}{1} = -1$$

$$\text{S.S} = \{(1, -1)\}$$

(iii) $x + 3y = 1$

$$2x + 8y = 0$$

Sol: $\begin{bmatrix} 1 & 3 \\ 2 & 8 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$

In matrix form

$$AX = B$$

$$A = \begin{bmatrix} 1 & 3 \\ 2 & 8 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 1 & 3 \\ 2 & 8 \end{vmatrix} = (1)(8) - (3)(2)$$

$$= 8 - 6 = 2$$

$$|D_1| = \begin{vmatrix} 1 & 3 \\ 0 & 8 \end{vmatrix} = (1)(8) - (3)(0)$$

$$= 8 - 0$$

$$= 8$$

$$|D_2| = \begin{vmatrix} 1 & 1 \\ 2 & 0 \end{vmatrix} = (1)(0) - (1)(2)$$

$$= 0 - 2$$

$$= -2$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

249

$$x = \frac{|D_1|}{|A|} = \frac{8}{2} = 4$$

$$y = \frac{|D_2|}{|A|} = \frac{-2}{2} = -1$$

$$S.S = \{(4, -1)\}$$

(iv) $-2x + 6y = 5$

$$x - 3y = -7$$

Sol: $\begin{bmatrix} -2 & 6 \\ 1 & -3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 5 \\ -7 \end{bmatrix}$ In matrix form
 $AX = B$

$$A = \begin{bmatrix} -2 & 6 \\ 1 & -3 \end{bmatrix}$$

$$|A| = \begin{vmatrix} -2 & 6 \\ 1 & -3 \end{vmatrix}$$

$$= (-2)(-3) - (6)(1)$$

$$= 6 - 6 = 0$$

$$A = \begin{bmatrix} -2 & 6 \\ 1 & -3 \end{bmatrix} \text{ is singular matrix, therefore S.S}$$

is not possible.

(v) $x - 3y = 5$

$$2x - 5y = 9$$

Sol: $\begin{bmatrix} 1 & -3 \\ 2 & -5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 5 \\ 9 \end{bmatrix}$ In matrix form

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

250

$$AX = B$$

$$A = \begin{bmatrix} 1 & -3 \\ 2 & -5 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 1 & -3 \\ 2 & -5 \end{vmatrix}$$

$$= (1)(-5) - (-3)(2)$$

$$= -5 + 6 = 1$$

$$|D_1| = \begin{vmatrix} 5 & -3 \\ 9 & -5 \end{vmatrix} = (5)(-5) - (-3)(9)$$

$$= -25 + 27$$

$$= 2$$

$$|D_2| = \begin{vmatrix} 1 & 5 \\ 2 & 9 \end{vmatrix}$$

$$= (1)(9) - (5)(2)$$

$$= 9 - 10 = -1$$

$$x = \frac{|D_1|}{|A|} = \frac{2}{1} = 2$$

$$y = \frac{|D_2|}{|A|} = \frac{-1}{1} = -1$$

$$S.S = \{(2, -1)\}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

Pilot Super One "General Math" 10th

251

(vi) $5x + 2y = 13$

$$2x + 5y = 17$$

Sol: $\begin{bmatrix} 5 & 2 \\ 2 & 5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 13 \\ 17 \end{bmatrix}$ In matrix form

$$AX = B$$

$$A = \begin{bmatrix} 5 & 2 \\ 2 & 5 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 5 & 2 \\ 2 & 5 \end{vmatrix} = (5)(5) - (2)(2)$$

$$= 25 - 4$$

$$= 21$$

$$|D_1| = \begin{vmatrix} 13 & 2 \\ 17 & 5 \end{vmatrix} = (13)(5) - (2)(17)$$

$$= 65 - 34$$

$$= 31$$

$$|D_2| = \begin{vmatrix} 5 & 13 \\ 2 & 17 \end{vmatrix} = (5)(17) - (13)(2)$$

$$= 85 - 26 = 59$$

$$x = \frac{|D_1|}{|A|} = \frac{31}{21}$$

$$y = \frac{|D_2|}{|A|} = \frac{59}{21}$$

$$\text{S.S} = \left\{ \left(\frac{31}{21}, \frac{59}{21} \right) \right\}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 6)

=====

Pilot Super One "General Math" 10th

252

5. Write the following matrices in the form of linear equations.

(i)
$$\begin{bmatrix} 2 & -1 \\ 5 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$$

Sol:
$$\begin{bmatrix} (2)(x) + (-1)(y) \\ (5)(x) + (2)(y) \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$$

then
$$\begin{bmatrix} 2x - y \\ 5x + 2y \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$$

and $2x - y = 2$

$$5x + 2y = 4$$

(ii)
$$\begin{bmatrix} -5 & 2 \\ 2 & -3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$$

Sol:
$$\begin{bmatrix} (-5)(x) + (2)(y) \\ (2)(x) + (-3)(y) \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$$

$$\begin{bmatrix} -5x + 2y \\ 2x - 3y \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$$

thus, $-5x + 2y = 2$

and $2x - 3y = -1$

(iii)
$$\begin{bmatrix} -4 & 1 \\ 5 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

Sol:
$$\begin{bmatrix} -4(x) + (1)(y) \\ 5(x) + (4)(y) \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$\begin{bmatrix} -4x + y \\ 5x + 4y \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

thus, $-4x + y = 1$

and $5x + 4y = -1$

(iv)
$$\begin{bmatrix} 0.8 & -0.6 \\ 0.6 & 0.8 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

Sol:
$$\begin{bmatrix} (0.8)(x) + (-0.6)(y) \\ (0.6)(x) + (0.8)(y) \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$
$$\begin{bmatrix} 0.8x - 0.6y \\ 0.6x + 0.8y \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

thus, $0.8x - 0.6y = 1$

and $0.6x + 0.8y = 2$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

262

مذہب (17) 10

1. Write down the angles marked with letters. Write whether the angles are complimentary or supplementary?

Sol:



(i)

$$\begin{aligned} \text{(i)} \quad m\angle a + 50^\circ &= 180^\circ \\ m\angle a &= 180^\circ - 50^\circ \\ m\angle a &= 130^\circ \end{aligned}$$

(Supplementary angles)



$$\begin{aligned} \text{(ii)} \quad m\angle b + 65^\circ &= 180^\circ \\ m\angle b &= 180^\circ - 65^\circ \\ m\angle b &= 115^\circ \end{aligned}$$


(Supplementary angles)



(iii)

$$\begin{aligned} \text{(iii)} \quad m\angle c + 138^\circ &= 180^\circ \\ m\angle c &= 180^\circ - 138^\circ \\ m\angle c &= 42^\circ \end{aligned}$$

(Supplementary angles)




$$\begin{aligned} \text{(iv)} \quad m\angle b + 60^\circ &= 90^\circ \\ m\angle b &= 90^\circ - 60^\circ \\ m\angle b &= 30^\circ \end{aligned}$$

(Complementary angles)

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

263




(v) $m\angle e + 72^\circ = 180^\circ$

$$m\angle e = 180^\circ - 72^\circ$$

$$m\angle e = 108^\circ$$

(Supplementary angles)



(vi) $m\angle b + 70^\circ = 90^\circ$

$$m\angle b = 90^\circ - 70^\circ$$

$$m\angle b = 20^\circ$$

(Complementary angles)

2. Two angles are supplementary and the greater exceeds the smaller by 30° . How many degrees are there in each angle?

Sol: The size of small angle = x

Let

The size of large angle = $x + 30^\circ$

According to statement

$$x + (x + 30^\circ) = 180^\circ$$

$$x + x + 30^\circ = 180^\circ$$

$$2x + 30^\circ = 180^\circ$$

$$2x = 180^\circ - 30^\circ$$

$$2x = 150^\circ$$

$$x = \frac{150^\circ}{2}$$

$$x = 75^\circ$$

The size of small angle = 75°

The size of large angle = $x + 30^\circ = 75^\circ + 30^\circ = 105^\circ$

angles = $75^\circ, 105^\circ$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

264

3. If 40° is added to an angle, the resulting angle is equal to the supplement of the original angle. Find the original angle.

Sol: Let the required angle = x

size of the angle after adding = $x + 40^\circ$

The supplement of 1st angle = $180^\circ - x$

According to statement

$$x + 40^\circ = 180^\circ - x$$

$$x + x = 180^\circ - 40^\circ$$

$$2x = 140^\circ$$

$$x = \frac{140^\circ}{2}$$

$$x = 70^\circ$$

4. The sum of two angles is 100° , and the difference between their supplements is 100° . Find the angles.

Let the size of 1st angle = x°

The size of 2nd angle = $100^\circ - x^\circ$

The supplement of $x^\circ = 180^\circ - x^\circ$

The supplement of $100^\circ - x^\circ = 180^\circ - (100^\circ - x^\circ)$

$$= 180^\circ - 100^\circ + x^\circ$$

According to the statement

$$(180^\circ - 100^\circ + x^\circ) - (180^\circ - x^\circ) = 100^\circ$$

$$180^\circ - 100^\circ + x^\circ - 180^\circ + x^\circ = 100^\circ$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

265

$$2x^{\circ} - 100^{\circ} = 100^{\circ}$$

$$2x = 100^{\circ} + 100^{\circ}$$

$$2x = 200^{\circ}$$

$$x = \frac{200^{\circ}}{2}$$

$$x = 100^{\circ}$$

The size of 1st angle = 100°

$$\begin{aligned}\text{The size of 2nd angle} &= 100^{\circ} - 100^{\circ} \\ &= 0^{\circ}\end{aligned}$$

5. The sum of two angles is 100° , the supplement of the first angle exceeds the supplement of the second angle 40° . Find the angles.

Sol: Let the size of 1st angle = x°

The size of 2nd angle = $100^{\circ} - x^{\circ}$

The supplement of 1st angle = $180^{\circ} - x^{\circ}$

$$\begin{aligned}\text{The supplement of 2nd angle} &= 180^{\circ} - (100^{\circ} - x) \\ &= 180^{\circ} - 100^{\circ} + x^{\circ} \\ &= 80^{\circ} + x^{\circ}\end{aligned}$$

According to the statement

$$180^{\circ} - x^{\circ} - 40^{\circ} = 80^{\circ} + x^{\circ}$$

$$-x^{\circ} - x^{\circ} = 80^{\circ} + 40^{\circ} - 180^{\circ}$$

$$-2x^{\circ} = -60^{\circ}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

266


$$x^{\circ} = 30^{\circ}$$

$$\text{1st angle} = 30^{\circ}$$

$$\text{2nd angle} = 100^{\circ} - 30^{\circ}$$

$$= 70^{\circ}$$

6. Write the equation for the given triangle and solve it.



Sol: The sum of angles of any triangles = 180°


$$\begin{aligned}\text{The sum of angles of } \triangle ABC &= x + 90^{\circ} + 30^{\circ} \\ &= x + 120^{\circ}\end{aligned}$$

According to the statement

$$\begin{aligned}x + 120^{\circ} &= 180^{\circ} \\ x &= 180^{\circ} - 120^{\circ} \\ x &= 60^{\circ}\end{aligned}$$

7. Write down the angles marked with letters.

- 7(i) $m\angle a = 40^{\circ}$ (vertically opp. angles)




GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

267


7(ii) $m\angle c = 35^\circ$ (vertically opp. angles)

$m\angle d = 145^\circ$ (vertically opp. angles)




7(iii) $m\angle e = 29^\circ$ (vertically opp. angles)

$m\angle f = 151^\circ$ (vertically opp. angles)



7(iv) $m\angle b = 135^\circ$ (vertically opp. angles)




7(v) $m\angle q = 70^\circ$ (vertically opp. angles)

$m\angle p = 180 - 77^\circ$ (supplementary angles)
 $= 103^\circ$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

268



(vertically opp. angles) $m\angle r = m\angle p$

$$\hookrightarrow (i) \quad = 103^\circ$$

7(vi)

(vertically opp. angles) $m\angle k = 150^\circ$

(supplementary angles) $m\angle i + 150^\circ = 180^\circ$

$$m\angle i = 180^\circ - 150^\circ$$



7(vi)

$$m\angle i = 30^\circ \quad (i)$$

(vertically opp. angles) $m\angle j = m\angle i$ and

$$\text{from (i)} \quad m\angle j = 30^\circ$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th


269

7(vii)

(vertically opp. angles) $m\angle h = 40''$

(supplementary angles) $m\angle 40 + \angle g = 180''$

$$m\angle g = 180'' - 40$$



7(vii)

$$m\angle g = 140'' \quad (i)$$


(vertically opp angles) $m\angle i = m\angle g$ and

from (i) $= 140''$

(vertically opp. angles) $m\angle k = 138''$ 7(viii)

(supplementary angles) $m\angle p + 138'' = 180''$

$$m\angle p = 180'' - 138''$$




7(viii)

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pink Super One "General Math" 10th

270

$m\angle p = 42''$ (i)
 (vertically opp. angles) $m\angle l = m\angle p$ and
 from (i) $m\angle l = 42''$
 (vertically opp. angles) $m\angle p = 58''$ 7(ix)
 (supplementary angles) $m\angle N + 58'' = 180''$
 $m\angle N = 180'' - 58$



$m\angle N = 122''$ (i)
 (vertically opp. angles) $m\angle M = m\angle N$ and
 from (i) $= 122''$ 7(x)



GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

271

(supplementary angles) $m\angle a + 22^\circ = 180^\circ$

$$m\angle a = 180^\circ - 22^\circ$$

$$m\angle a = 158^\circ$$

(supplementary angles) $m\angle b + 68^\circ = 180^\circ$


$$m\angle b = 180^\circ - 68^\circ$$

$$m\angle b = 112^\circ$$

(i) **PARALLEL LINES**

Parallel lines are two straight lines in the same plane which never meet.

The lines a and b are parallel, we write $a \parallel b$.




(ii) **Properties of Parallel Lines**

(a) Two lines parallel to a third are parallel to each other.



(b) If three parallel lines are intercepted by two transversals in such a way that the two intercepts on one transversal are equal to each other, the two intercepts on the second transversal are also equal.




(c) If a line bisects one side of a triangle and is parallel to a second side, then it bisects the third side.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th


272



i.e. if $\triangle ABC$ with $\overline{BD} = \overline{DA}$, $\overline{DE} \parallel \overline{BC}$ then $\overline{AE} = \overline{CE}$

Transversal

A transversal is a line that intersects two lines in different points.



If a transversal "n" intersects two parallel lines a and b, the angles formed are identified as follows:

1. Four interior angles : $\angle 1, \angle 2, \angle 3, \angle 4$
2. Four exterior angles : $\angle 5, \angle 6, \angle 7, \angle 8$
3. Two pairs of alternate interior angles $\angle 1$ and $\angle 3$; and $\angle 4$
4. Two pairs of alternate exterior angles $\angle 5$ and $\angle 7$; $\angle 6$ and $\angle 8$
5. Two pairs of interior angles on the same side of the transversal: $\angle 2$ and $\angle 3$; $\angle 1$ and $\angle 4$.
6. Four pairs of corresponding angles: $\angle 3$ and $\angle 7$; $\angle 4$ and $\angle 8$; $\angle 2$ and $\angle 6$; $\angle 1$ and $\angle 5$.

Relation Between the Pairs of Angles

If two parallel lines are cut by a transversal, the corresponding angles are equal.


$$[\angle 1 = \angle 2, \angle 2 = \angle 3, \angle 1 = \angle 3]$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

273

- d) If two parallel lines are cut by a transversal, the alternate interior angles are equal.



$a \parallel b$, lines a and b are cut by the transversal c at point M and N to form the pairs of alternate interior angles.

$(\angle 1, \angle 2)$ and $(\angle 3, \angle 4)$

$\angle 1 = \angle 2, \angle 3 = \angle 4$

- e) If two parallel lines are intercepted by a transversal, then pairs of interior angles on the same side of transversal are supplementary.

$AB \parallel CD$, lines are cut by the transversal t , angles a, b, c and d are formed.

(i) $m\angle 2 = m\angle 4$

$m\angle 1 = m\angle 3$

(ii) $m\angle 3 = m\angle 7$

$m\angle 4 = m\angle 8$

$m\angle 6 = m\angle 2$

$m\angle 5 = m\angle 1$

(iii) $m\angle 7 = m\angle 5$

$m\angle 6 = m\angle 8$

(iv) $m\angle 2 + m\angle 3 = 180^\circ$

$m\angle 1 + m\angle 4 = 180^\circ$

(v) $m\angle 5 + m\angle 8 = 180^\circ$

$m\angle 6 + m\angle 7 = 180^\circ$


GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

274



1. Look at the given figure and answer the following questions.



- (a) The pair of alternative angles
- (b) The pair of corresponding angles
- (c) The pair of complementary angles
- (d) The pair of supplementary angles
- (e) The pair of vertical angles

Answers:

- a) The pair of alternative interior angles: $(\angle 1, \angle 2)$ and $(\angle 3, \angle 4)$
- b) The pair of corresponding angles:
 $\angle 1, \angle 6; \angle 3, \angle 8; \angle 2, \angle 7; \angle 5, \angle 4$
- c) The pair of complementary angles: No one
- d) The pair of supplementary angles:
 $(\angle 4, \angle 1); (\angle 4, \angle 7); (\angle 7, \angle 8); (\angle 8, \angle 1)$
 $(\angle 5, \angle 6); (\angle 5, \angle 2); (\angle 2, \angle 3); (\angle 3, \angle 6)$
- e) The pair of vertical angles:


GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

275

$(\angle 5, \angle 3); (\angle 6, \angle 2); (\angle 4, \angle 6); (\angle 1, \angle 7)$

2. Look at the given figure and answer the following questions.



- (a) The pair of alternative interior angles
- (b) The pair of corresponding angles
- (c) The pair of complementary angles
- (d) The pair of supplementary angles
- (e) The pair of vertical angles

Answers:

The alternate interior angles: $(\angle n, \angle l), (\angle m, \angle r)$

The corresponding angles:

$(\angle p, \angle n); (\angle m, \angle s); (\angle q, \angle r); (\angle l, \angle t)$

The complementary angles: No one

The supplementary angles:

$(\angle p, \angle m); (\angle p, \angle q); (\angle q, \angle l); (\angle l, \angle m)$

$(\angle n, \angle s); (\angle n, \angle r); (\angle r, \angle t); (\angle t, \angle s)$

The vertical angles: $(\angle p, \angle l); (\angle m, \angle q); (\angle n, \angle t); (\angle r, \angle s)$


3. Take a point 'X' outside a line \overline{DE} . Draw a line through

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

276

'X' which cuts \overline{DE} at some point. Making corresponding angles congruent draw a line parallel to \overline{DE} .



Steps of Construction:

- (i) Draw a line \overline{DE} .
- (ii) Take any point "X" which lay outside of \overline{DE} .
- (iii) Take any point "A" on \overline{DE} .
- (iv) Join "A" with "X" then extend it
- (v) Draw corresponding angles $\angle 1$ & $\angle 2$ with the help of compasses.
- (vi) Extend \overline{XT} on both sides.

$\overline{PQ} \parallel \overline{DE}$ which passes through X.

CONGRUENT AND SIMILAR FIGURES

Congruent Figures


The word congruent comes from Latin meaning "together agree". Two geometrical figures which have the same size and shape are congruent.

One figure is congruent to the other. The symbol for congruent is \cong . Thus two segments are congruent when they have the same size.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

277



In figure (i) and (ii)

$$m\overline{AB} = m\overline{PQ}$$

$$m\overline{BC} = m\overline{QR}$$

$$m\overline{CA} = m\overline{RP}$$


$$m\angle B = m\angle Q$$

$$m\angle C = m\angle R$$

$$m\angle A = m\angle P$$

Similar Figures

In the polygons below, the members of each pair are similar to each other.



In figure (i) and (ii)

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

. 278

$$m\angle A = 50^\circ, \quad m\angle P = 50^\circ$$

$$m\angle B = 45^\circ, \quad m\angle Q = 45^\circ$$

$$m\angle C = 85^\circ, \quad m\angle R = 85^\circ$$

$$\frac{m\overline{AC}}{m\overline{PR}} = \frac{4}{2} = \frac{2}{1}$$

$$\frac{m\overline{AB}}{m\overline{PQ}} = \frac{6}{3} = \frac{2}{1}$$

$$\frac{m\overline{BC}}{m\overline{QR}} = \frac{4.6}{2.3} = \frac{2}{1}$$



Tell Whether or not the Figures in Question 1-3 are Similar:

1. All squares; Yes
 all rectangles; No
 all regular hexagons. Yes

2. Two rectangles with sides 8, 12, 10 and 15.

Ans. These are similar figures

$$\frac{10}{15} = \frac{8}{12}$$

$$\frac{2}{3} = \frac{2}{3}$$

3. Two rhombuses with angles of 55° and 125° .

Ans. These are similar figures because the four.

4. The sides of a polygon are 5cm, 6cm, 7cm, 8cm, and 9cm.
 In a similar polygon the sides corresponding to 6cm is 12cm. Find the other sides of the second polygon.

Ans. According to the given condition the ratio among

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

279

corresponding sides is $\frac{6}{12} = \frac{1}{2}$.

$$\frac{6}{12} = \frac{1}{2} = \frac{7}{a} = \frac{8}{b} = \frac{5}{c} = \frac{4}{d} = \frac{9}{e}$$

$$\frac{7}{a} = \frac{1}{2} \Rightarrow a = 7 \times 2 = 14 \text{ cm}$$

$$\frac{8}{b} = \frac{1}{2} \Rightarrow b = 8 \times 2 = 16 \text{ cm}$$

$$\frac{5}{c} = \frac{1}{2} \Rightarrow c = 5 \times 2 = 10 \text{ cm}$$

$$\frac{4}{d} = \frac{1}{2} \Rightarrow d = 4 \times 2 = 8 \text{ cm}$$

$$\frac{9}{e} = \frac{1}{2} \Rightarrow e = 9 \times 2 = 18 \text{ cm}$$

5. The sides of a quadrilateral are 2cm, 4cm, 6cm, and 7cm.
 The longest side of a similar quadrilateral is 21cm. Find
 the other sides.

Ans. The longest side of quadrilateral is 7cm. Thus, the ratio
 between the biggest sides $\frac{7}{21} = \frac{1}{3}$.

According to the statement the ratio between the
 corresponding sides is $\frac{1}{3}$.

$$\frac{1}{3} = \frac{6}{a} = \frac{5}{b} = \frac{4}{c} = \frac{2}{d} \quad \text{Thus}$$

$$\frac{6}{a} = \frac{1}{3} \Rightarrow a = 6 \times 3 = 18 \text{ cm} \quad \text{therefore}$$

$$\frac{5}{b} = \frac{1}{3} \Rightarrow b = 5 \times 3 = 15 \text{ cm}$$

$$\frac{4}{c} = \frac{1}{3} \Rightarrow c = 4 \times 3 = 12 \text{ cm}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

280

$$\frac{2}{d} = \frac{1}{3} \Rightarrow d = 2 \times 3 = 6 \text{ cm}$$

While a, b, c, d are the rest sides of quadrilateral.

6. The sides of a polygon are 5cm, 2cm, 7cm, 3cm, 4cm. Find the sides of a similar polygon whose side corresponding to 2cm is 6cm. What is the ratio of the perimeters of these two polygons?

Sol: According to the statement the ratio between the correspondign sides is $\frac{2}{6} = \frac{1}{3}$.

$$\frac{1}{3} = \frac{5}{a} = \frac{7}{b} = \frac{3}{c} = \frac{4}{d} \quad \text{thus}$$

$$\frac{5}{a} = \frac{1}{3} \Rightarrow a = 5 \times 3 = 15 \text{ cm}$$

$$\frac{7}{b} = \frac{1}{3} \Rightarrow b = 7 \times 3 = 21 \text{ cm}$$

$$\frac{3}{c} = \frac{1}{3} \Rightarrow c = 3 \times 3 = 9 \text{ cm}$$

$$\frac{4}{d} = \frac{1}{3} \Rightarrow d = 4 \times 3 = 12 \text{ cm}$$

While a, b, c, d are the rest sides of polygon.

$$\begin{aligned} \text{The perimeter of 1st polygon} &= 5 + 2 + 7 + 3 + 4 \\ &= 21 \text{ cm} \end{aligned}$$

$$\begin{aligned} \text{The perimeter of 2nd polygon} &= 15 + 21 + 9 + 12 + 6 \\ &= 63 \text{ cm} \end{aligned}$$

$$\text{The ratio between the perimeter} = \frac{21}{63} = \frac{1}{3} \quad \therefore 1 : 3$$


7. What are the congruent pairs of corresponding sides and

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

281

corresponding angles?



8. Are all similar figures congruent? Explain why?

Sol: All similar figures are equal in size and shape.
 Therefore, similar figures are congruent.

9. Are all congruent figures similar? Explain why?

Sol: All congruent figures have same shape but differ in size.
 Therefore, congruent figures are not similar.

Exercise 7.4

1. Fill in the blanks.

(a) If $\triangle ABC \cong \triangle FDE$, then

(i) $\overline{AB} =$ _____ (ii) $\overline{BC} =$ _____

(iii) $\overline{AC} =$ _____ (iv) $m\angle A =$ _____

(v) $m\angle B =$ _____ (vi) $m\angle C =$ _____

(b) In $\triangle PQR$, the angle included between side PR and QR is _____


GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

282

- (c) In $\triangle ABC$, the side included between $\angle E$ and $\angle F$ is _____
- (d) If $\overline{AB} \cong \overline{QP}$, $m\angle B = m\angle P$, $\overline{BC} \cong \overline{PR}$, then by _____ condition, $\triangle ABC \cong \triangle QPR$
- (e) If $m\angle A = m\angle R$, $m\angle B = m\angle P$, $\overline{AB} \cong \overline{RP}$ then by _____ congruence condition, $\triangle ABC \cong \triangle RPQ$.

Answers:



(a)


- (i) $\overline{AB} \cong \overline{FD}$ (ii) $\overline{BC} \cong \overline{DE}$
 (iii) $\overline{AC} \cong \overline{FE}$ (iv) $\angle A \cong \angle F$
 (v) $\angle B \cong \angle D$ (vi) $\angle C \cong \angle E$

mid angle $\angle R$



(b)

mid side \overline{FE}




(c)

$$m\angle A = m\angle R$$

$$m\angle B = m\angle P$$

$$\overline{AB} \cong \overline{RP}$$



(d)


$$\triangle ABC \cong \triangle RPQ \quad \text{ASA} \cong \text{ASA}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

283

2. In figure, the pairs of corresponding equal parts in a pair of triangles are shown with similar markings. Specify the two triangles which become congruent. Also, write the congruence of two triangles in symbolic form.



Sol:

- (i) $\triangle ABC \cong \triangle DEF$ SSS \cong SSS
 (ii) $\triangle XYZ \cong \triangle DFE$ RHS \cong RHS
 (iii) $\triangle ABC \cong \triangle CDA$ ASA \cong ASA
 (iv) $\triangle PQT \cong \triangle SRT$ SAS \cong SAS

3. In figure, ABC and DBC are two triangles on a common base BC such that $\overline{AB} = \overline{DC}$ and , where A and D lie on the same side of BC . In $\triangle ADB$ and $\triangle DAC$, state the corresponding parts so that $\triangle ADB \cong \triangle DAC$.
 Which condition do you use to establish the congruence?


GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

284

If $m\angle DCA = 40^\circ$ and $m\angle BAD = 100^\circ$.

Find $\angle ADB$.



Sol: Now $\triangle ABC$ and $\triangle DCB$

common $\overline{BC} \cong \overline{CB}$

given $\begin{cases} \overline{AB} \cong \overline{DC} \\ \overline{AC} \cong \overline{DB} \end{cases}$

Now $\triangle ADB$ and $\triangle DAC$

$\overline{DA} \cong \overline{AD}$

$\overline{DB} \cong \overline{AC}$

$\overline{BA} \cong \overline{CD}$

$\triangle ADB \cong \triangle DAC$

SSS \cong SSS

$m\angle DCA = 40^\circ$ now

$m\angle BAD = 100^\circ$ and

$m\angle ABD = 40^\circ$ therefore

$m\angle ADB = 180^\circ - 100^\circ - 40^\circ$

$= 180^\circ - 140^\circ$

$= 40^\circ$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

285

4. Identify the following figure as congruent, similar or neither.




- Sol: (i) congruent (ii) congruent
 (iii) congruent

No one is similar

5. Identify the corresponding parts in $\triangle MNO$ and $\triangle PQR$.

- (i) $\overline{MN} \leftrightarrow$
 (ii) $\overline{NO} \leftrightarrow$
 (iii) $\overline{PR} \leftrightarrow$
 (iv) \angle



Answers:

- (i) $\overline{MN} \leftrightarrow \overline{PQ}$
 (ii) $\overline{NO} \leftrightarrow \overline{QR}$
 (iii) $\overline{PR} \leftrightarrow \overline{MO}$
 (iv) $\angle 1 \leftrightarrow \angle 4$

Quadrilaterals:


GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

=====

Pilot Super One "General Math" 10th


286

A quadrilateral is a polygon with four sides.




Parallelogram:

A parallelogram is a quadrilateral with two pairs of parallel sides.




Rectangle

A rectangle is a parallelogram containing a right angle.



Square

A square is an equilateral rectangle.



Properties of Congruency

Four Sides of a Square are Equal


ABCD is a square. Measure \overline{AB} , \overline{BC} , \overline{CD} and \overline{DA} . We find that.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

287


$$m\overline{AB} = m\overline{BC} = m\overline{CD} = m\overline{DA} = 2.8\text{cm.}$$



Four Angles of a Square are Right Angles

ABCD is a square. Measure angle A, B, C, D with protractor. We find that

$$m\angle A = m\angle B = m\angle C = m\angle D = 90^\circ$$




Diagonals of a Square Bisect Each Other:

Consider a square ABCD, the diagonals intersect at 'O'. We find that

$$m\overline{OA} = m\overline{OC} = 1.9\text{cm and}$$

$$m\overline{OB} = m\overline{OD} = 1.9\text{cm}$$



7.5.2 Opposite Sides of a Rectangle are Equal

Consider Rectangle

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th


288

Let us consider a rectangle $ABCD$.

$\overline{AB}, \overline{CD}$ and $\overline{AD}, \overline{BC}$ are opposite pairs of rectangle $ABCD$.

We find that $m\overline{AB} = m\overline{CD} = 4.5\text{cm}$ and

$m\overline{AD} = m\overline{BC} = 2.8\text{cm}$



$$m\overline{AB} = m\overline{CD} \quad (i)$$

$$m\overline{AD} = m\overline{BC} \text{ and}$$

$$m\angle A = m\angle B = m\angle C = m\angle D = 90^\circ \quad (ii)$$

$$(iii)$$

$$m\overline{OA} = m\overline{OC},$$

$$m\overline{OB} = m\overline{OD}$$

$$m\overline{OA} = m\overline{OC} = m\overline{OB} = m\overline{OD}$$

Properties of Parallelogram

The opposite sides of a parallelogram are equal

$ABCD$ is a parallelogram. $\overline{AB}, \overline{CD}$ and $\overline{AD}, \overline{BC}$ are pairs of opposite sides.

We find that

$$m\overline{AB} = m\overline{CD} = 3.9\text{cm} \quad \text{and}$$

$$m\overline{AD} = m\overline{BC} = 2.0\text{cm}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

289



- *The opposite angles of a parallelogram are equal.*

$ABCD$ is a parallelogram. $\angle A, \angle C$ and $\angle B, \angle D$ are pairs of opposite angles.

We find that

$$m\angle A = m\angle C = 70^\circ \text{ and}$$

$$m\angle B = m\angle D = 110^\circ$$




- *The diagonals of a parallelogram bisect each other.*

A parallelogram $ABCD$, the diagonals \overline{AC} and \overline{BD}

intersect at O . We find that

$$m\overline{OA} = m\overline{OC} = 2.5\text{cm}$$

and $m\overline{OD} = m\overline{OB} = 2.5\text{cm}$



GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

290

Exercise 4

I. Fill in the blanks:


- (i) A parallelogram that contains a right angle is _____
- (ii) An equilateral rectangle is a _____
- (iii) A polygon with four sides is a _____
- (iv) The diagonals of a parallelogram _____ each other.
- (v) The opposite angles of a parallelogram are _____

Answers:

- (i) Rectangle (ii) Square (iii) Quadrilateral
- (iv) Bisect (v) Congruent



A circle is the set of points in a plane which are at a constant distance from a fixed point in the plane.



Centre

The fixed point C is called the centre of the circle.

Radial Segment

P is any point on the circumference of the circle with centre O. \overline{OP} is called the radial segment of the circle.

Radius

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

291

A radius of a circle is the length of a segment joining the centre to any point on the circle. In the given figure is $\overline{mC'P}$ the radius. Usually represented by 'r'.

Chord

A chord of a circle is a segment connecting any two points on the circle. In the given figure \overline{AB} is a chord.


Segment of a Circle

A chord \overline{AB} of a circle divides the circle in two parts.

These are called segment of the circle.

Minor Segment

The included area between minor arc and the chord is minor segment.



Major Segment

The included area between major arc and chord is called major segment.

Diameter

A diameter of a circle is a chord that passes through the centre. The length of a diameter of a circle is twice the length of the radius of the same circle.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

=====

Pilot Super One "General Math" 10th

292


$$\boxed{\text{Diameter} = 2 \times \text{radius}}$$

Equal Circles

Equal circles are circles having equal radii or equal diameter


Sector

A sector is the circular region bounded by an arc of a circle and its two corresponding radial segments is called a sector of the circle. In the figure, region AOB is the sector of the circle with centre at O .



Secant

A secant is a line which intersects a circle in two points.



GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

293

A tangent to a circle is the line perpendicular to radius of the circle at its outer extremity.

The point on the circle at which the radius and tangent meet is known as the Point of Contact or Point of Tangency.


Angle in a Semi-Circle is a Right Angle

- 1- Draw a line-segment \overline{AB} of any length. Mark the mid point of \overline{AB} as O .
- 2- Draw a semi-circle on \overline{AB} with radius \overline{OA} .
- 3- Take any point C on the semi-circle. Join A with C and B with C .

Thus, $\angle ACB$ is an angle in the semi-circle APB .

- 4- Now take a protractor and place it along \overline{AC} so that the centre of the protractor falls on C .

We note that the measure of the $\angle ACB$ by looking at the marking on the protractor corresponding to arm \overline{CB} of $m\angle ACB$ is of 90° , i.e. $m\angle ACB = 90^\circ$ or a right angle. Thus, angle in a semi-circle is a right angle.



Angles in the Same Segment are Equal:

Draw a circle with centre ' O '. Take two points B and C on the circle and join them. \overline{BC} divides the circle into two parts.

Draw angles, $\angle BAC$ and $\angle BDC$ in the same segment as shown in the figure. Take a sheet of tracing paper and

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)


Pilot Super One "General Math" 10th

294

make a trace copy of $\angle BAC'$. Place the trace copy of $\angle BAC'$ on $\angle BDC'$.


A falls on D and \overline{AB} falls on $\overline{DC'}$.

So that we observe that \overline{BD} falls on $\overline{AC'}$. Thus $\angle BAC' = \angle BDC'$, this shows that angles in the same segment are equal.



Central Angle

The central angle of a minor arc of a circle is double that the angle subtended by corresponding major arc.



$$m\angle AOB = m2\angle ACB$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 7)

Pilot Super One "General Math" 10th

295

10TH CLASS

1- Fill in the blanks:

- (i) In a plane the set of points whose distance from a fixed point is same is called _____.
- (ii) The distance of a point of a circle from its centre is called _____.
- (iii) A line segment whose end points lie on the circle is called _____.
- (iv) A chord that passes through the centre of the circle is called _____.
- (v) Half of a circle is called _____.
- (vi) An arc which is greater than a semicircle is called _____.
- (vii) One and only one circle can be constructed with a given centre and given _____.
- (viii) A region bounded by an arc and two of its radial segments is called _____.
- (ix) A straight line that intersects a circle at two points is called _____.
- (x) Angle in a semi-circle is a _____.

Answers:

- | | |
|------------------|-----------------|
| (i) circle | (ii) radius |
| (iii) chord | (iv) diameter |
| (v) semicircle | (vi) major arc |
| (vii) radius | (viii) sector |
| (ix) secant line | (x) right angle |

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)


Print Super One "General Math" 10th

302

EXERCISE 8.1

- Q.1. Draw a triangle ABC in which $\overline{BC} = 5.4\text{cm}$,
 $\overline{AB} = 4.3\text{cm}$ and $\overline{AC} = 3.9\text{cm}$. Find the in centre

Sol.



Steps of Construction:

- (i) Draw a line segment $\overline{BC} = 5.4\text{cm}$
- (ii) With B as centre draw an arc of radius 4.3 cm.
- (iii) With C as centre draw an arc of radius 3.9cm which intersect the first arc at A
- (iv) Join A with B and C.
 $\triangle ABC$ is the required triangle.
- (v) Draw perpendicular bisectors \overline{LM} and \overline{PQ} of the sides \overline{AB} and \overline{BC} which intersect each other at O.

Point O is the required incentre.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)

Pilot Super One "General Math" 10th

301

Practical Geometry

- ▶ Construction of a Triangle
- ▶ Construction of a Quadrilateral
- ▶ Tangent to a Circle

After completing of this unit, the students will be able to:
construct a triangle having:

Two sides and the included angle.

One side and two of the angles.

Two of its sides and angle opposite to one of them with all the three possibilities).

Draw:

Angle bisectors. ○ Altitudes.

Medians of a given triangle and verify their concurrency.

Construct a rectangle when:

○ Two sides are given.

○ Diagonal and one side are given.

Construct a square when its diagonal is given.

Construct a parallelogram when two adjacent sides and the angle included between them is given.

locate the centre of given circle.

draw a circle passing through three given non-collinear points.

draw a tangent to a given circle from a point P when P lies.

○ On the circumference,

○ Outside the circle.

Draw:

○ Direct common tangent or external tangent.

○ Transverse common tangent or internal tangent to two equal circles.

Draw a tangent to:

○ Two unequal touching circles.

○ Two unequal intersecting circles.


GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)

Pilot Super Time "General Math" 10th

303

- Q.2. Construct $\triangle ABC$ in which $m\overline{BC} = 4.6\text{cm}$,
 $\angle B = 110^\circ$ and $m\overline{AB} = 5\text{cm}$. Draw the
 perpendicular bisectors of its sides.

Sol.




Steps of Construction:

- (i) Draw a line segment $\overline{AB} = 5\text{cm}$.
- (ii) At point B, draw an angle 110° with the help of compasses.
- (iii) Cut $m\overline{BC} = 4.6\text{cm}$ at \overline{BH} .
- (iv) Join "C" with A. $\triangle ABC$ is the required triangle.
- (v) Draw perpendiculars EF, GH and LM of the sides \overline{AB} , \overline{BC} and \overline{AC} respectively. They meet each other at point "O".

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)

- Q.3.** Draw an equilateral $\triangle ABC$ in which
 $m\overline{AB} = m\overline{BC} = m\overline{AC} = 5\text{cm}$. Draw its altitudes and
 measure their lengths are they equal?

Sol.



Steps of Construction:

- (i) Draw a line segment $m\overline{BC} = 5\text{cm}$.
- (ii) Draw arcs of radius 5 cm with taking centre B and C, which intersect each other at A.

- (iii) Join A with B and C.

$\triangle ABC$ is the required equilateral.

$AB \perp BC$, $BQ \perp CA$ and $AB \perp CR$

And

- (v) $m\overline{AP} = m\overline{BC} = m\overline{CR} = 4.2\text{cm}$

All the altitudes are equal in lengths.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)

=====


Pilot Super One "General Math" 10th

305

Q.4. Construct a $\triangle ABC$ in which $m\overline{BC} = 5.4\text{cm}$,

$\angle B = 65^\circ$ and $m\angle C = 55^\circ$. Find the centroid of the triangle.

Sol.



Steps of Construction:

- (i) Draw a line segment \overline{BC} 5.4cm.
- (ii) Now draw an angle of 65° at point B & 55° at point C.
 \overline{BP} and \overline{CQ} intersect each other at point A.
- (iii) $\triangle ABC$ is the required triangle.
- (iv) M and N are the mid points of \overline{BC} and \overline{AC} .
- (v) \overline{AM} and \overline{BN} are the medians which intersect each other at point G.

Thus point "G" is the required centroid of triangle.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)


=====

Pilot Super One "General Math" 10th

306

- Q.5.** Draw an equilateral triangle each of whose sides is 5.3cm. Draw its medians. Are they equal?

Sol.



Steps of Construction:

- (i) Draw a line segment \overline{BC} 5.3cm.
- (ii) Taking B and C as centre draw two arcs which intersect each other at point A.
- (iii) Join point A with B and C.
 $\triangle ABC$ is the required equilateral.
- (iv) Draw medians of sides \overline{AB} , \overline{CA} , \overline{BC} at points P, Q & R.
- (v) Join A with P, B with Q and C with R.
 \overline{AP} , \overline{BQ} , \overline{CR} are the required medians.
- (vi) $m\overline{AP} = m\overline{BQ} = m\overline{CR} = 4.5\text{cm}$

So that the medians are equal in lengths.


GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)

Pilot Super One "General Math" 10th

307

Q.6. Draw an equilateral triangle with length of each side 6cm.

Sol.




Steps of Construction:

- (i) Draw a line segment $\overline{AB} = 6\text{cm}$.
 - (ii) Taking A and B as centre draw two arcs of radius 6cm each. They intersect each other at point C.
 - (iii) Join point C with A and B.
- $\triangle ABC$ is the required equilateral.

Q.7. Construct a triangle ABC with base length 5cm and the angles at both ends of the base are 45° and 60° respectively.

Sol.



GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)

Pilot Super One "General Math" 10th

308


Steps of Construction:

- (i) Draw a line segment $\overline{AB} = 5\text{cm}$.
- (ii) Draw $m\angle BAP = 60^\circ$ at point A.
- (iii) Draw an angle $m\angle ABQ = 45^\circ$ at point B.
- (iv) \overline{AP} and \overline{BQ} intersect each other at point C.

ABC is the required triangle.

- Q.8. Draw an isosceles triangle with length of the equal sides 5cm and the angle included between them is 60° .

Sol.



Steps of Construction:

- (i) Draw a line segment $\overline{BC} = 5\text{cm}$.
- (ii) At point B, draw $m\angle ABC = 60^\circ$ using compasses.
- (iii) Cut $\overline{BA} = 5\text{cm}$.
- (iv) Join point A with C.

ABC is the required isosceles.


GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)

Pilot Super One "General Math" 10th

309

Q.9. Construct a rectangle whose adjacent sides are 4cm and 3cm.

Sol.



Steps of Construction:

- (i) Draw a line segment $\overline{AB} = 4\text{cm}$.
- (ii) At points A and B, draw right angles with the help of compasses.
- (iii) Cut $m\overline{AD} = m\overline{BC} = 3\text{cm}$.
- (iv) Join point C with D.

ABCD is the required rectangle

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)


=====

Pilot Super One "General Math" 10th

310

Q.10. Construct a rectangle whose one side is 6cm and an adjacent diagonal of 9cm.

Sol.



Steps of Construction:

- (i) Draw a line segment $\overline{AB} = 6\text{cm}$.
- (ii) Draw right angle at points A and B.
- (iii) Taking centre as A draw an arc of radius 9cm which intersect \overline{BP} at C.
- (iv) Cut \overline{BC} , \overline{AD} at \overline{AQ} .
- (v) Join point C with D.

$ABCD$ is the required rectangle.


GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)

Pilot Super One "General Math" 10th

311

Q.11. Construct a square whose one side is 5cm.

Sol.



Steps of Construction:

- (i) Draw a line segment $\overline{AB} = 5\text{cm}$.
- (ii) At points A and B, draw right angle with the help of compasses.
- (iii) Cut $m\overline{BC} = m\overline{AD} = 5\text{cm}$ at \overline{AQ} and \overline{BP} .
- (iv) Join C with D

$ABCD$ is the required square.


GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)

Pilot Super One "General Math" 10th

312

Q.12. Construct a square whose one side is 3.5cm.

Sol.



Steps of Construction:

- (i) Draw a line segment $\overline{AB} = 3.5\text{cm}$.
- (ii) At points A and B, draw right angle with the help of compasses.
- (iii) Cut $m\overline{BC} = m\overline{AD} = 3.5\text{cm}$ at \overline{AQ} and \overline{BR} .
- (iv) Join C with D.

$ABCD$ is the required square.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)


=====

Pikt Super One "General Math" 10th

313

- Q.13. Construct a rectangle whose two adjacent sides measure 5cm and 4cm and their included angle is 90° .

Sol.



Steps of Construction:

- (i) Draw a line segment $\overline{AB} = 5\text{cm}$.
- (ii) At point A, draw right angle with the help of compasses.
- (iii) Cut $m\overline{BC} = m\overline{AD} = 4\text{cm}$ at \overline{AR} and \overline{BP} .
- (iv) Join C with D.

$ABCD$ is the required rectangle.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)


=====

Pilot Super One "General Math" 10th

313

- Q.13. Construct a rectangle whose two adjacent sides measure 5cm and 4cm and their included angle is 90° .

Sol.



Steps of Construction:

- (i) Draw a line segment $\overline{AB} = 5\text{cm}$.
- (ii) At point A, draw right angle with the help of compasses.
- (iii) Cut $m\overline{BC} = m\overline{AD} = 4\text{cm}$ at \overline{AR} and \overline{BP} .
- (iv) Join C with D.

$ABCD$ is the required rectangle.


GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)

Pilot Super One "General Math" 10th

314

Q.14. Draw a rectangle whose one side is 8cm and the length of each diagonal is 10cm.

Sol.



Steps of Construction:

- (i) Draw a line segment $\overline{AB} = 6\text{cm}$.
- (ii) At point A and B, draw right angle with the help of compasses.
- (iii) Draw an arc of radius 10cm with taking centre point "A" which intersect \overline{BP} at point C.
- (iv) Now, draw an arc of radius cm again with taking centre at point B. Which intersect \overline{AQ} at point D.
- (v) Joint point C with D.

$ABCD$ is the required rectangle.


GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)

Pilot Super One "General Math" 10th

315

Q.15. Draw a rectangle ABCD in which $m\overline{AB} = 6.5\text{cm}$ and $m\overline{AD} = 4.8\text{cm}$ and $m\angle BAD = 90^\circ$. Measure its diagonals.

Sol.



Steps of Construction:

- (i) Draw a line segment $\overline{AB} = 6.5\text{cm}$.
- (ii) At point A and B, draw right angle with the help of compasses.
- (iii) Intersect $m\overline{BC} = m\overline{AD} = 4.8\text{cm}$ at \overline{AP} and \overline{BQ} .
- (iv) Join C with D.
- (v) ABCD is the required rectangle.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)

Pilot Super One "General Math" 10th


316

Q.16. Name the following quadrilaterals when:

Questions	Answers
(i) The diagonals are equal and the adjacent sides are unequal	Rectangle
(ii) The diagonals are equal and the adjacent sides are equal	Square
(iii) All the sides are equal and one angle is 90°	Square
(iv) All the angles are equal and the adjacent sides are unequal	Rectangle

Q.17. Construct a rectangle with sides 10cm and 6cm.

Sol.



Steps of Construction:

- Draw a line segment $\overline{AB} = 10\text{cm}$.
- At point A and B, draw right angle with the help of compasses.
- Intersect $m\overline{BC} = m\overline{AD} = 6\text{cm}$ at \overline{AQ} and \overline{BP} .
- Join C with D.

$ABCD$ is the required rectangle.


GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)

Pilot Super One "General Math" 10th

317

Q.18. Construct a square with side of length 6cm.

Sol.



Steps of Construction:

- (i) Draw a line segment $\overline{AB} = 6\text{cm}$.
- (ii) Points A and B, draw right angle with the help of compasses.
- (iii) Cut $\overline{AP} = \overline{BR} = 6\text{cm}$ at \overline{AD} and \overline{BC} .
- (iv) Join C with D.

$ABCD$ is the required square.

Q.19. Name the following triangles.

Questions	Answers
(i) With all the three sides equal in length.	Equilateral triangle
(ii) With two sides equal in length.	Isosceles triangle
(iii) None of the sides is equal to the other.	Scalene triangle

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)


Pilot Super One "General Math" 10th

318

Q.20. Draw a circle with centre O and radius 5cm . Explain

the steps necessary to draw a segment of the circle.

Sol.



Steps of Construction:

- (i) Take any point O .
- (ii) Taking centre with " O ", draw an arc of radius 5cm .
- (iii) Now, take any diameter \overline{PQ} .

Conclusions: \overline{PQ} has divided the circle into two parts. The major segment part and minor segment part.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)


=====

Pilot Super One "General Math" 10th

319

Q.21. Draw a circle with center O and any radius. Draw the diameter AB and shade one semicircular region.

Sol.



Steps of Construction:

- (i) Take any point O .
- (ii) With taking " O " as centre, draw a circle with suitable radius
- (iii) Draw \overline{AOB} as diameter.

Conclusion: The circle has divided into two parts. Now, shaded the half part.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)


=====

Pilot Super One "General Math" 10th

320

Q.22. Show four angles in a semi-circular region of question 21.

Sol.



Steps of Construction:

- (i) Draw a circle with suitable radius and marks its centre as "O".
- (ii) Draw a diameter \overline{AOB} .
- (iii) Take a point C, D, E, F at half curved area.
- (iv) Join these points with A and B.

Conclusion: $\angle ACB, \angle ADB, \angle AEB, \angle AFB$ are the required four angles.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)


=====

Pilot Super One "General Math" 10th

321

Q.23. Draw a circle of radius 2cm with center O . Draw a chord and shade the portion showing major arc.

Sol.



Steps of Construction:

- (i) Take any point O .
- (ii) With taking " O " as centre, draw a circle with radius 2cm.
- (iii) Take \overline{AB} as chord.

Thus, \widehat{ACB} is the major arc.

- (iv) In figure \widehat{ACB} (major arc) is quite prominent.


GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)

Pilot Super One "General Math" 10th

322

- Q.24. Draw a circle of radius 2.5cm with center at O . Draw a chord and shade the portion showing the minor arc of the circle.


Sol.



Steps of Construction:

- (i) Take any point " O ".
 - (ii) With taking " O " as centre, draw a circle with radius 2.5cm
 - (iii) Draw \overline{AB} as chord.
 - (iv) The major arc \widehat{ACB} is quite prominent in the figure.
- Q.25. Draw a semi-circle with diameter 4cm and center at O .

Sol.



GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)


Pilot Super One "General Math" 10th

323

Steps of Construction:

- (i) Draw a line segment $\overline{AB} = 4\text{cm}$.
 - (ii) Now take "O" as centre, draw a semi-circle with radius $m\overline{AO}$ or $m\overline{OB}$.
 - (iii) With taking "O" as centre $m\overline{OA}$ or $m\overline{OB}$.
 - (iv) ACB is the required semi-circle.
- Q.26.** Draw a circle passing through the vertices of a square of side 3cm.

Sol.



Steps of Construction:

- (i) Draw a line segment $\overline{AB} = 3\text{cm}$.
- (ii) At points A and B, draw right angles at each
 (∴ $m\overline{AD} = m\overline{BC} = 3\text{cm}$)
- (iii) Join C with D.
- (iv) Draw two diagonals \overline{AC} and \overline{BD} which intersect each other at O.
- (v) Taking "O" as centre, draw a circle with radius $m\overline{OB}$ or $m\overline{OA}$ or $m\overline{OC}$ or $m\overline{OD}$.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)


=====

Pilot Super One "General Math" 10th

324

Q.27. In a right triangle ABC , $m\overline{AB} = 3cm$ and $m\overline{BC} = 4cm$
with right angle at B . Draw a circle through A , B and C .

Sol.



Steps of Construction:

- (i) Draw a line segment $\overline{AB} = 3cm$.
- (ii) At point B , draw right angle with the help of compasses.
- (iii) Cut $m\overline{BC} = 4cm$.
- (iv) Join C with A .
 ABC is the required triangle.
- (v) Now, draw perpendicular bisector of sides \overline{AB} and \overline{BC}
which cut each other at point " O ".
- (vi) With taking " O " as centre draw a circle with radius $m\overline{OB}$
or $m\overline{OC}$ or $m\overline{OA}$ respectively.

Conclusion: The circle is passing through the vertices (A , B and C)


GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)

Pilot Super One "General Math" 10th

325

Q.28. Draw a circle passing through the three vertices of an equilateral triangle with length of each side 4cm.

Sol.



Steps of Construction:

- (i) Draw a line segment $\overline{AB} = 4\text{cm}$.
- (ii) Taking A and B as centre, draw two arcs of radius 4 cm each. They intersect each other at point "C".
- (iii) Join C with points A and B.
 ABC is the equilateral triangle.
- (iv) Now, draw a perpendicular bisectors \overline{RT} and \overline{PQ} of sides \overline{AB} and \overline{BC} respectively.
 They meet each other at point "O".
- (v) With taking "O" as centre, draw a circle with radius $m\overline{OC}$ or $m\overline{OA}$ or $m\overline{OB}$ respectively.

Conclusion: The circle is passing through the vertices (A, B, C) of triangle.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)

Pilot Super One "General Math" 10th

326

REVIEW QUESTIONS

1- *Encircle the correct answer:*

1- The number of medians in a triangle is:

- (a) 1 (b) 2
(c) 3 (d) 4

2- The number of altitudes in a triangle is:

- (a) 1 (b) 2
(c) 3 (d) 4

3- The number of angle bisectors in a triangle is:

- (a) 1 (b) 2
(c) 3 (d) 4

4- The number perpendicular bisectors of the side of a triangle is:

- (a) 1 (b) 2
(c) 3 (d) 4

5- The angle bisectors of a triangle are:

- (a) concurrent (b) collinear
(c) perpendicular (d) non-concurrent

6- The medians of a triangle are:

- (a) concurrent (b) collinear
(c) non-concurrent (d) 4

7- The altitudes of a triangle are:

- (a) concurrent (b) collinear
(c) non-collinear (d) 5

8- A line joining one vertex of a triangle to the mid point of its opposite sides is called:

- (a) angle bisector (b) altitude
(c) median (d) side bisector

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 8)

Pilot Super One "General Math" 10th



327

- 9- A line joining one vertex of a triangle and perpendicular to its opposite side is called:
 (a) angle bisector (b) median
 (c) altitude (d) side bisector
- 10- A line coplanar with a circle and intersecting the circle at one point only is called:
 (a) tangent line (b) median
 (c) altitude (d) normal line

Answers:

1-	(c)	2-	(c)	3-	(c)	4-	(c)	5-	(a)
6-	(a)	7-	(a)	8-	(c)	9-	(c)	10-	(a)

11- Fill in the blanks.

- 1- The altitudes of a triangle are 
- 2- The medians of a triangle are 
- 3- The angle bisector of a triangle are _____.
- 4- The perpendicular bisector of the three sides of a triangle are _____.
- 5- The line joining one vertex of a triangle and perpendicular to its opposite side is called _____ of a triangle.
- 6- A line joining one vertex of a triangle to the midpoint of its opposite side is called _____ of a triangle.
- 7- A line bisecting the angle of a triangle is called the _____.
- 8- Every triangle has _____ altitudes.
- 9- Every triangle has _____ median.
- 10- Every triangle has _____ right bisectors.

Answers:

1-	concurrent	2-	concurrent	3-	concurrent	4-	concurrent	5-	altitude
6-	median	7-	angle bisector	8-	three	9-	three	10-	three

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

=====



I. Find the third side of each right triangle with legs a and b and hypotenuse c .

(i) $a = 3, b = 4, c = ?$

(ii) $a = 5, c = 13, b = ?$

(iii) $b = 5, c = 61, a = ?$

Solution:

(i) $a = 3$

$b = 4$

$c = ?$

$c^2 = a^2 + b^2$ By Pythagoras theorem

$c^2 = (3)^2 + (4)^2$ Putting values of a, b

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

331

$$c^2 = 9 + 16$$

$$c^2 = 25$$

$$\sqrt{c^2} = \sqrt{25} \quad \text{Taking square root}$$

$$\boxed{c=5}$$

(ii) $a = 5, \quad c = 13, \quad b = ?$

By Pythagoras theorem

$$c^2 = a^2 + b^2$$

$$b^2 = c^2 - a^2 \quad \text{or}$$

Putting values of a, c

$$b^2 = (13)^2 - (5)^2$$

$$b^2 = 169 - 25$$

$$b^2 = 144$$

$$\sqrt{b^2} = \sqrt{144} \quad \text{Taking square root}$$

$$\boxed{b=12}$$

(iii) $b = 5$

$$c = 61$$

$$a = ?$$

By Pythagoras theorem

$$c^2 = a^2 + b^2$$

$$a^2 = c^2 - b^2 \quad \text{therefore}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Paper Super One "General Math" 10th

332

Putting values of b, c

$$a^2 = (61)^2 - (5)^2$$

$$a^2 = 3721 - 25$$


$$a^2 = 3696$$

$$a^2 = 3696 \quad \text{Taking square root}$$

$$\sqrt{a} = \sqrt{16 \times 231}$$

$$a = 4\sqrt{231}$$

2. If the legs of a right triangle are $2ab$ and $a^2 - b^2$, prove that hypotenuse is $a^2 + b^2$.



sol

$$\text{one side} = 2ab$$

$$\text{2nd side} = a^2 - b^2$$

$$\text{Hypotenuse} = ?$$

By Pythagoras theorem

$$(\text{Hypotenuse})^2 = (2ab)^2 + (a^2 - b^2)^2$$

$$\begin{aligned} \text{Hypotenuse} &= \sqrt{(2ab)^2 + (a^2 - b^2)^2} \\ &= \sqrt{a^4 + b^4 + 2a^2b^2 + 4a^2b^2} \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th


333

$$= \sqrt{a^4 + b^4 + 2a^2b^2}$$

$$= \sqrt{(a^2 + b^2)^2}$$

$$\text{Hypotenuse} = (a^2 + b^2)$$

3. Find the hypotenuse of the right isosceles triangle each of whose legs is l .



Sol:

$$m\overline{AC} =$$

$$m\overline{BC} =$$

$$m\overline{AB} = ?$$

By Pythagoras theorem

$$(m\overline{AB})^2 = (m\overline{AC})^2 + (m\overline{BC})^2$$

$$= l^2 + l^2$$

$$= 2l^2$$

$$\sqrt{m\overline{AB}^2} = \sqrt{2l^2} \quad \text{Taking square root}$$


$$m\overline{AB} = \sqrt{2} l \text{ units}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

334

4. Find the hypotenuse of a right isosceles triangle whose legs are 8cm



Sol:

$$a = 8 \text{ cm} \quad \text{here}$$

$$b = 8 \text{ cm}$$

$$c = ?$$

By Pythagoras theorem

$$c^2 = a^2 + b^2$$

Putting values of a, b

$$c^2 = (8)^2 + (8)^2$$

$$c^2 = 64 + 64$$

$$c^2 = 128$$

$$\sqrt{c^2} = \sqrt{128} \quad \text{Taking square root}$$

$$c = \sqrt{8 \times 8 \times 2}$$

$$c = 8\sqrt{2} \text{ cm}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

335

5. If the numbers represent the lengths of the sides of a triangle, which triangles are right triangles?

(i) 3, 4, 5

(ii) 9, 17, 25

(iii) 11, 61, 60

Sol: Length of sides 3, 4, 5

We observe that, is the square of big side equal or not equal the square of the other two sides? Then these sides will be right angle Δ otherwise not.

$$\begin{array}{l|l} (5)^2 = 25 & (3)^2 + (4)^2 \\ & = 9 + 16 \\ & = 25 \end{array}$$

These sides are sides of right angle Δ .

(ii)

$$\begin{array}{l|l} (25)^2 = 625 & (9)^2 + (17)^2 \\ & = 81 + 289 \\ & = 370 \end{array}$$

$625 \neq 370$

Sides 9, 17, 25 are not the sides of right angle triangle.

(iii)

$$\begin{array}{l|l} (61)^2 = 3721 & (11)^2 + (60)^2 \\ & = 121 + 3600 \\ & = 3721 \end{array}$$

Sides 11, 61, 60 are the sides of right angle triangle.


GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

336

6. $\triangle ABC$ is right angled at C . If $m\overline{AC} = 9\text{ m}$ and $m\overline{BC} = 12\text{ m}$, find the length \overline{AB} , using Pythagoras theorem

Sol In $\triangle ABC$



$$b = 9\text{ cm}$$

$$a = 12\text{ cm}$$

$$c = ?$$

By Pythagoras theorem

$$c^2 = a^2 + b^2$$

Putting values of a, b

$$c^2 = (12)^2 + (9)^2$$

$$= 144 + 81$$

$$c^2 = 225$$

$$\sqrt{c^2} = \sqrt{225} \quad \text{Taking square root}$$


$$\boxed{c = 15} \text{ cm}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

337

7. The hypotenuse of a right triangle is 25cm. If one of the sides are of length 24cm, find the length of the other side.
Sol:



$$c = 25 \text{ cm} \quad \text{here}$$

$$b = 24 \text{ cm}$$

$$a = ?$$

By Pythagors theorem

$$c^2 = a^2 + b^2$$

$$a^2 = c^2 - b^2 \quad \text{or}$$

Putting values of b, c

$$a^2 = (25)^2 - (24)^2$$

$$= 625 - 576$$

$$a^2 = 49$$

$$\sqrt{a^2} = \sqrt{49} \quad \text{Taking square root}$$

$$\boxed{a = 7 \text{ cm}}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)


=====

Pilot Super One "General Math" 10th

338

8. A ladder 17m long when set against the wall of a house just reaches a window at a height of 15m from the ground. How far is the lower end of the ladder from the base of the wall?

Sol:



Lower end of the ladder from the wall = a

$$b = 15\text{m} \quad \text{here}$$

$$c = 17\text{m}$$

$$a = ?$$

By Pythagoras theorem

$$a^2 = c^2 - b^2$$

Putting values of c, b

$$a^2 = (17)^2 - (15)^2$$

$$= 289 - 225$$

$$a^2 = 64$$

$$\sqrt{a^2} = \sqrt{64} \quad \text{Taking square root}$$

$$\boxed{a = 8} \text{m}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)


=====

Pilot Super One "General Math" 10th

339

9. The two legs of a right triangle are equal and the square of the hypotenuse is 50. Find the length of each leg.

Sol:



$$c = 50 \quad \text{here}$$

$$a = b$$

By Pythagoras theorem

$$c^2 = a^2 + b^2$$

$$c^2 = a^2 + a^2 \quad (a = b)$$

$$c^2 = 2a^2$$

$$2a^2 = c^2 \quad \text{or}$$

Putting values of c^2

$$2a^2 = (50)^2$$

$$a^2 = \frac{2500}{2}$$

$$a^2 = 1250$$

$$\sqrt{a^2} = \sqrt{1250} \quad \text{taking square root}$$

$$a = \sqrt{625 \times 2}$$

$$a = 25\sqrt{2}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

340

Length of each side = $25\sqrt{2}$ units

10. The sides of a triangle are 15cm, 36cm and 39cm. Show that it is a right angled triangle.

Sol:

The length of big side = 39 cm

The length of small side = 15cm, 36cm

$$(\text{Lengths of big side})^2 = (39)^2$$

$$= 1521 \quad (i)$$

$$\text{The sum of the square of small sides} = (15)^2 + (36)^2$$

$$= 225 + 1296$$

$$\text{Now its prove that} = 1521 \quad (ii)$$

These lengths are the lengths of right angle triangle.

Area

The surface inside the boundary of a shape is called area.

We see that, Is the big side of square equal or not equal the square of the other two sides? Then these sides will be sides of right angle triangle otherwise not.

Area of a Triangle when all the three sides are given

A triangle ABC with sides a, b, c and

$$2S = a + b + c \Rightarrow S = \frac{a + b + c}{2},$$

where 'S' is half the perimeter of a triangle.

Then area of any triangle is $A = \sqrt{S(S-a)(S-b)(S-c)}$.

This is called *Hero's Formula* for finding the area of a triangle.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

341

Example 1:

1. A verandah 40m long, 15m wide is to be paved with stones each measuring 6cm by 5cm. Find the number of stones.

Sol: Length of verandah = 40m

Width of verandah = 15m

Area of veranda = 40×15

= 600 sq. m

Area of one stone = 6×5 Now

= 30 sq. m


$$\text{Number of stones} = \frac{\text{Total area}}{\text{area of one stone}}$$

$$= \frac{600}{30}$$

Number of stones = 20 tiles

2. How many tiles of 40cm² will be required to pave the footpath 1m wide carried round the outside of a grassy plot 28m by 18m?

Sol: Tiles fixed round the out side of a plot therefore,



External length = $28 + 1 + 1$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

342

$$= 30 \text{ m}$$

$$\text{External width} = 18 + 2 = 20$$

$$= 20 \text{ m}$$

$$\text{Area of plot with path} = 30 \times 20$$

$$= 600 \text{ sq. m}$$

$$\text{The area of plot} = 28 \times 18$$

$$= 504 \text{ sq. m}$$

$$\text{Area of path} = 600 - 504$$

$$= 96 \text{ sq. m}$$

$$\text{Area of one tile} = 40 \text{ sq. m}$$

$$= \frac{40}{100 \times 100} \text{ sq. m}$$

$$\text{Number of tiles} = \frac{\text{Area of Path}}{\text{Area of one tile}}$$

$$= \frac{96}{\frac{40}{100 \times 100}} = \frac{96 \times 100 \times 100}{40}$$

$$\text{Number of tiles} = 24000$$

3. Find the area of a room 5.49m long and 3.87m wide. What is the cost of carpeting the room if the rate of carpet is Rs 10.50 per m²?

Sol. Length of room = 5.49m

$$\text{Width of room} = 3.87\text{m}$$

$$\text{Area of room} = \text{Length} \times \text{Width}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

343

$$= 5.49 \times 3.87$$

$$= 21.2463 \text{ sq. m}$$

Cost of carpeting one square m = Rs. 10.50

Cost of carpeting 21.2463 sq m = 10.50×21.2463

$$= 223.08615$$

Cost of carpeting = Rs. 223 Approx

4. The area of a rectangular rice field is 2.5 hectares and its sides are in the ratio 3 : 2. Find the perimeter of the field

Area of field = 2.5 hectares

Area of field = 25000 sq m

Length of field = $3x$ m

Width of field = $2x$ m and

$$\text{Area} = 3x \times 2x$$

$$\text{Area} = 6x^2 \text{ sq m}$$

from (i) and (ii)

$$6x^2 = 25000$$

$$x^2 = \frac{25000}{6}$$

$$x^2 = 4166.67 \text{ Taking square root}$$

$$x = 64.55$$

$$\text{Length } 3x = 193.65$$

$$\text{Width } 2x = 129.10$$

$$\text{Perimeter} = 2(3x + 2x) = 2(193.65 + 129.10)$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)


Pilot Super One "General Math" 11th

344

$$2(322.75)$$

$$= 645.50 \text{ m}$$

5. The area of a square playground is 4500 m². How long will a man take to cross it diagonally at the speed of 3 km per hour?



$$= 4500 \text{ sq m}$$

$$= \sqrt{4500} \text{ m}$$

$$\text{Length of diagonal of a square} = m\overline{BD} = \sqrt{(\sqrt{4500})^2 + (\sqrt{4500})^2}$$

$$\text{By Pythagoras theorem} = 4500 + 4500$$

$$= \sqrt{9000}$$

$$m\overline{BD} = 94.87 \text{ m}$$

$$\text{Time spent to cross 3 km distance} = 60 \text{ min}$$

$$\text{Time spent to cross 3000 km distance} = 60 \text{ min}$$

$$\text{Time spent to cross 94.87 m distance} = \frac{60}{3000} \times 94.87 \text{ min}$$

$$= \frac{94.87}{50}$$

$$= 1.8974 \text{ min}$$

$$= 1 \text{ min } 54 \text{ sec}$$


GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

=====

Pilot Super One "General Math" 10th

345

6. The diagonal of a square is 14cm Find its area.



Diagonal of square = 14cm

$$\begin{aligned}\text{Area of a square} &= \frac{14 \times 14}{2} \\ &= 14 \times 7 \\ &= 98 \text{ sq. cm}\end{aligned}$$

2nd method:

Suppose that side of a square = x cm

By Pythagoras theorem

$$x^2 + x^2 = (14)^2$$

$$2x^2 = 14 \times 14$$

$$x^2 = \frac{14 \times 14}{2}$$

$$= 14 \times 7 = 98$$

and x^2 is area of a square.

Thus, Area of square = 98 sq.cm

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

346

7. Find the area of a triangle whose sides are.

(i) 120cm, 150cm and 200cm

(ii) 50dm, 78dm and 112dm

Sol: $a = 200$ here

$$b = 150$$

$$c = 120$$

$$S = \frac{a+b+c}{2}$$

$$S = \frac{200+150+120}{2}$$

$$S = \frac{470}{2}$$

$$S = 235$$

$$S - a = 235 - 200 = 35$$

$$S - b = 235 - 150 = 85$$

$$S - c = 235 - 120 = 115$$

$$\text{Area of } \Delta A = \sqrt{S(S-a)(S-b)(S-c)}$$

$$A = \sqrt{235 \times 35 \times 85 \times 115}$$

$$= \sqrt{5 \times 47 \times 5 \times 7 \times 5 \times 17 \times 5 \times 23}$$

$$= \sqrt{5^2 \times 47 \times 7 \times 5^2 \times 17 \times 23}$$

$$= 5 \times 5 \sqrt{47 \times 7 \times 17 \times 23}$$

$$= 25 \sqrt{128639}$$

$$= 25 \times 358.66$$

$$= 8967 \text{ sq. cm approx}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

=====

Pilot Super One "General Math" 10th

347

(ii) $a = 112$ dm here

$$b = 78 \text{ dm}$$

$$c = 50 \text{ dm}$$

$$S = \frac{a+b+c}{2}$$

$$= \frac{112+78+50}{2}$$

$$= \frac{240}{2}$$

$$S = 120$$

$$\text{Area (A)} = A = \sqrt{S(S-a)(S-b)(S-c)}$$

$$= \sqrt{120(120-112)(120-78)(120-50)}$$

$$= \sqrt{120 \times 8 \times 42 \times 70}$$

$$A = \sqrt{3 \times 5 \times 8 \times 8 \times 7 \times 2 \times 3 \times 7 \times 5 \times 2}$$

$$= \sqrt{3^2 \times 5^2 \times 8^2 \times 7^2 \times 2^2}$$

$$= 3 \times 5 \times 8 \times 7 \times 2$$

$$= 1680 \text{ sq. dm}$$

8. The perimeter of a triangular field is 540m and its sides are in the ratio 25 : 17 : 12. Find the area of triangle.

Hint: Let the sides be $25x$, $17x$, $12x$ meters.

$$\text{Then } 25x + 17x + 12x = 540 \Rightarrow x = 10$$

Sol: Let sides are $25x$, $17x$, $12x$

$$\text{perimeter} = 25x + 17x + 12x$$

$$= 54x$$

$$\text{perimeter} = 540 \text{ m} \quad \text{but}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th 348

then $540 = 54x$

$x = 10$

sides $25 \times 10, 17 \times 10, 12 \times 10$

$= 250, 170, 120 \text{ m}$

$$S = \frac{a+b+c}{2}$$

$$= \frac{250+170+120}{2}$$

$$= \frac{540}{2} = 270$$

$S - a = 270 - 250 = 20$

$S - b = 270 - 170 = 100$

$S - c = 270 - 120 = 150$

$$\text{Area (A)} = A = \sqrt{S(S-a)(S-b)(S-c)}$$

$$= \sqrt{270 \times 20 \times 100 \times 150}$$

$$= \sqrt{27 \times 10 \times 2 \times 10 \times 2 \times 50 \times 3 \times 50}$$

$$= \sqrt{10^2 \times 2^2 \times 50^2 \times 3 \times 27}$$

$$= 10 \times 2 \times 50 \times \sqrt{3 \times 27}$$

$$= 1000 \times \sqrt{3 \times 3 \times 3 \times 3}$$

$$= 1000 \times 3 \times 3$$

$$= 9000 \text{ sq. meter}$$

9. Find the area of a parallelogram if its two adjacent sides are 12cm and 14cm and diagonal is 18cm.

Hints:

Let ABCD is a//gm in which

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)


Pilot Super One "General Math" 10th

349

Find area of ΔABC .

Area of $\parallel^m = 2$ (Area of ΔABC)

Sol: Find area of ΔACB



$$a = 18 \text{ cm}$$

$$b = 14 \text{ cm}$$

$$c = 12 \text{ cm}$$

$$S = \frac{a+b+c}{2}$$

$$S = \frac{18+14+12}{2}$$

$$S = \frac{44}{2}$$

$$S = 22$$

$$S - a = 22 - 18 = 4$$

$$S - b = 22 - 14 = 8$$

$$S - c = 22 - 12 = 10$$

$$\text{Area of } \Delta ABC = A = \sqrt{S(S-a)(S-b)(S-c)}$$

$$= \sqrt{22 \times 4 \times 8 \times 10}$$

$$= \sqrt{11 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 5}$$

$$= \sqrt{11 \times 2^2 \times 2^2 \times 2^2 \times 2 \times 5}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

350

$$= 2 \times 2 \times 2 \sqrt{110}$$

$$= 8 \sqrt{110} \text{ sq. cm} \dots\dots\dots (i)$$

Area of ΔACD = Area of ΔABC and

$$= 8 \sqrt{110} \text{ sq. cm} \dots\dots\dots (ii)$$

Area of $\square ABCD$ = (i) + (ii)

$$= 8 \sqrt{110} + 8 \sqrt{110}$$

$$= (8+8) \sqrt{110}$$


$$= 16 \sqrt{110} \text{ sq. cm}$$

10. Find the area of the following washers whose external and internal diameters are:

(i) 15cm and 13cm

(ii) 1.2m and 0.9m

(iii) 40mm and 33mm.



$$m AB = 15 \text{ cm}$$

$$m PQ = 13 \text{ cm}$$

Sol: External diameter = $\frac{15}{2} \text{ cm}$

Internal diameter = $\frac{13}{2} \text{ cm}$

$$\text{Total area} = \pi^2$$

$$= \pi \left(\frac{15}{2} \right)^2$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pink Super Line "General Math" 11ph

351

$$\text{Internal area} = \pi r^2$$

$$= \pi \left(\frac{13}{2} \right)^2$$

$$\text{Area of washers} = \pi \left(\frac{15}{2} \right)^2 - \pi \left(\frac{13}{2} \right)^2$$

$$= \pi \left[\left(\frac{15}{2} \right)^2 - \left(\frac{13}{2} \right)^2 \right]$$

$$= \frac{22}{7} \left[\frac{225}{4} - \frac{169}{4} \right]$$

$$= \frac{22}{7} \left[\frac{225 - 169}{4} \right]$$

$$= \frac{22}{7} \left(\frac{56}{4} \right)$$

$$44 \text{ sq. cm}$$

$$\text{Sol: External diameter} = \frac{1.2}{2} = 6 \text{ m} = \frac{6}{10} \text{ m}$$

$$\text{Internal diameter} = \frac{0.9}{2} = \frac{9}{10 \times 2}$$

$$= \frac{9}{20} \text{ m}$$

$$\text{Total area} = \pi r^2$$

$$= \pi \left(\frac{6}{10} \right)^2$$

$$\text{Internal area} = \pi r^2$$

$$= \pi \left(\frac{9}{20} \right)^2$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

352

$$\begin{aligned}
 \text{Area of washers} &= \pi \left[\left(\frac{6}{10} \right)^2 - \left(\frac{9}{20} \right)^2 \right] \\
 &= \pi \left[\left(\frac{6}{10} \right)^2 - \left(\frac{9}{20} \right)^2 \right] \\
 &= \frac{22}{7} \left[\frac{36}{100} - \frac{81}{400} \right] \\
 &= \frac{22}{7} \left[\frac{144 - 81}{400} \right] \\
 &= \frac{22}{7} \left(\frac{63}{400} \right) \\
 &= \frac{198}{400} \\
 &= 0.495 \text{ sq. mm}
 \end{aligned}$$

Sol:

$$\text{Diameter of big half circle} = \frac{40}{2} = 20 \text{ mm}$$

$$\text{Internal diameter} = \frac{33}{2} \text{ mm}$$

$$\begin{aligned}
 \text{Total area} &= \pi r^2 \\
 &= \pi (20)^2
 \end{aligned}$$

$$\begin{aligned}
 \text{Internal area} &= \pi r^2 \\
 &= \pi \left(\frac{33}{2} \right)^2
 \end{aligned}$$

$$\begin{aligned}
 \text{Area of washers} &= \pi (20)^2 - \pi \left(\frac{33}{2} \right)^2 \\
 &= \pi \left[(20)^2 - \left(\frac{33}{2} \right)^2 \right]
 \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)


Pilot Super One "General Math" 10th

353

$$\begin{aligned}
 &= \frac{22}{7} \left[400 - \frac{1089}{4} \right] \\
 &= \frac{22}{7} \left[\frac{1600 - 1089}{4} \right] \\
 &= \frac{22}{7} \left(\frac{511}{4} \right) \\
 &= \frac{11 \times 73}{2}
 \end{aligned}$$

$$= 401.5 \text{ sq. mm}$$

11. Find the area of the shaded region.



$$\text{Diameter of big half circle} = \frac{28}{2} = 14 \text{ m}$$

$$\begin{aligned}
 \text{Area of big half circle} &= \frac{1}{2} \pi r^2 \\
 &= \frac{1}{2} \pi (14)^2 \\
 &= \frac{1}{2} \times \frac{22}{7} \times 14^2 \times 14 \\
 &= 308 \text{ sq. m}
 \end{aligned}$$

$$\text{Diameter of every small half circle} = \frac{14}{2} = 7 \text{ m}$$

$$\text{Total area of both} = \frac{1}{2} \pi r^2$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

354

$$= \frac{1}{2} \times \frac{22}{7} \times (7)^2$$

$$= \frac{1}{2} \times \frac{22}{7} \times 7 \times 7$$

$$= 77 \text{ sq m}$$

$$\text{small circles} = 2 \times 77$$

$$= 154 \text{ sq m}$$

$$\text{Area of shaded region} = 308 - 154$$

$$= 154 \text{ sq m}$$

12. Find the area of an equilateral triangle whose side is 8m.

Sol $a = b = c = 8\text{m}$

$$S = \frac{a+b+c}{2} = \frac{8+8+8}{2} = \frac{24}{2} = 12$$

$$S - a = 12 - 8 = 4$$

$$S - b = 12 - 8 = 4$$

$$S - c = 12 - 8 = 4$$

$$\text{Area of } \triangle = A = \sqrt{S(S-a)(S-b)(S-c)}$$

$$= \sqrt{12 \times 4 \times 4 \times 4}$$

$$= \sqrt{3 \times 4 \times 4 \times 4 \times 4}$$

$$= 4 \times 4 \sqrt{3}$$

$$= 16\sqrt{3} \text{ sq. m}$$

13. The side of an equilateral triangle is 6cm. Find its area.

$$a = b = c = 6 \quad \text{here}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

355

$$S = \frac{a+b+c}{2} = \frac{6+6+6}{2} = \frac{18}{2} = 9$$

$$S - a = 9 - 6 = 3$$

$$S - b = 9 - 6 = 3$$

$$S - c = 9 - 6 = 3$$

$$\begin{aligned} \text{Area of } \Delta &= A = \sqrt{S(S-a)(S-b)(S-c)} \\ &= \sqrt{9 \times 3 \times 3 \times 3} \\ &= \sqrt{3 \times 3 \times 3 \times 3 \times 3} \\ &= 3 \times 3\sqrt{3} = 9\sqrt{3} \text{ sq. cm} \end{aligned}$$

14. Find the area of the right triangle with legs 12cm and 35cm.

Length of one side = 12 cm

Length of 2nd side = 35 cm

$$\begin{aligned} \text{Area of right angles} &= \frac{\text{base} \times \text{altitude}}{2} \\ &= \frac{35 \times 12}{2} \\ &= 35 \times 6 \\ &= 210 \text{ sq. cm} \end{aligned}$$

15. The base of a rectangle is three times its altitude. The area is 147cm². Find the dimensions of the rectangle.

Sol Let altitude = x

Length of base = 3x cm

$$\begin{aligned} \text{Area} &= (x)(3x) \\ &= 3x^2 \end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

356

$$\text{Area} = 147 \text{ sq. cm} \quad \text{but}$$

$$3x^2 = 147 \quad \text{therefore}$$

$$x^2 = \frac{147}{3}$$

$$x^2 = 49$$

$$\sqrt{x^2} = \sqrt{49} \quad \text{Taking square root}$$

$$x = 7$$

$$\text{altitude} = 7 \text{ cm}$$

$$\begin{aligned} \text{Base} &= 3x \\ &= 3 \times 7 \end{aligned}$$

$$\text{Base} = 21 \text{ cm}$$

16. Find the base of the parallelogram whose altitude is 18cm and whose area is 3m².

$$\text{Length of altitude} = 18 \text{ cm}$$

$$\text{Area} = 3 \text{ sq.m}$$

$$\text{Area} = 3 \times 100 \times 100 \text{ sq.m}$$

$$\begin{aligned} \text{Length of base} &= \frac{\text{area}}{\text{altitude}} \\ &= \frac{3 \times 100 \times 100 \text{ m}}{18} \\ &= \frac{5000}{3} \end{aligned}$$

$$\text{Length of base} = 1666.67 \text{ cm}$$

17. The area of a parallelogram is 144cm². Find the altitude if the base is 2m long.

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

357

Area of a parallelogram = 144 sq. m

Length of base = 2 cm

$$\begin{aligned}\text{Length of altitude} &= \frac{\text{area}}{\text{base}} \\ &= \frac{144}{2}\end{aligned}$$

Length of altitude = 72 cm

18. Find the area of the rectangle 2m long and 18cm wide.

Length of a rectangle = 2m

= 200 cm

Width = 18 cm

Area = Length × Width

= 200 × 18

= 3600 sq.cm

19. The area of an equilateral triangle is $4\sqrt{3}\text{cm}^2$. Find the length of a side.

Sol: If side of an any equilateral triangle are "a" units then its

area will be $\frac{\sqrt{3} a^2}{4}$.

• therefore $\frac{\sqrt{3} a^2}{4} = 4\sqrt{3}$

$$a^2 = (4\sqrt{3}) \left(\frac{4}{\sqrt{3}} \right)$$

$$a^2 = 4 \times 4$$

$$\sqrt{a^2} = \sqrt{4 \times 4} \quad \text{Taking square root}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

358

$$a = 4 \text{ cm}$$

$$\begin{aligned} \text{Volume of a cube} &= V = l \times l \times l \\ &= (l)^3 \end{aligned}$$

Volume of a cuboid of length l , breadth b and height h is

$$\text{Volume of cuboid} = V = l \times b \times h$$

Volume of a cuboid of length l , breadth b and height h is

$$\text{Volume of cylinder} = \pi r^2 h$$

where ' r ' is radius of the base and ' h ' is height.

$$\text{Volume of cylinder} = \frac{1}{3} \pi r^2 h$$

where ' r ' is radius of base and ' h ' is height.

$$\text{Volume of sphere} = \frac{4}{3} \pi r^3$$

where ' r ' is radius of sphere.

Remember that:

$$1. \quad \text{As } 1 \text{ cm} = 10 \text{ mm},$$

$$\text{therefore, } 1 \text{ cm}^3 = 10 \times 10 \times 10 \text{ mm}^3$$

$$1 \text{ cm}^3 = 1000 \text{ mm}^3$$

$$2. \quad 1 \text{ m}^3 = 100 \times 100 \times 100 \text{ cm}^3$$

$$= 1000000 \text{ cm}^3$$

$$\boxed{1 \text{ m}^3 = 10^6 \text{ cm}^3}$$

$$\text{also } 1 \text{ m}^3 = 1000 \times 1000 \times 1000 \text{ mm}^3$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

359

$$1m^3 = 10^9 mm^3$$

- 3- For measurement of volumes of liquids, we use the terms
liters (*l*) and milliliters (*ml*).

$$1cm^3 = 1ml$$

$$1000cm^3 = 1l$$

$$\text{and } 1m^3 = 1000000 cm^3 = 1000 l$$

$$1m^3 = 1kl \text{ (1 kiloliter)}$$

$$10^3 \text{ liters} = 1kl$$

Find the Volume of the Solids

1. A cube of a side 4cm

Sol: A cube of a side = 4

$$\text{Volume of cube} = (l)^3$$

$$= (4)^3$$

$$= 64 \text{ cubic cm}$$

2. A cube whose total area is $96cm^2$.

$$\text{Total area of cube} = 96$$

$$\text{areas of cube} = 6$$

$$\text{area of 1 face} = \frac{96}{6}$$

$$= 16 \text{ sq. cm}$$

$$\text{Length of edge} = \sqrt{16}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

=====

Pilot Super One "General Math" 10th

360

$$= 4 \text{ cm}$$

$$\text{volume of cube} = (l)^3$$

$$= (4)^3$$

$$= 64 \text{ cm}^3$$

3. A rectangular box with length 4m breadth 3m and height 2m.

$$\text{length of rectangular} = 4 \text{ m}$$

$$\text{Breadth of rectangular} = 3 \text{ m}$$

$$\text{Height of rectangular} = 2 \text{ m}$$

$$\text{volume of cuboid} = l \times b \times h$$

$$= 4 \times 3 \times 2$$

$$= 24 \text{ (m)}^3$$

4. Right cylinder, with radius of base 4cm, altitude 10cm, use

$$\pi = \frac{22}{7}$$

Sol. Radius of base = (r) = 4 cm

$$\text{altitude} = (h) = 10 \text{ cm}$$

$$\text{volume of cylinder} = \pi r^2 h$$

$$= \frac{22}{7} \times (4)^2 (10)$$

$$= \frac{22}{7} \times 16 \times 10$$

$$= \frac{3520}{7}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

361

$$= 502.86 \text{ cm}^3$$

5. Circular cone, with radius of base 3cm, altitude 10cm.

$$\text{Radius of circular base} = r = 3 \text{ cm}$$

$$\text{Height of altitude} = h = 10 \text{ cm}$$

$$\begin{aligned}\text{Volume of circular cone} &= \frac{1}{3} \pi r^2 h \\ &= \frac{1}{3} \left(\frac{22}{7} \right) (3)^2 (10) \\ &= \frac{1}{3} \times \frac{22}{7} \times 3 \times 3 \times 10 \\ &= \frac{660}{7} \\ &= 94.3 \text{ cm}^3\end{aligned}$$

6. Sphere, with radius 3cm.

$$\text{Radius of sphere} = r = 3 \text{ cm}$$

$$\begin{aligned}\text{Volume of sphere} &= \frac{4}{3} \pi r^3 \\ &= \frac{4}{3} \times \frac{22}{7} \times (3)^3 \\ &= \frac{4}{3} \times \frac{22}{7} \times 3 \times 3 \times 3 \\ &= \frac{4 \times 22 \times 3 \times 3}{7} \\ \text{Volume of sphere} &= \frac{792}{7} \\ &= 113.14 \text{ cm}^3\end{aligned}$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pilot Super One "General Math" 10th

362

7. Right circular cylinder, with circumferences of base 4cm, altitude 1m

Sol: Circumference of base = 4cm

Circumference of base = $2\pi r$

then $2\pi r = 4$

$$r = \frac{4}{2\pi}$$

$$= \frac{2}{\pi} \text{ cm} \quad (i)$$

Length of cylinder (h) = 1m

$$= 100 \text{ cm}$$

Volume of cylinder = $\pi r^2 \times h$

$$\text{from (i)} \quad = \pi \left(\frac{2}{\pi} \right)^2 (100)$$

$$= \pi \times \frac{2 \times 2 \times 100}{\pi \times \pi}$$

$$= \frac{7 \times 2 \times 2 \times 100}{2 \times 2}$$

$$= \frac{1400}{11}$$

$$= 127.3 \text{ cm}^3 \quad \text{approx}$$

8. Cone with altitude 9cm, radius of base 6cm.

Sol: Cone with altitude = h = 9 cm

Cone with radius of base = r = 6 cm

$$\text{Volume of cone} = \frac{1}{3} \pi r^2 \times h$$

GENERAL MATHEMATICS FOR 10TH CLASS (UNIT # 9)

Pink Super One "General Math" 10th

363

$$= \frac{1}{3} \left(\frac{22}{7} \right) (6)^2 \times 9$$

$$= \frac{1}{3} \times \frac{22}{7} \times 36 \times 9$$

$$= \frac{22 \times 12 \times 9}{7}$$

$$= \frac{2376}{7}$$

$$= 339.4 \text{ cm}^3$$

www.downloadclassnotes.com

